

3-3-2006

# Evidence for the Exclusive Decay $B_c^\pm \rightarrow J/\psi\pi^\pm$ and Measurement of the Mass of the $B_c^\pm$ Meson

A. Abulencia

*University of Illinois, Urbana, Illinois 61801, USA*

Kenneth Bloom

*University of Nebraska - Lincoln, kenbloom@unl.edu*

CDF Collaboration

Follow this and additional works at: <http://digitalcommons.unl.edu/physicsbloom>



Part of the [Physics Commons](#)

---

Abulencia, A.; Bloom, Kenneth; and Collaboration, CDF, "Evidence for the Exclusive Decay  $B_c^\pm \rightarrow J/\psi\pi^\pm$  and Measurement of the Mass of the  $B_c^\pm$  Meson" (2006). *Kenneth Bloom Publications*. Paper 192.

<http://digitalcommons.unl.edu/physicsbloom/192>

This Article is brought to you for free and open access by the Research Papers in Physics and Astronomy at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Kenneth Bloom Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

**Evidence for the Exclusive Decay  $B_c^\pm \rightarrow J/\psi\pi^\pm$  and Measurement of the Mass of the  $B_c^\pm$  Meson**

A. Abulencia,<sup>23</sup> D. Acosta,<sup>17</sup> J. Adelman,<sup>13</sup> T. Affolder,<sup>10</sup> T. Akimoto,<sup>53</sup> M. G. Albrow,<sup>16</sup> D. Ambrose,<sup>16</sup> S. Amerio,<sup>42</sup> D. Amidei,<sup>33</sup> A. Anastassov,<sup>50</sup> K. Anikeev,<sup>16</sup> A. Annovi,<sup>44</sup> J. Antos,<sup>1</sup> M. Aoki,<sup>53</sup> G. Apollinari,<sup>16</sup> J.-F. Arguin,<sup>32</sup> T. Arisawa,<sup>55</sup> A. Artikov,<sup>14</sup> W. Ashmanskas,<sup>16</sup> A. Attal,<sup>8</sup> F. Azfar,<sup>41</sup> P. Azzi-Bacchetta,<sup>42</sup> P. Azzurri,<sup>44</sup> N. Bacchetta,<sup>42</sup> H. Bachocou,<sup>28</sup> W. Badgett,<sup>16</sup> A. Barbaro-Galtieri,<sup>28</sup> V. E. Barnes,<sup>46</sup> B. A. Barnett,<sup>24</sup> S. Baroiant,<sup>7</sup> V. Bartsch,<sup>30</sup> G. Bauer,<sup>31</sup> F. Bedeschi,<sup>44</sup> S. Behari,<sup>24</sup> S. Belforte,<sup>52</sup> G. Bellettini,<sup>44</sup> J. Bellinger,<sup>57</sup> A. Belloni,<sup>31</sup> E. Ben-Haim,<sup>16</sup> D. Benjamin,<sup>15</sup> A. Beretvas,<sup>16</sup> J. Beringer,<sup>28</sup> T. Berry,<sup>29</sup> A. Bhatti,<sup>48</sup> M. Binkley,<sup>16</sup> D. Bisello,<sup>42</sup> M. Bishai,<sup>16</sup> R. E. Blair,<sup>2</sup> C. Blocker,<sup>6</sup> K. Bloom,<sup>33</sup> B. Blumenfeld,<sup>24</sup> A. Bocci,<sup>48</sup> A. Bodek,<sup>47</sup> V. Boisvert,<sup>47</sup> G. Bolla,<sup>46</sup> A. Bolshov,<sup>31</sup> D. Bortoletto,<sup>46</sup> J. Boudreau,<sup>45</sup> S. Bourov,<sup>16</sup> A. Boveia,<sup>10</sup> B. Brau,<sup>10</sup> C. Bromberg,<sup>34</sup> E. Brubaker,<sup>13</sup> J. Budagov,<sup>14</sup> H. S. Budd,<sup>47</sup> S. Budd,<sup>23</sup> K. Burkett,<sup>16</sup> G. Busetto,<sup>42</sup> P. Bussey,<sup>20</sup> K. L. Byrum,<sup>2</sup> S. Cabrera,<sup>15</sup> M. Campanelli,<sup>19</sup> M. Campbell,<sup>33</sup> F. Canelli,<sup>8</sup> A. Canepa,<sup>46</sup> D. Carlsmith,<sup>57</sup> R. Carosi,<sup>44</sup> S. Carron,<sup>15</sup> M. Casarsa,<sup>52</sup> A. Castro,<sup>5</sup> P. Catastini,<sup>44</sup> D. Cauz,<sup>52</sup> M. Cavalli-Sforza,<sup>3</sup> A. Cerri,<sup>28</sup> L. Cerrito,<sup>41</sup> S. H. Chang,<sup>27</sup> J. Chapman,<sup>33</sup> Y. C. Chen,<sup>1</sup> M. Chertok,<sup>7</sup> G. Chiarelli,<sup>44</sup> G. Chlachidze,<sup>14</sup> F. Chlebana,<sup>16</sup> I. Cho,<sup>27</sup> K. Cho,<sup>27</sup> D. Chokheli,<sup>14</sup> J. P. Chou,<sup>21</sup> P. H. Chu,<sup>23</sup> S. H. Chuang,<sup>57</sup> K. Chung,<sup>12</sup> W. H. Chung,<sup>57</sup> Y. S. Chung,<sup>47</sup> M. Cijliak,<sup>44</sup> C. I. Ciobanu,<sup>23</sup> M. A. Ciocci,<sup>44</sup> A. Clark,<sup>19</sup> D. Clark,<sup>6</sup> M. Coca,<sup>15</sup> A. Connolly,<sup>28</sup> M. Convery,<sup>48</sup> J. Conway,<sup>7</sup> B. Cooper,<sup>30</sup> K. Copic,<sup>33</sup> M. Cordelli,<sup>18</sup> G. Cortiana,<sup>42</sup> J. Cranshaw,<sup>2</sup> A. Cruz,<sup>17</sup> J. Cuevas,<sup>11</sup> R. Culbertson,<sup>16</sup> D. Cyr,<sup>57</sup> S. Da Ronco,<sup>42</sup> S. D'Auria,<sup>20</sup> M. D'Onofrio,<sup>19</sup> D. Dagenhart,<sup>6</sup> P. de Barbaro,<sup>47</sup> S. De Cecco,<sup>49</sup> A. Deisher,<sup>28</sup> G. De Lentdecker,<sup>47</sup> M. Dell'Orso,<sup>44</sup> S. Demers,<sup>47</sup> L. Demortier,<sup>48</sup> J. Deng,<sup>15</sup> M. Deninno,<sup>5</sup> D. De Pedis,<sup>49</sup> P. F. Derwent,<sup>16</sup> T. Devlin,<sup>50</sup> C. Dionisi,<sup>49</sup> J. R. Dittmann,<sup>4</sup> P. DiTuro,<sup>50</sup> C. Dörr,<sup>25</sup> A. Dominguez,<sup>28</sup> S. Donati,<sup>44</sup> M. Donega,<sup>19</sup> P. Dong,<sup>8</sup> J. Donini,<sup>42</sup> T. Dorigo,<sup>42</sup> S. Dube,<sup>50</sup> K. Ebina,<sup>55</sup> J. Efron,<sup>38</sup> J. Ehlers,<sup>19</sup> R. Erbacher,<sup>7</sup> D. Errede,<sup>23</sup> S. Errede,<sup>23</sup> R. Eusebi,<sup>47</sup> H. C. Fang,<sup>28</sup> S. Farrington,<sup>29</sup> I. Fedorko,<sup>44</sup> W. T. Fedorko,<sup>13</sup> R. G. Feild,<sup>58</sup> M. Feindt,<sup>25</sup> J. P. Fernandez,<sup>46</sup> R. Field,<sup>17</sup> G. Flanagan,<sup>34</sup> L. R. Flores-Castillo,<sup>45</sup> A. Foland,<sup>21</sup> S. Forrester,<sup>7</sup> G. W. Foster,<sup>16</sup> M. Franklin,<sup>21</sup> J. C. Freeman,<sup>28</sup> Y. Fujii,<sup>26</sup> I. Furic,<sup>13</sup> A. Gajjar,<sup>29</sup> M. Gallinaro,<sup>48</sup> J. Galyardt,<sup>12</sup> J. E. Garcia,<sup>44</sup> M. Garcia Sciveres,<sup>28</sup> A. F. Garfinkel,<sup>46</sup> C. Gay,<sup>58</sup> H. Gerberich,<sup>23</sup> E. Gerchtein,<sup>12</sup> D. Gerdes,<sup>33</sup> S. Giagu,<sup>49</sup> P. Giannetti,<sup>44</sup> A. Gibson,<sup>28</sup> K. Gibson,<sup>12</sup> C. Ginsburg,<sup>16</sup> K. Giolo,<sup>46</sup> M. Giordani,<sup>52</sup> M. Giunta,<sup>44</sup> G. Giurgiu,<sup>12</sup> V. Glagolev,<sup>14</sup> D. Glenzinski,<sup>16</sup> M. Gold,<sup>36</sup> N. Goldschmidt,<sup>33</sup> J. Goldstein,<sup>41</sup> G. Gomez,<sup>11</sup> G. Gomez-Ceballos,<sup>11</sup> M. Goncharov,<sup>51</sup> O. González,<sup>46</sup> I. Gorelov,<sup>36</sup> A. T. Goshaw,<sup>15</sup> Y. Gotra,<sup>45</sup> K. Goulianos,<sup>48</sup> A. Gresele,<sup>42</sup> M. Griffiths,<sup>29</sup> S. Grinstein,<sup>21</sup> C. Grosso-Pilcher,<sup>13</sup> U. Grundler,<sup>23</sup> J. Guimaraes da Costa,<sup>21</sup> C. Haber,<sup>28</sup> S. R. Hahn,<sup>16</sup> K. Hahn,<sup>43</sup> E. Halkiadakis,<sup>47</sup> A. Hamilton,<sup>32</sup> B.-Y. Han,<sup>47</sup> R. Handler,<sup>57</sup> F. Happacher,<sup>18</sup> K. Hara,<sup>53</sup> M. Hare,<sup>54</sup> S. Harper,<sup>41</sup> R. F. Harr,<sup>56</sup> R. M. Harris,<sup>16</sup> K. Hatakeyama,<sup>48</sup> J. Hauser,<sup>8</sup> C. Hays,<sup>15</sup> H. Hayward,<sup>29</sup> A. Heijboer,<sup>43</sup> B. Heinemann,<sup>29</sup> J. Heinrich,<sup>43</sup> M. Hennecke,<sup>25</sup> M. Herndon,<sup>57</sup> J. Heuser,<sup>25</sup> D. Hidas,<sup>15</sup> C. S. Hill,<sup>10</sup> D. Hirschbuehl,<sup>25</sup> A. Hocker,<sup>16</sup> A. Holloway,<sup>21</sup> S. Hou,<sup>1</sup> M. Houlden,<sup>29</sup> S.-C. Hsu,<sup>9</sup> B. T. Huffman,<sup>41</sup> R. E. Hughes,<sup>38</sup> J. Huston,<sup>34</sup> K. Ikado,<sup>55</sup> J. Incandela,<sup>10</sup> G. Introzzi,<sup>44</sup> M. Iori,<sup>49</sup> Y. Ishizawa,<sup>53</sup> A. Ivanov,<sup>7</sup> B. Iyutin,<sup>31</sup> E. James,<sup>16</sup> D. Jang,<sup>50</sup> B. Jayatilaka,<sup>33</sup> D. Jeans,<sup>49</sup> H. Jensen,<sup>16</sup> E. J. Jeon,<sup>27</sup> M. Jones,<sup>46</sup> K. K. Joo,<sup>27</sup> S. Y. Jun,<sup>12</sup> T. R. Junk,<sup>23</sup> T. Kamon,<sup>51</sup> J. Kang,<sup>33</sup> M. Karagoz-Unel,<sup>37</sup> P. E. Karchin,<sup>56</sup> Y. Kato,<sup>40</sup> Y. Kemp,<sup>25</sup> R. Kephart,<sup>16</sup> U. Kerzel,<sup>25</sup> V. Khotilovich,<sup>51</sup> B. Kilminster,<sup>38</sup> D. H. Kim,<sup>27</sup> H. S. Kim,<sup>27</sup> J. E. Kim,<sup>27</sup> M. J. Kim,<sup>12</sup> M. S. Kim,<sup>27</sup> S. B. Kim,<sup>27</sup> S. H. Kim,<sup>53</sup> Y. K. Kim,<sup>13</sup> M. Kirby,<sup>15</sup> L. Kirsch,<sup>6</sup> S. Klimentenko,<sup>17</sup> M. Klute,<sup>31</sup> B. Knuteson,<sup>31</sup> B. R. Ko,<sup>15</sup> H. Kobayashi,<sup>53</sup> K. Kondo,<sup>55</sup> D. J. Kong,<sup>27</sup> J. Konigsberg,<sup>17</sup> K. Kordas,<sup>18</sup> A. Korytov,<sup>17</sup> A. V. Kotwal,<sup>15</sup> A. Kovalev,<sup>43</sup> J. Kraus,<sup>23</sup> I. Kravchenko,<sup>31</sup> M. Kreps,<sup>25</sup> A. Kreymer,<sup>16</sup> J. Kroll,<sup>43</sup> N. Krumnack,<sup>4</sup> M. Kruse,<sup>15</sup> V. Krutelyov,<sup>51</sup> S. E. Kuhlmann,<sup>2</sup> Y. Kusakabe,<sup>55</sup> S. Kwang,<sup>13</sup> A. T. Laasanen,<sup>46</sup> S. Lai,<sup>32</sup> S. Lami,<sup>44</sup> S. Lammel,<sup>16</sup> M. Lancaster,<sup>30</sup> R. L. Lander,<sup>7</sup> K. Lannon,<sup>38</sup> A. Lath,<sup>50</sup> G. Latino,<sup>44</sup> I. Lazzizzera,<sup>42</sup> C. Lecci,<sup>25</sup> T. LeCompte,<sup>2</sup> J. Lee,<sup>47</sup> J. Lee,<sup>27</sup> S. W. Lee,<sup>51</sup> R. Lefèvre,<sup>3</sup> N. Leonardo,<sup>31</sup> S. Leone,<sup>44</sup> S. Levy,<sup>13</sup> J. D. Lewis,<sup>16</sup> K. Li,<sup>58</sup> C. Lin,<sup>58</sup> C. S. Lin,<sup>16</sup> M. Lindgren,<sup>16</sup> E. Lipeles,<sup>9</sup> T. M. Liss,<sup>23</sup> A. Lister,<sup>19</sup> D. O. Litvintsev,<sup>16</sup> T. Liu,<sup>16</sup> Y. Liu,<sup>19</sup> N. S. Lockyer,<sup>43</sup> A. Loginov,<sup>35</sup> M. Loreti,<sup>42</sup> P. Loverre,<sup>49</sup> R.-S. Lu,<sup>1</sup> D. Lucchesi,<sup>42</sup> P. Lujan,<sup>28</sup> P. Lukens,<sup>16</sup> G. Lungu,<sup>17</sup> L. Lyons,<sup>41</sup> J. Lys,<sup>28</sup> R. Lysak,<sup>1</sup> E. Lytken,<sup>46</sup> P. Mack,<sup>25</sup> D. MacQueen,<sup>32</sup> R. Madrak,<sup>16</sup> K. Maeshima,<sup>16</sup> P. Maksimovic,<sup>24</sup> G. Manca,<sup>29</sup> F. Margaroli,<sup>5</sup> R. Marginean,<sup>16</sup> C. Marino,<sup>23</sup> A. Martin,<sup>58</sup> M. Martin,<sup>24</sup> V. Martin,<sup>37</sup> M. Martínez,<sup>3</sup> T. Maruyama,<sup>53</sup> H. Matsunaga,<sup>53</sup> M. E. Mattson,<sup>56</sup> R. Mazini,<sup>32</sup> P. Mazzanti,<sup>5</sup> K. S. McFarland,<sup>47</sup> D. McGivern,<sup>30</sup> P. McIntyre,<sup>51</sup> P. McNamara,<sup>50</sup> R. McNulty,<sup>29</sup> A. Mehta,<sup>29</sup> S. Menzemer,<sup>31</sup> A. Menzione,<sup>44</sup> P. Merkel,<sup>46</sup> C. Mesropian,<sup>48</sup> A. Messina,<sup>49</sup> M. von der Mey,<sup>8</sup> T. Miao,<sup>16</sup> N. Miladinovic,<sup>6</sup> J. Miles,<sup>31</sup> R. Miller,<sup>34</sup> J. S. Miller,<sup>33</sup> C. Mills,<sup>10</sup> M. Milnik,<sup>25</sup> R. Miquel,<sup>28</sup> S. Miscetti,<sup>18</sup> G. Mitselmakher,<sup>17</sup> A. Miyamoto,<sup>26</sup> N. Moggi,<sup>5</sup> B. Mohr,<sup>8</sup> R. Moore,<sup>16</sup> M. Morello,<sup>44</sup> P. Movilla Fernandez,<sup>28</sup> J. Mülmenstädt,<sup>28</sup> A. Mukherjee,<sup>16</sup> M. Mulhearn,<sup>31</sup> Th. Muller,<sup>25</sup> R. Mumford,<sup>24</sup> P. Murat,<sup>16</sup> J. Nachtman,<sup>16</sup> S. Nahn,<sup>58</sup> I. Nakano,<sup>39</sup>

A. Napier,<sup>54</sup> D. Naumov,<sup>36</sup> V. Necula,<sup>17</sup> C. Neu,<sup>43</sup> M. S. Neubauer,<sup>9</sup> L. Nicolas,<sup>16,20</sup> J. Nielsen,<sup>28</sup> T. Nigmanov,<sup>45</sup>  
 L. Nodulman,<sup>2</sup> O. Norriella,<sup>3</sup> T. Ogawa,<sup>55</sup> S. H. Oh,<sup>15</sup> Y. D. Oh,<sup>27</sup> T. Okusawa,<sup>40</sup> R. Oldeman,<sup>29</sup> R. Orava,<sup>22</sup>  
 K. Osterberg,<sup>22</sup> C. Pagliarone,<sup>44</sup> E. Palencia,<sup>11</sup> R. Paoletti,<sup>44</sup> V. Papadimitriou,<sup>16</sup> A. Papikononou,<sup>25</sup> A. A. Paramonov,<sup>13</sup>  
 B. Parks,<sup>38</sup> S. Pashapour,<sup>32</sup> J. Patrick,<sup>16</sup> G. Pauletta,<sup>52</sup> M. Paulini,<sup>12</sup> C. Paus,<sup>31</sup> D. E. Pellett,<sup>7</sup> A. Penzo,<sup>52</sup> T. J. Phillips,<sup>15</sup>  
 G. Piacentino,<sup>44</sup> J. Piedra,<sup>11</sup> K. Pitts,<sup>23</sup> C. Plager,<sup>8</sup> L. Pondrom,<sup>57</sup> G. Pope,<sup>45</sup> X. Portell,<sup>3</sup> O. Poukhov,<sup>14</sup> N. Pounder,<sup>41</sup>  
 F. Prakoshyn,<sup>14</sup> A. Pronko,<sup>16</sup> J. Proudfoot,<sup>2</sup> F. Ptohos,<sup>18</sup> G. Punzi,<sup>44</sup> J. Pursley,<sup>24</sup> J. Rademacker,<sup>41</sup> A. Rahaman,<sup>45</sup>  
 A. Rakitin,<sup>31</sup> S. Rappoccio,<sup>21</sup> F. Ratnikov,<sup>50</sup> B. Reiser,<sup>16</sup> V. Rekovic,<sup>36</sup> N. van Remortel,<sup>22</sup> P. Renton,<sup>41</sup> M. Rescigno,<sup>49</sup>  
 S. Richter,<sup>25</sup> F. Rimondi,<sup>5</sup> K. Rinnert,<sup>25</sup> L. Ristori,<sup>44</sup> W. J. Robertson,<sup>15</sup> A. Robson,<sup>20</sup> T. Rodrigo,<sup>11</sup> E. Rogers,<sup>23</sup> S. Rolli,<sup>54</sup>  
 R. Roser,<sup>16</sup> M. Rossi,<sup>52</sup> R. Rossin,<sup>17</sup> C. Rott,<sup>46</sup> A. Ruiz,<sup>11</sup> J. Russ,<sup>12</sup> V. Rusu,<sup>13</sup> D. Ryan,<sup>54</sup> H. Saarikko,<sup>22</sup> S. Sabik,<sup>32</sup>  
 A. Safonov,<sup>7</sup> W. K. Sakumoto,<sup>47</sup> G. Salamanna,<sup>49</sup> O. Salto,<sup>3</sup> D. Saltzberg,<sup>8</sup> C. Sanchez,<sup>3</sup> L. Santi,<sup>52</sup> S. Sarkar,<sup>49</sup> K. Sato,<sup>53</sup>  
 P. Savard,<sup>32</sup> A. Savoy-Navarro,<sup>16</sup> T. Scheidle,<sup>25</sup> P. Schlabach,<sup>16</sup> E. E. Schmidt,<sup>16</sup> M. P. Schmidt,<sup>58</sup> M. Schmitt,<sup>37</sup>  
 T. Schwarz,<sup>33</sup> L. Scodellaro,<sup>11</sup> A. L. Scott,<sup>10</sup> A. Scribano,<sup>44</sup> F. Scuri,<sup>44</sup> A. Sedov,<sup>46</sup> S. Seidel,<sup>36</sup> Y. Seiya,<sup>40</sup> A. Semenov,<sup>14</sup>  
 F. Semeria,<sup>5</sup> L. Sexton-Kennedy,<sup>16</sup> I. Sfiligoi,<sup>18</sup> M. D. Shapiro,<sup>28</sup> T. Shears,<sup>29</sup> P. F. Shepard,<sup>45</sup> D. Sherman,<sup>21</sup>  
 M. Shimojima,<sup>53</sup> M. Shochet,<sup>13</sup> Y. Shon,<sup>57</sup> I. Shreyber,<sup>35</sup> A. Sidoti,<sup>44</sup> A. Sill,<sup>16</sup> P. Sinervo,<sup>32</sup> A. Sisakyan,<sup>14</sup> J. Sjolín,<sup>41</sup>  
 A. Skiba,<sup>25</sup> A. J. Slaughter,<sup>16</sup> K. Sliwa,<sup>54</sup> D. Smirnov,<sup>36</sup> J. R. Smith,<sup>7</sup> F. D. Snider,<sup>16</sup> R. Snihur,<sup>32</sup> M. Soderberg,<sup>33</sup> A. Soha,<sup>7</sup>  
 S. Somalwar,<sup>50</sup> V. Sorin,<sup>34</sup> J. Spalding,<sup>16</sup> M. Spezziga,<sup>16</sup> F. Spinella,<sup>44</sup> P. Squillacioti,<sup>44</sup> M. Stanitzki,<sup>58</sup>  
 A. Staveris-Polykalas,<sup>44</sup> R. St. Denis,<sup>20</sup> B. Stelzer,<sup>8</sup> O. Stelzer-Chilton,<sup>32</sup> D. Stentz,<sup>37</sup> J. Strologas,<sup>36</sup> D. Stuart,<sup>10</sup>  
 J. S. Suh,<sup>27</sup> A. Sukhanov,<sup>17</sup> K. Sumorok,<sup>31</sup> H. Sun,<sup>54</sup> T. Suzuki,<sup>53</sup> A. Taffard,<sup>23</sup> R. Tafirout,<sup>32</sup> R. Takashima,<sup>39</sup>  
 Y. Takeuchi,<sup>53</sup> K. Takikawa,<sup>53</sup> M. Tanaka,<sup>2</sup> R. Tanaka,<sup>39</sup> M. Tecchio,<sup>33</sup> P. K. Teng,<sup>1</sup> K. Terashi,<sup>48</sup> S. Tether,<sup>31</sup> J. Thom,<sup>16</sup>  
 A. S. Thompson,<sup>20</sup> E. Thomson,<sup>43</sup> P. Tipton,<sup>47</sup> V. Tiwari,<sup>12</sup> S. Tkaczyk,<sup>16</sup> D. Toback,<sup>51</sup> K. Tollefson,<sup>34</sup> T. Tomura,<sup>53</sup>  
 D. Tonelli,<sup>44</sup> M. Tönnemann,<sup>34</sup> S. Torre,<sup>44</sup> D. Torretta,<sup>16</sup> S. Tourneur,<sup>16</sup> W. Trischuk,<sup>32</sup> R. Tsuchiya,<sup>55</sup> S. Tsuno,<sup>39</sup>  
 N. Turini,<sup>44</sup> F. Ukegawa,<sup>53</sup> T. Unverhau,<sup>20</sup> S. Uozumi,<sup>53</sup> D. Usynin,<sup>43</sup> L. Vacavant,<sup>28</sup> A. Vaiciulis,<sup>47</sup> S. Vallecorsa,<sup>19</sup>  
 A. Varganov,<sup>33</sup> E. Vataga,<sup>36</sup> G. Velev,<sup>16</sup> G. Veramendi,<sup>23</sup> V. Veszpremi,<sup>46</sup> T. Vickey,<sup>23</sup> R. Vidal,<sup>16</sup> I. Vila,<sup>11</sup> R. Vilar,<sup>11</sup>  
 I. Vollrath,<sup>32</sup> I. Volobouev,<sup>28</sup> F. Würthwein,<sup>9</sup> P. Wagner,<sup>51</sup> R. G. Wagner,<sup>2</sup> R. L. Wagner,<sup>16</sup> W. Wagner,<sup>25</sup> R. Wallny,<sup>8</sup>  
 T. Walter,<sup>25</sup> Z. Wan,<sup>50</sup> M. J. Wang,<sup>1</sup> S. M. Wang,<sup>17</sup> A. Warburton,<sup>32</sup> B. Ward,<sup>20</sup> S. Waschke,<sup>20</sup> D. Waters,<sup>30</sup> T. Watts,<sup>50</sup>  
 M. Weber,<sup>28</sup> W. C. Wester III,<sup>16</sup> B. Whitehouse,<sup>54</sup> D. Whiteson,<sup>43</sup> A. B. Wicklund,<sup>2</sup> E. Wicklund,<sup>16</sup> H. H. Williams,<sup>43</sup>  
 P. Wilson,<sup>16</sup> B. L. Winer,<sup>38</sup> P. Wittich,<sup>43</sup> S. Wolbers,<sup>16</sup> C. Wolfe,<sup>13</sup> S. Worm,<sup>50</sup> T. Wright,<sup>33</sup> X. Wu,<sup>19</sup> S. M. Wynne,<sup>29</sup>  
 A. Yagil,<sup>16</sup> K. Yamamoto,<sup>40</sup> J. Yamaoka,<sup>50</sup> Y. Yamashita,<sup>39</sup> C. Yang,<sup>58</sup> U. K. Yang,<sup>13</sup> W. M. Yao,<sup>28</sup> G. P. Yeh,<sup>16</sup> K. Yi,<sup>24</sup>  
 J. Yoh,<sup>16</sup> K. Yorita,<sup>13</sup> T. Yoshida,<sup>40</sup> I. Yu,<sup>27</sup> S. S. Yu,<sup>43</sup> J. C. Yun,<sup>16</sup> L. Zanello,<sup>49</sup> A. Zanetti,<sup>52</sup> I. Zaw,<sup>21</sup> F. Zetti,<sup>44</sup>  
 X. Zhang,<sup>23</sup> J. Zhou,<sup>50</sup> and S. Zucchelli<sup>5</sup>

(CDF Collaboration)

<sup>1</sup>*Institute of Physics, Academia Sinica, Taipei, Taiwan 11529, Republic of China*<sup>2</sup>*Argonne National Laboratory, Argonne, Illinois 60439, USA*<sup>3</sup>*Institut de Física d'Altes Energies, Universitat Autònoma de Barcelona, E-08193, Bellaterra (Barcelona), Spain*<sup>4</sup>*Baylor University, Waco, Texas 76798, USA*<sup>5</sup>*Istituto Nazionale di Fisica Nucleare, University of Bologna, I-40127 Bologna, Italy*<sup>6</sup>*Brandeis University, Waltham, Massachusetts 02254, USA*<sup>7</sup>*University of California, Davis, Davis, California 95616, USA*<sup>8</sup>*University of California, Los Angeles, Los Angeles, California 90024, USA*<sup>9</sup>*University of California, San Diego, La Jolla, California 92093, USA*<sup>10</sup>*University of California, Santa Barbara, Santa Barbara, California 93106, USA*<sup>11</sup>*Instituto de Física de Cantabria, CSIC-University of Cantabria, 39005 Santander, Spain*<sup>12</sup>*Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA*<sup>13</sup>*Enrico Fermi Institute, University of Chicago, Chicago, Illinois 60637, USA*<sup>14</sup>*Joint Institute for Nuclear Research, RU-141980 Dubna, Russia*<sup>15</sup>*Duke University, Durham, North Carolina 27708, USA*<sup>16</sup>*Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA*<sup>17</sup>*University of Florida, Gainesville, Florida 32611, USA*<sup>18</sup>*Laboratori Nazionali di Frascati, Istituto Nazionale di Fisica Nucleare, I-00044 Frascati, Italy*<sup>19</sup>*University of Geneva, CH-1211 Geneva 4, Switzerland*<sup>20</sup>*Glasgow University, Glasgow G12 8QQ, United Kingdom*<sup>21</sup>*Harvard University, Cambridge, Massachusetts 02138, USA*

- <sup>22</sup>*Division of High Energy Physics, Department of Physics, University of Helsinki and Helsinki Institute of Physics, FIN-00014 Helsinki, Finland*
- <sup>23</sup>*University of Illinois, Urbana, Illinois 61801, USA*
- <sup>24</sup>*The Johns Hopkins University, Baltimore, Maryland 21218, USA*
- <sup>25</sup>*Institut für Experimentelle Kernphysik, Universität Karlsruhe, 76128 Karlsruhe, Germany*
- <sup>26</sup>*High Energy Accelerator Research Organization (KEK), Tsukuba, Ibaraki 305, Japan*
- <sup>27</sup>*Center for High Energy Physics, Kyungpook National University, Taegu 702-701; Seoul National University, Seoul 151-742; and SungKyunKwan University, Suwon 440-746, Korea*
- <sup>28</sup>*Ernest Orlando Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA*
- <sup>29</sup>*University of Liverpool, Liverpool L69 7ZE, United Kingdom*
- <sup>30</sup>*University College London, London WC1E 6BT, United Kingdom*
- <sup>31</sup>*Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*
- <sup>32</sup>*Institute of Particle Physics, McGill University, Montréal, Canada H3A 2T8; and University of Toronto, Toronto, Canada M5S 1A7*
- <sup>33</sup>*University of Michigan, Ann Arbor, Michigan 48109, USA*
- <sup>34</sup>*Michigan State University, East Lansing, Michigan 48824, USA*
- <sup>35</sup>*Institution for Theoretical and Experimental Physics, ITEP, Moscow 117259, Russia*
- <sup>36</sup>*University of New Mexico, Albuquerque, New Mexico 87131, USA*
- <sup>37</sup>*Northwestern University, Evanston, Illinois 60208, USA*
- <sup>38</sup>*The Ohio State University, Columbus, Ohio 43210, USA*
- <sup>39</sup>*Okayama University, Okayama 700-8530, Japan*
- <sup>40</sup>*Osaka City University, Osaka 588, Japan*
- <sup>41</sup>*University of Oxford, Oxford OX1 3RH, United Kingdom*
- <sup>42</sup>*Istituto Nazionale di Fisica Nucleare, University of Padova, Sezione di Padova-Trento, I-35131 Padova, Italy*
- <sup>43</sup>*University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA*
- <sup>44</sup>*Istituto Nazionale di Fisica Nucleare Pisa, Universities of Pisa, Siena and Scuola Normale Superiore, I-56127 Pisa, Italy*
- <sup>45</sup>*University of Pittsburgh, Pittsburgh, Pennsylvania 15260, USA*
- <sup>46</sup>*Purdue University, West Lafayette, Indiana 47907, USA*
- <sup>47</sup>*University of Rochester, Rochester, New York 14627, USA*
- <sup>48</sup>*The Rockefeller University, New York, New York 10021, USA*
- <sup>49</sup>*Istituto Nazionale di Fisica Nucleare, Sezione di Roma 1, University of Rome “La Sapienza,” I-00185 Roma, Italy*
- <sup>50</sup>*Rutgers University, Piscataway, New Jersey 08855, USA*
- <sup>51</sup>*Texas A&M University, College Station, Texas 77843, USA*
- <sup>52</sup>*Istituto Nazionale di Fisica Nucleare, University of Trieste/Udine, Italy*
- <sup>53</sup>*University of Tsukuba, Tsukuba, Ibaraki 305, Japan*
- <sup>54</sup>*Tufts University, Medford, Massachusetts 02155, USA*
- <sup>55</sup>*Waseda University, Tokyo 169, Japan*
- <sup>56</sup>*Wayne State University, Detroit, Michigan 48201, USA*
- <sup>57</sup>*University of Wisconsin, Madison, Wisconsin 53706, USA*
- <sup>58</sup>*Yale University, New Haven, Connecticut 06520, USA*

(Received 23 May 2005; revised manuscript received 14 November 2005; published 28 February 2006)

We report the first evidence for a fully reconstructed decay mode of the  $B_c^\pm$  meson in the channel  $B_c^\pm \rightarrow J/\psi\pi^\pm$ , with  $J/\psi \rightarrow \mu^+\mu^-$ . The analysis is based on an integrated luminosity of  $360 \text{ pb}^{-1}$  in  $p\bar{p}$  collisions at 1.96 TeV center of mass energy collected by the Collider Detector at Fermilab. We observe  $14.6 \pm 4.6$  signal events with a background of  $7.1 \pm 0.9$  events, and a fit to the  $J/\psi\pi^\pm$  mass spectrum yields a  $B_c^\pm$  mass of  $6285.7 \pm 5.3(\text{stat}) \pm 1.2(\text{syst}) \text{ MeV}/c^2$ . The probability of a peak of this magnitude occurring by random fluctuation in the search region is estimated as 0.012%.

DOI: [10.1103/PhysRevLett.96.082002](https://doi.org/10.1103/PhysRevLett.96.082002)

PACS numbers: 13.25.Hw, 14.40.Lb, 14.40.Nd

Within the standard model of elementary particles, five of the six different kinds of quarks combine in quark-antiquark pairs to make mesons. The  $B_c^\pm$  meson combines the two heaviest of these quarks as a bottom-charm quark-antiquark pair. Although it has been observed in semileptonic decay modes [1,2], up to now no evidence for the  $B_c^\pm$  has been found in fully reconstructed decay modes [3–6]. Consequently, its mass  $M(B_c)$  has not been measured with good precision.

Nonrelativistic potential models predict the  $\bar{b}$  and  $c$  quarks to be tightly bound with a ground state mass in the approximate range  $6200\text{--}6300 \text{ MeV}/c^2$  [7–9]. Recent QCD-based perturbative computations up to  $\mathcal{O}(\alpha_s^4)$  predict  $M(B_c)$  to be  $6307 \pm 17 \text{ MeV}/c^2$  [10,11]. Most recently, a three-flavor lattice QCD calculation obtains  $M(B_c) = 6304 \pm 12(\text{stat} \oplus \text{syst})_0^{+18}(\text{cutoff effects}) \text{ MeV}/c^2$  [12].

Several of the predicted  $B_c^\pm$  decay modes contain a  $J/\psi$  meson [13]. These are among the most easily reconstruc-

tible  $B_c^\pm$  decays at CDF, owing to an efficient dimuon trigger giving high purity  $J/\psi \rightarrow \mu^+ \mu^-$  reconstruction. The CDF Collaboration made the first observation of the  $B_c^\pm$  meson in the semileptonic decay channels  $B_c^\pm \rightarrow J/\psi l^\pm \nu_l X$ , in a sample of  $110 \text{ pb}^{-1}$  of data at  $\sqrt{s} = 1.8 \text{ TeV}$  in run I at the Tevatron [1]. The symbol  $X$  denotes possible undetected decay particles. With a signal of  $20.4_{-5.5}^{+6.2}$  events, the  $B_c^\pm$  mass was measured to be  $6.40 \pm 0.39(\text{stat}) \pm 0.13(\text{syst}) \text{ GeV}/c^2$ . Recently, the D0 Collaboration reported a preliminary observation of a  $B_c^\pm$  signal in the decay channel  $B_c^\pm \rightarrow J/\psi \mu^\pm \nu_\mu X$  in a sample of  $210 \text{ pb}^{-1}$  of run II data [2].

In this Letter we report first evidence for the  $B_c^\pm$  meson in the fully reconstructed decay channel  $B_c^\pm \rightarrow J/\psi \pi^\pm$ , with  $J/\psi \rightarrow \mu^+ \mu^-$ . The analysis is based on a data set of  $360 \text{ pb}^{-1}$  in  $p\bar{p}$  collisions collected at  $\sqrt{s} = 1.96 \text{ TeV}$  by CDF at the Tevatron during run II.

The CDF II detector consists of a magnetic spectrometer surrounded by calorimeters and muon chambers and is described in detail elsewhere [14]. The components most relevant to this analysis are briefly described here. The tracking system is in a 1.4 T axial magnetic field and consists of a silicon microstrip detector (L00, SVX, ISL, in increasing order of radius) [15–17] surrounded by an open-cell wire drift chamber (COT) [18]. The muon detectors used for this analysis are the central muon drift chambers (CMU), covering the pseudorapidity range  $|\eta| < 0.6$  [19,20], and the extension muon drift chambers (CMX), covering  $0.6 < |\eta| < 1.0$ . Cylindrical coordinates are used with the  $+z$  axis in the proton beam direction.

This measurement uses events containing pairs of muons, each with  $|\eta| < 1.0$ , selected with a three-level trigger. At the first trigger level, muon-candidate track segments in CMU and CMX are matched to COT tracks obtained with a hardware processor [21]. Dimuon triggers use combinations of CMU-CMU and CMU-CMX muons with  $p_T > 1.5(2.0) \text{ GeV}/c$  for CMU (CMX) muons, where  $p_T$  is the momentum transverse to the beam line. At the second level, opening angle and opposite-charge cuts are imposed on the muon pairs. At the third level, three-dimensional (3D) tracking is performed to select muon pairs with invariant mass,  $M(\mu^+ \mu^-)$ , between 2700 and 4000  $\text{MeV}/c^2$ .

To reconstruct the  $B_c^\pm \rightarrow J/\psi \pi^\pm$  decay offline, we make several requirements on the quality of the tracks and the  $J/\psi$  candidate. To ensure good primary and secondary vertex resolution, each track must have an  $r - \phi$  position measurement on at least three of five SVX layers. For  $J/\psi$  identification, we require matching between the COT muon tracks and the muon chamber track segments. In addition, we require that  $3042 < M(\mu^+ \mu^-) < 3152 \text{ MeV}/c^2$ , the average  $J/\psi$  mass resolution in our sample being  $14 \text{ MeV}/c^2$ . Each other charged particle track with  $p_T > 400 \text{ MeV}/c$  is treated as a pion candidate to be combined with the  $J/\psi$ . The pion candidate and the

two muons are then fitted to a common 3D vertex, with  $M(\mu^+ \mu^-)$  constrained to the world average  $J/\psi$  mass value [22]. All combinations for which the vertex fit converged are retained. The primary vertex position is calculated from the other tracks in each event.

The  $B_c^\pm$  search was performed using the following analysis method. The mass values of the  $J/\psi \pi^\pm$  combinations in the search window  $5600 < M(J/\psi \pi^\pm) < 7200 \text{ MeV}/c^2$ , referred to as  $B_c^\pm$  candidates, were temporarily hidden. The search window was chosen to correspond to the  $\pm 2$  standard deviation region around the CDF run I measurement of the  $B_c^\pm$  mass [1]; it is approximately 100 times wider than the expected  $B_c^\pm$  mass resolution.

In order to optimize the significance of a possible  $B_c^\pm$  signal, we varied the selection criteria to maximize the function  $Q = S_F/(1.5 + \sqrt{B_{\text{av}}})$  [23]. Here,  $S_F$  is the accepted fraction of signal events, in this case taken from a Monte Carlo sample, and the background  $B_{\text{av}}$  is the number of selected  $B_c^\pm$  candidates within the search window, scaled to correspond to a mass range of 63  $\text{MeV}/c^2$ , based on the average mass resolution of a  $B_c^\pm$  candidate within the search window. The term 1.5 is appropriate for optimizing a search for a signal at least  $3\sigma$  above background fluctuations. The distributions of the selection variables for the signal events were evaluated using samples of simulated  $B_c^\pm \rightarrow J/\psi \pi^\pm$  decays. These were generated with a  $B_c^\pm$  mass of 6400  $\text{MeV}/c^2$ , a lifetime of 0.46 ps [1], and  $p_T$  and rapidity distributions according to a leading order perturbative QCD calculation [24]. A harder  $p_T$  spectrum [25] was used as an alternative to check the stability of the optimal selection criteria; these were not very sensitive to variations of the  $p_T$  spectrum or the assumed lifetime within its experimental uncertainty. The Monte Carlo  $B_c^\pm$  decays were processed with full detector simulation and the same trigger and reconstruction criteria as the data. The distributions of the selection variables for the background were taken from the data in the search window, in which the contribution from a signal is expected to be small.

Optimized cuts were determined for the following selection variables: the  $J/\psi \pi^\pm$  three-track 3D vertex fit ( $\chi^2 < 9$  for 4 degrees of freedom), the pion track contribution to the vertex fit ( $\chi_\pi^2 < 2.6$ ), the impact parameter in  $r - \phi$  of the  $B_c^\pm$  candidate with respect to the primary vertex ( $< 65 \mu\text{m}$ ), the maximum  $ct$  where  $t$  is the proper decay time of the  $B_c^\pm$  candidate ( $< 750 \mu\text{m}$ ), the transverse momentum of the pion ( $> 1.8 \text{ GeV}/c$ ), the 3D angle between the momentum of the  $B_c^\pm$  candidate and the vector joining the primary to the secondary vertex ( $\beta < 0.4 \text{ rad}$ ), and the significance of the projected decay length of the  $B_c^\pm$  candidate onto its transverse momentum direction [ $L_{xy}/\sigma(L_{xy}) > 4.4$ ]. After these selections, 390 candidates remain in the search window, with no two candidates from the same beam crossing.

A sample of  $B^\pm$  mesons, reconstructed in the decay mode  $B^\pm \rightarrow J/\psi K^\pm$ , was analyzed as a control sample

in order to check our understanding of the reconstruction of the relevant variables in the simulation. The  $B^\pm \rightarrow J/\psi K^\pm$  decay topology is the same as that of  $B_c^\pm \rightarrow J/\psi \pi^\pm$ , apart from the different masses and lifetimes. The  $B^\pm$  mass distribution, shown in Fig. 1, was obtained using the same selection requirements as optimized for the  $B_c^\pm$  candidates, but without the cut on maximum  $ct$ . A total of  $2378 \pm 57$   $B^\pm \rightarrow J/\psi K^\pm$  signal events is found, with a fitted mass of  $5279.0 \pm 0.3$  MeV/ $c^2$ . The fit takes into account a small contribution from the Cabibbo-suppressed decay  $B^\pm \rightarrow J/\psi \pi^\pm$ . The average mass resolution is  $11.5 \pm 0.3$  MeV/ $c^2$ , in agreement with the simulation, which can thus be used with confidence to evaluate the expected mass resolution for  $B_c^\pm$  decays. The  $B^\pm$  yield is used to calculate the expected  $B_c^\pm$  yield. The relative trigger and reconstruction efficiency,  $\epsilon_{B_c^\pm}/\epsilon_{B^\pm}$ , is in the range 35%–85%, with uncertainties arising from the  $B_c^\pm$   $p_T$  spectrum and the  $B_c^\pm$  lifetime. On the basis of the  $B^\pm$  yield, previous CDF cross section measurements [1], and theoretical calculations [13,26–31] of the branching fractions of the  $B_c^\pm \rightarrow J/\psi \pi^\pm$  and  $B_c^\pm \rightarrow J/\psi l^\pm \nu$  decay modes, a  $B_c^\pm$  yield in the range of 10 to 50 events is expected.

A search procedure was then defined to identify any possible signal in the data and to estimate its significance. This was based on a scan of the search region in 10 MeV/ $c^2$  intervals, with a sliding fit window extending from  $-100$  to  $+200$  MeV/ $c^2$  in mass around each nominal peak position,  $m$ . This window was chosen to minimize possible contributions from partially reconstructed  $B_c^\pm$  decays below the peak position (e.g., into  $J/\psi$  and more than one additional particle). For each value of  $m$ , a fit function was defined as a Gaussian signal with mean  $m$ , combined with a linear background term. The Gaussian resolution was fixed as a linear function of  $m$  based on Monte Carlo simulation, and varied from 13 to 19 MeV/ $c^2$  over the search region. The three fit parameters were the number of signal ( $S$ ) and background ( $B$ ) events and the linear back-




FIG. 1 (color online). The invariant mass distribution of the  $B^\pm \rightarrow J/\psi K^\pm$  candidates. The curve is a fit to the data.

ground slope. The output of a scan was defined to be the largest value of  $\Sigma = S/(1.5 + \sqrt{B})$ ,  $\Sigma_{\max}$ , obtained from the 131 fits performed in the mass interval  $5700 \leq M(J/\psi \pi^\pm) \leq 7000$  MeV/ $c^2$ .

The distribution of  $\Sigma_{\max}$  for the null hypothesis was obtained from Monte Carlo experiments [32], in which the mass spectra were derived from a smooth background model. This model was necessarily approximate owing to the initially hidden mass distribution. The model consisted of a linear background, to describe combinatoric events, and a “physical” background to describe partially reconstructed  $B_c^\pm$  decays in the mass range below 6400 MeV/ $c^2$ . Studies showed that the main source of combinatoric background are events in which a genuine  $J/\psi$  is paired with an uncorrelated track. The shape of the physical background was based on Monte Carlo simulations of inclusive  $B_c^\pm \rightarrow J/\psi X$  decays, with branching ratios taken from Ref. [13].

Applied to the 390 event data sample, the scan procedure found a  $\Sigma_{\max}$  near  $m = 6290$  MeV/ $c^2$ , which is compatible with a  $B_c^\pm$  signal of  $19 \pm 6$  events. Using a large set of Monte Carlo simulations, we modeled the shape of the observed background, and, analyzing it in the same way as the data, evaluated the probability that a random enhancement has a  $\Sigma_{\max}$  value exceeding that of the data. This probability was found to be 0.17%.

After the above steps had been performed, further checks on the previously hidden events revealed that the existing pion selection allowed two classes of fitted tracks that were unsuitable for the  $B_c^\pm$  search. The first class had insufficient number of COT hits to give good mass resolution and so was not compatible with a search for a narrow Gaussian signal; the second class had poor SVX resolution in the  $z$  direction and was dominated by combinatorial background. The above two classes of events contributed 10% to the  $B^\pm$  signal; they would be expected to contribute fewer than two events to the  $B_c^\pm$  signal, but they increase the combinatorial background by about 40% over the  $J/\psi \pi^\pm$  mass range. After removal of both classes of poor quality tracks in addition to the original optimized cut selection, 220 candidates remained. These were required to have good SVX  $z$  resolution on both the pion track and at least one of the muon tracks. This final track selection, which maximizes  $\mathcal{Q}$ , is therefore not fully blind; it is based also on the observed properties of the  $B^\pm$  signal and the overall properties of the  $B_c^\pm$  candidate sample.

Figure 2 shows the mass spectrum for the 220 event sample. The main features are the  $B_c^\pm \rightarrow J/\psi \pi^\pm$  signal peak near 6290 MeV/ $c^2$ , a linear combinatorial background above this peak, and a broad enhancement below the peak which can be attributed to the physical background from partially reconstructed  $B_c^\pm$  decays. We perform a global unbinned likelihood fit over the entire mass range to obtain the mass and yield for the  $B_c^\pm$  signal. The fit included a Gaussian signal with a variable mass but with a resolution whose mass-dependent value was determined by






FIG. 2. The invariant mass distribution of the  $J/\psi\pi^\pm$  candidates and results of an unbinned likelihood fit in the search window. The inset shows the peak section of the distribution. The broad enhancement below  $6.2 \text{ GeV}/c^2$  is attributable to partially reconstructed  $B_c^\pm$  mesons.

the Monte Carlo simulation, together with background modeled as a linear combinatorial term and a broad low-mass Gaussian contribution for the physical background. A signal of  $14.6 \pm 4.6$  events is obtained centered at a mass of  $6285.7 \pm 5.3 \text{ MeV}/c^2$ . The standard deviation of the Gaussian at the central value of the signal mass is  $15.5 \text{ MeV}/c^2$ . The background within a region of  $\pm 2$  standard deviations from this mass value is  $7.1 \pm 0.9$  events. The statistical significance of the signal is discussed below. Within the signal region, the distributions of the selection variables agree within statistics with those of the Monte Carlo simulation.

Systematic uncertainties on the  $B_c^\pm$  mass determination due to measurement uncertainties on the track parameters ( $\pm 0.3 \text{ MeV}/c^2$ ) and the momentum scale ( $\pm 0.6 \text{ MeV}/c^2$ ) are evaluated from the corresponding uncertainties on the  $B^\pm$  mass analysis [33]. Further uncertainties are due to the possible differences in the  $p_T$  spectra of the  $B^\pm$  and  $B_c^\pm$  mesons ( $\pm 0.5 \text{ MeV}/c^2$ ) and our limited knowledge of the background shape used in the final mass fit as well as uncertainty in the signal width ( $\pm 0.9 \text{ MeV}/c^2$ ) [34]. The total systematic uncertainty is evaluated to be  $\pm 1.2 \text{ MeV}/c^2$ .

The signal peak is robust under variations of the pion track quality selection. We have investigated several methods for determining the best figure of significance for such a peak over a broad mass range. The method that gives the best sensitivity to a real signal is based on the standard significance measure  $S/\sqrt{B}$ . We repeated the Monte Carlo scans for the new track selection to determine the null hypothesis distribution for  $S/\sqrt{B}$ . Applying to the Monte Carlo simulations the same global fit method as to the data, we find that the probability that a random enhancement anywhere in the range  $5800\text{--}7000 \text{ MeV}/c^2$  exceeds the value of  $S/\sqrt{B}$  for the experimental peak is  $0.012\%$ .

In view of the limited statistics of the observed mass peak, an independent consistency check was performed. If the mass peak is due to fully reconstructed  $B_c^\pm \rightarrow J/\psi\pi^\pm$  decays, partially reconstructed  $B_c^\pm \rightarrow J/\psi + \text{track} + X$  decays should be detectable in the mass region below the peak but not in the region above. The pion candidate in partially reconstructed decays should have a small impact parameter  $d_{xy}$  relative to the  $J/\psi$  vertex, consistent with being physically associated with it, whereas the pion candidate in combinatorial background events should have a broad  $d_{xy}$  distribution reflecting random association with the  $J/\psi$  vertex.

To investigate this, we relax the cuts on  $\beta$ , the impact parameter of the  $B_c^\pm$  candidate, and the  $\chi^2$  of the 3D vertex fit, so as to make a signal in the  $d_{xy}$  distribution visible over the broader combinatorial background. We compare the distribution of  $d_{xy}$  of the pion candidate in the region  $5600 < M(B_c) < 6190 \text{ MeV}/c^2$  (lower side band) to that in the region  $6390 < M(B_c) < 7200 \text{ MeV}/c^2$  (upper side band), where the main contribution should be combinatorial.

Figure 3 (top panel) shows the difference between the lower ( $4900\text{--}5100 \text{ MeV}/c^2$ ) and upper ( $5400\text{--}5700 \text{ MeV}/c^2$ ) sidebands for the  $d_{xy}$  distribution in the  $B^\pm$  data sample, with a large excess of events visible at small  $d_{xy}$  values. Figure 3 (bottom panel) shows the corresponding plot obtained using the  $B_c^\pm$  candidate sample. An enhancement is visible with a shape compatible with that seen in the  $B^\pm$  sample. The  $B^\pm$  curve, rescaled to fit the  $B_c^\pm$  data, provides a good description of this distribution. The excess of low  $d_{xy}$  events in the  $B_c^\pm$  sample is evaluated to be  $244 \pm 59$ , where the uncertainty is statistical only. This result is consistent with Monte Carlo esti-




FIG. 3 (color online). Impact parameter of the third track relative to the  $J/\psi$  vertex for the lower sideband region, after subtraction of the same distribution for the upper sideband: (top panel) the curve is the sum of two Gaussians, fitted to the  $B^\pm$  data points; (bottom panel) the  $B_c^\pm$  data points, overlaid with the above curve, rescaled. In both cases the selection criteria were relaxed.

mates based on the calculations of [13]. This supports the hypothesis that the broad physical background below the signal peak, evident in Fig. 2, is in fact associated with partially reconstructed  $B_c^\pm$  decays.

In conclusion, we observe a peak in the  $J/\psi\pi^\pm$  mass spectrum at a mass of  $6285.7 \pm 5.3(\text{stat}) \pm 1.2(\text{syst}) \text{ MeV}/c^2$ . This peak is consistent with a narrow, weakly decaying particle state and is interpreted as the first evidence for fully reconstructed decays of the  $B_c^\pm$  meson. The mass value has much improved precision over the results obtained in  $B_c^\pm$  semileptonic decays [1,2]. There is also good agreement with recent theoretical predictions for the  $B_c^\pm$  mass around  $6300 \text{ MeV}/c^2$  [10–12].

We thank the Fermilab staff and the technical staffs of the participating institutions for their vital contributions. We also thank A. V. Berezhnoy, C. H. Chang, and X. G. Wu for making available their calculations of  $B_c^\pm$  production spectra. This work was supported by the U.S. Department of Energy and the National Science Foundation, the Italian Istituto Nazionale di Fisica Nucleare, the Ministry of Education, Culture, Sports, Science and Technology of Japan, the Natural Sciences and Engineering Research Council of Canada, the National Science Council of the Republic of China, the Swiss National Science Foundation, the A. P. Sloan Foundation, the Bundesministerium für Bildung und Forschung, Germany, the Korean Science and Engineering Foundation and the Korean Research Foundation, the Particle Physics and Astronomy Research Council and the Royal Society, U.K., the Russian Foundation for Basic Research, the Comision Interministerial de Ciencia y Tecnologia, Spain, in part by the European Community's Human Potential Programme under Contract No. HPRN-CT-2002-00292, and the Academy of Finland.

- 
- [1] F. Abe *et al.* (CDF Collaboration), Phys. Rev. Lett. **81**, 2432 (1998); F. Abe *et al.*, Phys. Rev. D **58**, 112004 (1998).
  - [2] E. Cheu (D0 Collaboration), Int. J. Mod. Phys. A **20**, 3664 (2005).
  - [3] F. Abe *et al.* (CDF Collaboration), Phys. Rev. Lett. **77**, 5176 (1996).
  - [4] P. Abreu *et al.* (DELPHI Collaboration), Phys. Lett. B **398**, 207 (1997).
  - [5] R. Barate *et al.* (ALEPH Collaboration), Phys. Lett. B **402**, 213 (1997).
  - [6] K. Ackerstaff *et al.* (OPAL Collaboration), Phys. Lett. B **420**, 157 (1998).

- [7] W. Kwong and J. Rosner, Phys. Rev. D **44**, 212 (1991).
- [8] E. Eichten and C. Quigg, Phys. Rev. D **49**, 5845 (1994).
- [9] S. Godfrey, Phys. Rev. D **70**, 054017 (2004).
- [10] N. Brambilla, Y. Sumino, and A. Vairo, Phys. Rev. D **65**, 034001 (2002).
- [11] N. Brambilla *et al.*, hep-ph/0412158, and references therein.
- [12] I. F. Allison *et al.*, Phys. Rev. Lett. **94**, 172001 (2005); Nucl. Phys. B, Proc. Suppl. **140**, 440 (2005).
- [13] V. V. Kiselev, Phys. At. Nucl. **67**, 1559 (2004).
- [14] D. Acosta *et al.* (CDF Collaboration), Phys. Rev. D **71**, 032001 (2005).
- [15] A. Sill *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **447**, 1 (2000).
- [16] A. Affolder *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **453**, 84 (2000).
- [17] C. S. Hill *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **530**, 1 (2004).
- [18] T. Affolder *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **526**, 249 (2004).
- [19] G. Ascoli *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **268**, 33 (1988).
- [20] T. Dorigo *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **461**, 560 (2001).
- [21] E. J. Thomson *et al.*, IEEE Trans. Nucl. Sci. **49**, 1063 (2002).
- [22] S. Eidelman *et al.*, Phys. Lett. B **592**, 1 (2004).
- [23] G. Punzi, in *Proceedings of PhyStat2003 (SLAC, 2003)*, econf C030908, MODT002 (2003).
- [24] C. H. Chang and X. G. Wu (private communication); C. H. Chang *et al.*, Comput. Phys. Commun. **159**, 192 (2004); C. H. Chang and X. G. Wu, Eur. Phys. J. C **38**, 267 (2004).
- [25] A. V. Berezhnoy (private communication); A. V. Berezhnoy, V. V. Kiselev, and A. K. Likhoded, Z. Phys. A **356**, 79 (1996); A. V. Berezhnoy, A. K. Likhoded, and M. V. Shevlyagin, Phys. At. Nucl. **58**, 1732 (1995).
- [26] C. H. Chang *et al.*, Phys. Rev. D **49**, 3399 (1994).
- [27] A. V. Berezhnoy, V. V. Kiselev, and A. K. Likhoded, Phys. At. Nucl. **61**, 252 (1998).
- [28] A. Yu. Anisimov *et al.*, Phys. At. Nucl. **62**, 1739 (1999).
- [29] P. Colangelo *et al.*, Phys. Rev. D **61**, 034012 (2000).
- [30] A. Abd El-Hady *et al.*, Phys. Rev. D **62**, 014019 (2000).
- [31] K. Anikeev *et al.*, hep-ph/0201071.
- [32] W. A. Rolke and A. M. Lopez, in *Proceedings of PhyStat2003 (SLAC, 2003)*, econf C030908, MODT002 (2003).
- [33] D. Acosta *et al.* (CDF Collaboration), hep-ex/0508022 (to be published).
- [34] L. Nicolas, Ph.D. dissertation, University of Glasgow, 2005.