

1-1-1995

Increasing Concentrations of 17β -Estradiol Has Differential Effects on Secretion of Luteinizing Hormone and Follicle-Stimulating Hormone and Amounts of mRNA for Gonadotropin Subunits during the Follicular Phase of the Bovine Estrous Cycle

Andrea S. Cupp

University of Nebraska-Lincoln, acupp2@unl.edu

F. N. Kojima

University of Nebraska-Lincoln

M. S. Roberson

Oregon Health Sciences University, Portland, OR

T. T. Stumpf

University of Missouri, Columbia

M. F. Wolfe

Case Western Reserve University

Cupp, Andrea S.; Kojima, F. N.; Roberson, M. S.; Stumpf, T. T.; Wolfe, M. F.; Werth, L. A.; Kittok, Roger J.; Grotjan, H. E.; and Kinder, J. E., "Increasing Concentrations of 17β -Estradiol Has Differential Effects on Secretion of Luteinizing Hormone and Follicle-Stimulating Hormone and Amounts of mRNA for Gonadotropin Subunits during the Follicular Phase of the Bovine Estrous Cycle" (1995). *Faculty Papers and Publications in Animal Science*. 186.
<http://digitalcommons.unl.edu/animalscifacpub/186>

This Article is brought to you for free and open access by the Animal Science Department at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Papers and Publications in Animal Science by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

See next page for additional authors

Follow this and additional works at: <http://digitalcommons.unl.edu/animalscifacpub>

 Part of the [Animal Sciences Commons](#)

Authors

Andrea S. Cupp, F. N. Kojima, M. S. Roberson, T. T. Stumpf, M. F. Wolfe, L. A. Werth, Roger J. Kittok, H. E. Grotjan, and J. E. Kinder

Increasing Concentrations of 17 β -Estradiol Has Differential Effects on Secretion of Luteinizing Hormone and Follicle-Stimulating Hormone and Amounts of mRNA for Gonadotropin Subunits during the Follicular Phase of the Bovine Estrous Cycle¹

A.S. CUPP, F.N. KOJIMA, M.S. ROBERSON,³ T.T. STUMPF,⁴ M.W. WOLFE,⁵ L.A. WERTH,⁶ R.J. KITTOK, H.E. GROTTJAN, and J.E. KINDER²

Department of Animal Science, University of Nebraska, Lincoln, Nebraska 68583–0908

ABSTRACT

The hypothesis tested was that 17 β -estradiol (E₂) would increase amounts of mRNA for α , LH β , and FSH β subunits during the follicular phase of the estrous cycle prior to the preovulatory surge of gonadotropins in cows. On Day 16 (Day 0 = estrus) of the estrous cycle, all cows were treated with prostaglandin F_{2 α} (PGF_{2 α}). Cows served as intact controls (CONT, n = 4) were ovariectomized (OVX, n = 5), or were ovariectomized and administered E₂ (OVXE, n = 6) in increasing doses starting at the time of treatment with PGF_{2 α} . Cows were bled for 6 h before and for 40 h after PGF_{2 α} treatment to characterize pulsatile secretion of LH and FSH. Forty hours after PGF_{2 α} treatment, pituitaries were collected for evaluation of amounts of mRNA for α , LH β , and FSH β subunits. Amplitude of LH pulses was greater ($p < 0.05$) in cows from the OVXE than from the CONT group. Concentrations of FSH were greater in cows from both the OVXE and OVX ($p < 0.01$) groups than from the CONT group. Amounts of mRNA for α and FSH β subunits were greater ($p < 0.01$) in pituitaries of cows from the OVX than from the CONT or OVXE groups. Amounts of mRNA for LH β subunit in pituitaries of cows from the OVX group tended to be greater ($p < 0.08$) than from the CONT group. Cows in the OVXE group tended ($p < 0.08$) to have greater amounts of mRNA for FSH β subunit than did CONT cows. Amounts of mRNA for α and LH β subunits in cows from the OVXE and CONT groups did not differ ($p > 0.10$). Pituitary weight and content of LH and FSH were not different ($p > 0.10$) among cows of the different groups. Ovariectomy resulted in enhanced secretion of gonadotropins and increased amounts of mRNA for gonadotropin subunits above values detected in CONT cows. Ovariectomized cows administered E₂ in follicular-phase patterns had amounts of mRNA for α and LH β similar to those in CONT cows even though secretion of LH was enhanced in the OVXE group. We reject our hypothesis and conclude that E₂ has a divergent role in regulation of gonadotropins. Release of LH is enhanced by E₂, but E₂ reduced mRNA for gonadotropin subunits (tended to reduce mRNA for FSH β subunit) in ovariectomized cows to amounts detected in intact cows during the follicular phase of the bovine estrous cycle.

INTRODUCTION

During the follicular phase of the estrous cycle in cows, increased 17 β -estradiol (E₂) enhanced secretion of LH [1, 2]. Estradiol modulates LH secretion in the ewe by influencing secretion of LHRH [3] and up-regulating receptors for LHRH at the pituitary [4, 5]. Enhanced secretion of LHRH in rodents results in a similar up-regulation of receptors for LHRH [6, 7], and enhanced secretion of LH is a consequence of the increased population of receptors for LHRH.

Stimulation of gonadotrophs by LHRH contributes to the production of mRNA for gonadotropin subunits (ewe [8]; rat [9, 10]). Pulsatile secretion of LHRH is required to maintain greater amounts of mRNA for LH β (ewe [11]; rat [12]) and FSH β subunits (ewe [11]). Increases in the pulse fre-

quency of LHRH (rat [13]) or continuous infusion of LHRH (ewe [11]) results in preferential synthesis of α subunit mRNA. Therefore, alterations in pattern of LHRH secretion from the hypothalamus results in changes in amounts of mRNA for gonadotropin subunits.

Modulation of pulse frequency of LHRH occurs during different physiological states. Pulse frequency of LHRH is increased during sexual maturation in rats [14], during the follicular phase of the menstrual cycle in primates [15–17], and prior to the preovulatory surge of gonadotropins in ewes [18, 19]. Administration of progesterone, E₂, or both steroids in combination is hypothesized to regulate the pattern of secretion of LHRH or up-regulate receptors for LHRH to cause increased production of mRNA for the LH or FSH subunit [20, 21].

In the ewe, amounts of mRNA for both α and LH β subunits increase around estrus, whereas amounts of mRNA for FSH β subunit are minimal [21]. In contrast, in experiments where E₂ was administered chronically to ewes, suppression of mRNA for α , LH β [22], and FSH β subunits occurred [23]. Experiments have not been conducted in cows to determine the effect of a physiological increase in E₂ during the follicular phase of the estrous cycle on the amount of mRNA for gonadotropin subunits. In the present experiment we tested the hypothesis that elevated concentrations of E₂ (similar to follicular phase increases) that result in

Accepted September 21, 1994.

Received March 16, 1994.

¹Published as paper No. 10525, Journal Ser. Nebraska Agr. Res. Div., Research supported by USDA CRGO 88–37240–3966.

²Correspondence: James E. Kinder, A224J Animal Sciences, University of Nebraska, Lincoln, NE 68583–0908. FAX: (402) 472–6362.

³Current address: Department of Cell Biology and Anatomy, Oregon Health Sciences University, Portland, OR 97201.

⁴Current address: Department of Animal Science, University of Missouri, Columbia, MO 65211.

⁵Current address: Department of Pharmacology, Case Western Reserve University, Cleveland, OH 44106.

⁶Current address: Department of Veterinary and Biomedical Sciences, University of Nebraska, Lincoln, NE 68583–0905.

increased secretion of LH during the follicular phase of the bovine estrous cycle prior to the preovulatory surge of gonadotropins would increase amounts of mRNA for α and LH β subunits. In addition, we evaluated the influence of this change in E₂ on the amount of mRNA for FSH β subunit.

MATERIALS AND METHODS

Experimental Protocol

All procedures used in this experiment were approved by the Institutional Animal Care and Use Committee of the University of Nebraska, Lincoln. Fifteen cows of composite breeding (1/4 Hereford, 1/4 Angus, 1/4 Red Poll, 1/4 Pinzgauer) were synchronized to a common day of the estrous cycle with two injections of prostaglandin F_{2 α} (PGF_{2 α} ; Lutalyse® Sterile Solution; The Upjohn Co., Kalamazoo, MI). On Day 16 (Day 0 = estrus) of the estrous cycle, all cows were treated with PGF_{2 α} . Cows of one group served as controls (CONT, n = 4), and cows in the other two groups were either ovariectomized (OVX, n = 5) or ovariectomized and administered increasing concentrations of E₂ (OVXE, n = 6) via vaginal implants starting at the time of treatment with PGF_{2 α} . Ovariectomies were performed via high lumbar laparotomy (0 h). After ovariectomy, cows in the OVXE group received intravaginal implants (polydimethylsiloxane intravaginal implants: 3.35 mm i.d. \times 4.65 mm o.d. \times 13.5 cm; Dow-Corning, Midland, MI) containing E₂ (Sigma Chemical Co., St. Louis, MO). The number of implants was increased (one per time period) at 0, 10, 15, 20, and 30 h to mimic the elevation of E₂ that occurs during the follicular phase of the bovine estrous cycle (CONT).

Before PGF_{2 α} treatment, indwelling jugular catheters were inserted, and blood samples were collected at 12-min intervals for 6 h to determine initial concentrations of hormones in circulation. After treatment with PGF_{2 α} and initiation of treatments with E₂ in cows from the OVX and OVXE groups, cows were bled every 12-min for four 6-h periods with 4 hourly samples between each serial blood collection. The last 6-h period of blood collection ended 40 h after administration of PGF_{2 α} and initiation of treatments with E₂. Blood samples were collected from cows in the control group after the same protocol of blood collection as that for ovariectomized cows. These blood samples were collected to characterize changes in concentration and pulsatile secretion of LH and FSH. Additional blood samples were collected every 5 h to determine plasma concentrations of E₂ and progesterone in circulation.

After the last 6-h period of serial blood collection, pituitaries were collected from all cows within 15 min after exsanguination. Anterior pituitaries were hemisected, transferred to separate vials, frozen in liquid nitrogen, and stored at -70°C until amounts of mRNA for α , LH β , and FSH β subunits were determined.

RIAs

Blood samples for quantification of LH were allowed to clot at room temperature and were stored at 4°C for 24 h. Serum was collected after centrifugation at $1520 \times g$ for 15 min and stored at -20°C until assayed for LH. Concentrations of LH in serum samples collected at 12-min intervals and in pituitary extracts were determined by RIA [24, 25]. Intra- and interassay coefficients of variation for serum LH assays were 2.5 and 10.5%, respectively. Intra- and interassay coefficients of variation for pituitary extracts were 3.0 and 13.2%, respectively. Sensitivities of the LH assays were approximately 28 pg/ml in serum and 30 pg/ml in pituitary homogenates.

Concentrations of FSH were analyzed in all serum samples and pituitary extracts by RIA [26, 27]. Intra- and interassay coefficients of variation for FSH assays in serum were 2.1 and 11.2% and in pituitaries were 2.4 and 12.7%, respectively. Sensitivity of the FSH assays were 44 pg/ml in serum and 74 pg/ml in pituitary homogenates.

Blood samples taken every 5 h were collected in heparinized tubes and centrifuged at $1520 \times g$ for 20 min, and plasma was stored at -20°C until assayed for progesterone [28] and E₂ [29]. Concentrations of progesterone were determined in a single assay, and the intraassay coefficient of variation was 5%. Intra- and interassay coefficients of variation for E₂ were 2.4 and 17.5%, respectively. Sensitivity of the progesterone assay was 600 pg/ml, and sensitivities of the E₂ assays were 0.5 pg/ml.

Pituitary Extracts

To determine pituitary content of gonadotropins, hemipituitaries were homogenized in 150 mM NaCl buffered with 50 mM Tris, pH 7.4, containing 0.5% (vol/vol) Triton X-100, 5 mM Na₂ EDTA, 1 mM phenylmethylsulphonyl fluoride, 0.5 mg/L leupeptin, and 200 U/ml aprotinin (1.0 ml/100 mg wet tissue weight) for 30 sec in a polytron homogenizer (Brinkman Instruments, Westbury, NY). Pituitary extracts were clarified by centrifugation at $100\,000 \times g$ for 1 h, aliquoted into 0.5-ml portions (equivalent to 50 mg tissue), and stored at -70°C until assayed for LH [24, 25] and FSH [25, 26]. Pituitary extracts were also used to quantify differences in isoforms of LH and FSH among treatments; these data are presented in a companion paper [30].

Northern and Slot Blot Analysis of Total Cellular RNA (tcRNA)

Frozen hemipituitaries were weighed, and tcRNA was isolated by homogenization in buffer and centrifugation through a cesium chloride gradient [22]. Precipitates of tcRNA were maintained under ethanol at -70°C until Northern (α and LH β subunits) and slot (FSH β subunit) blots were prepared and hybridized to appropriate subunit probes. Procedures to determine amounts of mRNA for α and LH β subunits in pituitaries of cows using Northern blots were described previously [29].




FIG. 1. Mean concentrations (in pg/ml) of E₂ in blood samples for each group collected every 5 h starting with PGF_{2α} treatment and/or ovariectomy and ending at pituitary collection. Pooled SEM = 0.50 pg/ml.




FIG. 2. Mean concentrations (in ng/ml) of progesterone in blood samples for each group collected every 5 h starting with PGF_{2α} treatment and/or OVX and ending at pituitary collection. Pooled SEM = 0.19 ng/ml.

Briefly, Northern blots were prepared by separating 5 μ g of tcrRNA from individual hemi-pituitaries in 1.5% agarose gels containing formaldehyde, where 18s and 28s bands were visualized for efficiency of RNA loading per well. Within each gel, samples of tcrRNA from cows in each treatment group were represented. Total cellular RNA isolated from cerebellum and steer pituitaries were used as negative and positive controls, respectively. Size-separated tcrRNA was subsequently transferred to nitrocellulose membranes (Beth Research Laboratory, Gaithersburg, MD) by capillary action and baked at 80°C for 2 h. Plasmids containing cDNA fragments encoding bovine α - [31] and LH β [32] subunits (generously provided by Dr. Richard Maurer) were used to derive riboprobes that were labeled with α -³²P-CTP (Amersham Corp., Arlington Heights, IL).

Slot blots to determine amounts of mRNA for FSH β subunit were prepared with 5 μ g of tcrRNA and diethylpyrocarbonate distilled water to 10 μ l total volume. RNA was denatured by addition of 20 μ l 100% formamide, 7 μ l 37% formaldehyde, and 2 μ l 20-strength saline sodium citrate (SSC; single-strength SSC is 0.15 M NaCl and 0.015 M sodium citrate), heated to 68°C for 15 min, and then placed on ice [33]. Before RNA was loaded onto slot blots, wells were washed twice with 10-strength SSC (250 μ l/wash for each well). Two volumes of 20-strength SSC containing bromophenol blue dye were added to RNA samples and applied to nylon filters (Zeta probe, Bio-Rad Labs., Richmond, CA) in individual wells on slot blots. The nylon filters were rinsed four times with 10-strength SSC, dried and baked for 30 min at 80°C. The cDNA probe for FSH β subunit [34] was prepared by random-primer labeling [35] utilizing (α -³²P-dCTP, Amersham Corp.). Heterologous oligonucleotide radioactive probes to rat α -tubulin (NEN DuPont, Boston, MA;

nucleotides +958 to +987 of rat cDNA) were prepared by 3' end labeling (3' end labeling kit, NEN DuPont) with ³⁵S-dATP (Amersham Corp.) according to manufacturers' instructions.

Total cellular RNA bound to nylon and nitrocellulose filters were hybridized to α and LH β subunits and α -tubulin probes for 46 h at 60°C. All treatments were represented on one slot blot that was hybridized to FSH β subunit at 45°C for 24 h. Specific activity of probes was 1.5–2.0 \times 10⁹ dpm/ μ g. The concentration of radioactive probes was approximately 10⁶ total cpm/ml hybridization buffer (5 ml/blot). After hybridization, filters were washed at room temperature (3 washes; 20 min/wash) in double-strength SSC, 0.1% SDS and then in 0.2-strength SSC, 0.1% SDS at 60°C.

Specific radiolabeled bands from Northern and slot blots were visualized and quantified by use of an Ambis Radioanalytic Imaging System. Quantification of the radiolabeled probe hybridizing to mRNA bound to nitrocellulose or nylon was expressed as net counts per minute detected within the discrete area of the radiolabeled band.

Statistical Analysis

Values for each characteristic of LH and FSH secretion were analyzed for the five 6-h periods of serial blood collection. Mean concentrations of LH and FSH (ng/ml), frequency of LH and FSH pulses (pulses/6 h), and amplitude of LH and FSH pulses (ng/ml) were determined through the use of algorithms (Pulsar Software, Univ. of Illinois, Urbana, IL, modified for the IBM-PC by J.R. Gitzen and V.D. Ramirez, Univ. of Illinois, Urbana, IL, [G-values for LH were G1—5.00, G2—2.80, G3—2.30, G4—10.00, G5—10.00; G-values for FSH were: G1—3.20, G2—2.80, G3—2.30, G4—10.00, G5—10.00]).

TABLE 1. Concentrations of LH prior to and during treatment period.

Group	Mean LH (ng/ml)		LH pulse amplitude (ng/ml)		LH Pulse frequency/12 h	
	Pretreatment	Treatment	Pretreatment	Treatment	Pretreatment	Treatment
CONT	0.87 ^a	1.74 ^c	1.11 ^a	1.65 ^{a,c}	1.75 ^a	2.74 ^{a,c}
OVX	0.89 ^a	2.30 ^{b,c}	1.00 ^a	2.65 ^b	1.40 ^a	4.05 ^b
OVXE	0.75 ^a	2.70 ^b	1.41 ^a	3.07 ^b	1.00 ^a	3.62 ^{b,c}
SEM	0.20	0.32	0.47	0.38	0.54	0.65
	P > 0.10	P < 0.05	P > 0.10	P < 0.01	P > 0.10	P < 0.05

^{a,b,c}All values with different superscripts within columns are different at P-values listed and within rows at $p < 0.05$.

Data regarding the effect of treatment on characteristics of pulsatile secretion of LH and FSH and plasma samples of E₂ and progesterone were fitted to a linear model containing the fixed effect of blood collection period and the random effect of cow. Variance components were estimated by means of Restricted Maximum Likelihood algorithms of PROC MIXED in SAS [36, 37]. Repeated measures were accounted for by assuming an autocorrelation error structure for residual error that estimates the correlation among repeated measurements on an individual cow. Therefore, all parameters for LH and FSH were calculated over time (four 6-h blood collection periods after PGF_{2α}) during the treatment period to determine differences between treatments. The plasma samples collected for E₂ and progesterone (every 5 h) were also analyzed over time to determine differences among groups.

The means of the characteristics for LH and FSH secretion from the combined blood collection periods after PGF_{2α} were compared to the initial blood collection (since this blood collection occurred before administration of treatments). In addition, comparisons for steroid hormone concentration (E₂ and progesterone) were made between the one sample collected before treatments and all collections after treatment administration. Amount of mRNA for gonadotropin subunits and α-tubulin were compared according to PROC MIXED procedures with contrast statements to determine differences among treatments [36, 37].

RESULTS

Stage of Estrous Cycle

There were no differences in characteristics of hormones in serum during the initial blood collection period prior to administration of PGF_{2α} and/or ovariectomy. No cows in the CONT group exhibited a surge of LH during the blood collection periods, and large ovulatory follicles with no evidence of functional CL were present when ovaries were evaluated after collection of pituitary tissue. Therefore, we conclude that all CONT cows were in the follicular phase of the estrous cycle throughout the experimental period.

Concentrations of Steroid Hormones

Data for mean concentrations of E₂ are depicted in Figure 1 for each group during the period of blood collection.

Concentrations of E₂ were similar in all cows prior to PGF_{2α} treatment. Cows in the OVXE group had concentrations of E₂ in circulation similar to those of CONT cows ($p > 0.10$; Fig. 1). Cows in both the OVXE and CONT groups had greater ($p > 0.05$) concentrations of E₂ than did OVX cows. Concentrations of E₂ remained low in cows from the OVX group (less than level of sensitivity of the assay) throughout the experiment. Mean concentrations of progesterone for each group are depicted in Figure 2. Concentrations of progesterone were similar in cows of all groups ($p > 0.10$).


Pattern of LH Secretion

Mean concentrations of LH were greater ($p < 0.05$) in cows in the OVXE than in the CONT group when compared over all blood collection periods after PGF_{2α} treatment (Table 1). Mean concentrations of LH were similar among cows in the OVX and CONT groups for each of the four 6-h periods of serial blood collection after PGF_{2α} administration ($p > 0.10$). Frequency of LH pulses were greater ($p < 0.05$) in cows in the OVX group than in the CONT group during the combined blood collection periods after PGF_{2α} treatment (Table 1). Amplitude of LH pulses was greater ($p < 0.01$) in cows in the OVXE group than in the CONT group during the blood collection periods after PGF_{2α} was administered (Table 1). The profiles of pulsatile LH secretion during each of the five periods of serial blood collection are depicted for representative cows from each treatment in Figure 3.

Pattern of FSH Secretion

Mean concentrations of FSH were greater in cows in the OVX and OVXE ($p < 0.01$) groups than in the CONT cows during the blood collections after PGF_{2α} treatment (Table 2). However, FSH remained similar ($p > 0.10$) in cows in the OVX and OVXE groups during the same time period (Table 2). During the 6-h period of blood sampling prior to pituitary collection, cows in the OVX group had greater concentrations of FSH in circulation than did the cows of either the OVXE ($p < 0.01$) or CONT group ($p < 0.001$; OVX = 3.38; OVXE = 2.73; CONT = 1.67 ng/ml). In addition, cows in the OVXE group had greater concentrations of FSH than did cows in the CONT group ($p < 0.001$).

The patterns of secretion of FSH in cows from the ovariectomized groups were different from the pattern in the



CONT group (Fig. 3). During the four blood collection periods after initiation of treatment, cows in the OVX and OVXE groups had a greater ($p < 0.01$) amplitude of FSH pulses than did cows from the CONT group ($0.57, 0.58, \text{ and } 0.05 \pm 0.14$ ng/ml, respectively). Cows in the OVXE group had a greater frequency of FSH pulses than did cows in either the CONT or OVX groups ($1.40, 0.12, 0.97 \pm 0.33$ pulses/6 h, respectively). Only two cows in the CONT group had pulses of FSH during the four serial blood collections after initiation of treatment, and in both cows pulses of FSH were low in amplitude ($0.32, 0.43$ ng/ml). Only two cows (one OVX and one OVXE) had pulses of FSH during the serial blood collection prior to treatment.

Gonadotropin Content and Relative Amounts of mRNA in Pituitaries

Hemisected pituitary weights (CONT = 929.0, OVXE = 860.3, OVX = 899.4 ± 78.2 mg) and concentrations of LH (CONT = 3.3, OVXE = 3.2, OVX = 2.7 ± 0.4 $\mu\text{g}/\text{mg}$ tissue) and FSH (CONT = 0.088, OVXE = 0.099, OVX = 0.098 ± 0.009 $\mu\text{g}/\text{mg}$ of tissue) were not different ($p > 0.10$) among cows in the different treatment groups. Concentrations of tRNA isolated from pituitaries were not affected by treatment ($p > 0.10$) and averaged 5 ± 1 μg tRNA/mg in pituitary tissue. Absorbance ratios of A260/A280 and A260/A230 were similar ($p > 0.05$) among pituitary tissues from cows in different groups and averaged 1.62 ± 0.03 and 2.47 ± 0.06 , respectively. Total cellular RNA isolated from cow cerebellum did not specifically hybridize to either α , LH β , or FSH β subunit probes.

Relative amounts of α -tubulin detected in tRNA were similar ($p > 0.10$) among pituitary tissues of cows from all groups (Fig. 4d), indicating that effects of E₂ were specific in modulating amounts of mRNA for the gonadotropin subunits. If there were nonspecific actions of estradiol in increasing amounts of mRNA in the pituitary, amount of mRNA for α -tubulin would have been influenced by treatment with E₂. There were greater amounts of mRNA for α -tubulin in cerebellum compared to pituitary tissue ($p < 0.05$).

Amounts of mRNA for α subunit were greater in cows in the OVX than in the CONT group (630 ± 55 and 406 ± 62 counts/5 μg tRNA; $p < 0.01$; Fig. 4a). Amounts of mRNA for LH β subunit tended to be greater in pituitaries of cows in the OVX than in the CONT or OVXE groups (225 ± 27 and 150 ± 30 or 178 ± 25 counts/5 μg tRNA, respectively; $p < 0.08$; Fig. 4b). Cows in the OVXE group had amounts of mRNA for α and LH β subunits similar to amounts in CONT cows ($p > 0.10$). Amounts of mRNA for FSH β subunit were greater in cows of the OVX than in those of the CONT (p

FIG. 3. Profiles of LH and FSH secretion in representative cows from each treatment group for each of the five periods of serial blood collection, which followed the protocol described in *Material and Methods*. Top) CONT, middle) OVXE, bottom) OVX.

TABLE 2. Concentrations of FSH prior to and during treatment period.

Group	Mean FSH (ng/ml)		FSH pulse amplitude (ng/ml)		FSH pulse frequency/12 h	
	Pretreatment	Treatment	Pretreatment	Treatment	Pretreatment	Treatment
CONT	1.75 ^a	1.77 ^a	0.00 ^a	0.05 ^a	0.00 ^a	0.12 ^a
OVX	1.90 ^a	2.79 ^b	0.11 ^a	0.56 ^b	0.40 ^a	0.97 ^{**}
OVXE	2.01 ^a	2.55 ^b	0.12 ^a	0.57 ^b	0.16 ^a	1.39 ^b
SEM	0.22	0.24	0.22	0.15	0.54	0.33
	P > 0.10	P < 0.01	P > 0.10	P < 0.01	P > 0.10	P < 0.01

^{a,b}All values with different superscripts within columns are different at P-values listed and within rows at $p < 0.05$.

*Tends to be different than CONT; ($p < 0.08$).

< 0.001) or OVXE ($p < 0.01$) groups (13043 ± 1525 and 2833 ± 1705 or 7079 ± 1392 count/5 μg tcRNA, respectively, Fig. 4c). Cows in the OVXE group tended ($p < 0.08$) to have greater amounts of FSH β subunit mRNA than did cows from the CONT group.

DISCUSSION

In the present experiment, ovariectomy and treatment with E_2 resulted in enhanced secretion of LH and FSH. Cows of the OVX group tended to have enhanced amounts of mRNA for all gonadotropin subunits (α , LH β and FSH β) compared to cows of the CONT group. However, amounts of mRNA for α and LH β subunits were similar in cows of the OVXE and CONT groups. Although E_2 enhanced secretion of LH, it did not enhance amounts of mRNA for LH subunits during the follicular phase of the bovine estrous cycle as we had hypothesized. Therefore, we reject our hypothesis that increasing concentrations of E_2 during the follicular phase of the bovine estrous cycle enhance amounts of mRNA for the LH subunits coincident with the increase in secretion of LH that occurs during this part of the estrous cycle.

Many experiments have been conducted to determine the effect of E_2 on steady-state or relative amounts of mRNA for gonadotropin subunits, with mixed results. Chronic administration of E_2 reduced amounts of mRNA for gonadotropin subunits in vivo [22, 23, 38, 39]. During periods of the estrous cycle when concentrations of E_2 are greater in circulation (such as the follicular phase and around estrus), amounts of mRNA for α and LH β subunits were increased whereas mRNA for FSH β subunit was minimal [21].

In the present experiment, when E_2 was administered to ovariectomized cows in a pattern that mimicked that of the follicular phase of the bovine estrous cycle, amounts of mRNA for all gonadotropin subunits were suppressed compared to amounts in ovariectomized cows not treated with E_2 . This demonstrates the negative effect of physiological concentrations of E_2 on amounts of mRNA for α , LH β , and FSH β subunits. Our results in cattle concur with those from experiments in other species (rat [40]; ewe [22, 41]) where chronically ovariectomized animals were administered either a constant high or low dose of E_2 for relatively long periods of time.

Secretion of LH (mean concentration and pulse amplitude of LH) was enhanced in ovariectomized cows treated with increasing concentrations of E_2 compared to CONT cows. Because the pattern of circulating concentrations of steroids was similar in cows of the OVXE and CONT groups, ovarian factors may exist that regulate LH secretion other than ovarian steroids. Possible ovarian factors that could elicit this effect are unknown; thus, it would be pure conjecture to speculate on their identity.

Secretion of FSH differs from that of LH because it is coordinately regulated by ovarian peptides—inhibin and




FIG. 4. Mean concentrations of (a) α subunit, (b) LH β subunit, (c) FSH β subunit, and (d) α -tubulin subunit mRNA expressed in net counts of radioactivity per minute for each group at pituitary collection 40 h after PGF $_{2\alpha}$ treatment. SEM for α subunit mRNA = 57 counts/min. SEM for LH β mRNA = 27 counts/min. SEM for FSH β mRNA = 1525 counts/min. SEM for α -tubulin mRNA = 69 counts/min. **Differences from CONT ($p < 0.01$). *Differences from CONT ($p < 0.08$).

activin—as well as LHRH [42]. In sheep, suppression of secretion of FSH is under control of both E_2 and inhibin. In combination, both of these hormones can function synergistically to suppress FSH secretion in ewes [43, 44]. It has also been suggested that inhibin and E_2 function at different times during the estrous cycle to regulate secretion of FSH in ewes [23]. Concentrations of plasma FSH during the follicular phase of the ovine estrous cycle could be regulated primarily by E_2 ; however, during the 24-h period following the preovulatory surge of LH, E_2 may be relatively unimportant with regard to modulation of FSH secretion [45].

Interestingly, cows in both the OVX and OVXE groups had profiles of FSH secretion with distinct pulses compared to cows in the CONT group. This difference in pulsatile secretion of FSH is probably due to the absence of inhibin or other factors following ovariectomy. Previous research from our laboratory [46] has indicated that release of FSH in intact cows is not pulsatile, but the baseline of FSH secretion may be altered during specific days of the estrous cycle. According to results from the present experiment, E_2 does not affect mean concentration of FSH (because there was no difference in OVXE and OVX groups); however, it does affect the pattern of FSH secretion. Pulse frequency of FSH was enhanced by E_2 in the absence of other ovarian factors without increasing mean concentration of FSH. It is possible that E_2 acts at the gonadotroph to regulate release of FSH as well as LH.

In most animal systems, the amount of mRNA for FSH β subunit and the secretion of FSH are tightly coupled [47]. Unlike the common α or LH β subunit mRNA, FSH β subunit mRNA is regulated by ovarian factors other than steroids. Progesterone and E_2 decrease steady-state amounts of FSH β subunit mRNA coordinately with FSH secretion in ovine pituitary cultures [48]. Amounts of FSH β subunit mRNA in the pituitary are suppressed by administration of follicular fluid or inhibin [47]. Perfusion of pituitaries with inhibin suppresses, while perfusion with activin or LHRH increases, the amount of mRNA for FSH β subunit [49]. It has been postulated that activin and inhibin alter the amount of mRNA for FSH β subunit by modulating production of mRNA at the pituitary. In contrast, E_2 is hypothesized to elicit actions at the hypothalamus through altering LHRH secretion, which causes suppression of mRNA for FSH β subunit in ewes [47]. The relationship between the amount of mRNA for FSH β subunit present in the pituitary to the amount of FSH released from the pituitary of cows has not been evaluated.

In the present experiment, amounts of mRNA for FSH β subunit were lowest in the CONT group; however, administration of E_2 reduced amounts of mRNA in the OVXE group as well, compared to amounts in the OVX group. These results are similar to those in rats, where treatment with E_2 at the time of ovariectomy reduced mRNA for FSH β subunit by up to 67% when compared to that of ovariectomized rats not treated with E_2 [50]. Estradiol, therefore, has an important role in modulating amounts of mRNA for FSH β sub-

unit, which is independent of other ovarian hormones (such as inhibin) in the cow. In the present study, E_2 had little effect on mean concentrations of FSH (OVX compared to OVXE), but it reduced amounts of mRNA for FSH β subunit independent of other ovarian factors.

The actions of E_2 may be elicited post-transcriptionally instead of through regulation of hypothalamic factors by altering the half-life of the mRNA for gonadotropin subunits in the pituitary. Steroid hormones may synergize with ovarian peptide hormones to alter post-transcriptional processing (through stabilization or destabilization). Activin stabilizes mRNA for FSH β subunit, and inhibin decreases stability of mRNA for FSH β subunit [48]. Data from the present experiment indicates that E_2 can affect amount of mRNA for all gonadotropin subunits. Whether this is through transcriptional or post-transcriptional processes still remains to be determined.

Pituitary content of either LH or FSH was not different in any of the three treatment groups. This was presumably due to the relatively short duration of treatment (40 h) compared to that in previous studies. In addition, pituitaries were collected before the preovulatory surge of gonadotropins. Considering both of these factors, secretion of hormones (LH and FSH) did not deplete the pituitary to the extent that differences in stores of the gonadotropins were detected. Collection of pituitaries at earlier time periods (12 or 24 h after PGF $_{2\alpha}$ treatment) may have resulted in different amounts of mRNA for gonadotropin subunits than we observed in the present experiment with pituitary collection at 40 h after PGF $_{2\alpha}$ treatment. The purpose of collecting pituitaries at 40 h after PGF $_{2\alpha}$ was to evaluate amounts of mRNA for gonadotropin subunits at a time when the pituitary had been exposed to E_2 for maximal amounts of time yet before initiation of the preovulatory surge of gonadotropins.

We reject our hypothesis because treatment with E_2 , in a pattern typical of what occurs during the follicular phase of the estrous cycle, did not result in enhanced amounts of mRNA for the gonadotropins. Instead, cows of the OVXE group had amounts of mRNA for α and LH β subunits similar to amounts in cows of the CONT group, whereas cows of the OVX group not treated with E_2 had greater amounts of mRNA for all three subunits than cows of the CONT group. This occurred at the same time that secretion of LH and FSH was greater in cows of the OVX and OVXE groups than in cows of the CONT group. An increased amplitude of LH pulses released from the pituitary results from treating cows with E_2 . This results in greater concentrations of LH in circulation. The amount of mRNA for LH subunits is decreased as a result of treatment of cows with E_2 . Estradiol, therefore, enhances LH release but suppresses amount of mRNA for LH synthesis in cows. Therefore, during the follicular phase of the bovine estrous cycle when secretion of LH is enhanced, there is not a corresponding increase in amount of mRNA for the gonadotropin subunits in cows.

ACKNOWLEDGMENTS

We thank Karl Moline, Jeff Bergman, and Bob Browleit for management of experimental animals; Ken Pearson for assistance with hormone analyses; Georgette Caddy for assistance with isolation of mRNA from pituitaries. We also thank each of the following for materials that allowed us to complete our study: Dr. Debora Hamernik for preparing the FSH β probe and reagents to conduct FSH β slot blots; Dr. Rich Maurer for providing bovine cDNAs for α , LH β , and FSH β subunit mRNA; Dr. Jerry Reeves for LH antisera; Dr. Leo Reichert, Jr. for purified LH and FSH; Dr. Jim Dias for FSH antisera; Dr. Norman Mason for E₂ antisera; and Dr. H. Edward Grotjan, Jr. for the Four Fit program for E₂ analysis.

REFERENCES

- Imakawa K, Day ML, Zalesky DD, Garcia-Winder M, Kittok RJ, Kinder JE. Regulation of pulsatile LH secretion by ovarian steroids in the heifer. *J Anim Sci* 1986; 63:162-168.
- Stumpf TT, Day ML, Wolfe MW, Clutter AC, Stotts JA, Wolfe PL, Kittok RJ, Kinder JE. Effect of estradiol on secretion of luteinizing hormone during the follicular phase of the bovine estrous cycle. *Biol Reprod* 1989; 41:91-97.
- Moenter SM, Caraty A, Karsch FJ. The estradiol-induced surge of gonadotropin-releasing hormone in the ewe. *Endocrinology* 1990; 127:1375-1384.
- Crowder ME, Nett TM. Pituitary content of gonadotropins and receptors for gonadotropin-releasing hormone (GnRH) and hypothalamic content of GnRH during the periovulatory period of the ewe. *Endocrinology* 1984; 114:234-239.
- Gregg DW, Nett TM. Direct effects of estradiol-17 β on the number of gonadotropin-releasing hormone receptors in the ovine pituitary. *Biol Reprod* 1989; 40:288-293.
- Clayton RN, Solano AR, Garcia-Vela A, Dufau ML, Catt KJ. Regulation of pituitary receptors for gonadotropin-releasing hormone during the rat estrous cycle. *Endocrinology* 1980; 107:699-705.
- Katt JA, Duncan JA, Herbon L, Barkan A, Marshall JC. The frequency of GnRH stimulation determines the number of pituitary GnRH receptors. *Endocrinology* 1985; 116:2113-2115.
- Hamernik DL, Crowder ME, Nilson JH, Nett TM. Measurement of messenger ribonucleic acid for luteinizing hormone beta, alpha subunit, growth hormone and prolactin after hypothalamic pituitary disconnection in ovariectomized ewes. *Endocrinology* 1986; 119:2704-2710.
- Andrews WV, Maurer RA, Conn PM. Stimulation of rat luteinizing hormone beta messenger RNA levels by gonadotropin releasing hormone: apparent role for protein kinase C. *J Biol Chem* 1988; 263:13755-13761.
- Rodin DA, Lalozz MR, Clayton RN. Gonadotropin-releasing hormone regulates follicle-stimulating hormone β -subunit gene expression in the male rat. *Endocrinology* 1989; 125:1282-1289.
- Mercer JE, Clarke IJ. Regulation of anterior pituitary gonadotrophin subunit mRNA levels by gonadotrophin-releasing hormone in the ewe. *J Neuroendocrinol* 1989; 1:327-331.
- Shupnik MA. Effect of gonadotropin-releasing hormone on rat gonadotropin gene transcription in vitro: requirement for pulsatile administration for luteinizing hormone- β gene stimulation. *Mol Endocrinol* 1990; 4:1444-1450.
- Haisenleder DJ, Dalkin AC, Ortolano GA, Marshall JC, Shupnik MA. A pulsatile gonadotropin-releasing hormone stimulus is required to increase transcription of the gonadotropin subunit genes: evidence for differential regulation of transcription by pulse frequency in vivo. *Endocrinology* 1991; 128:509-517.
- Bourguignon JP, Franchimont P. Puberty-related increase in episodic LHRH release from rat hypothalamus in vitro. *Endocrinology* 1984; 114:1941-1943.
- Reame N, Sauder SE, Kelch RP, Marshall JC. Pulsatile gonadotropin secretion during the human menstrual cycle: evidence for altered frequency of GnRH secretion. *J Clin Endocrinol & Metab* 1984; 59:328-337.
- Norman RL, Linstrom SA, Bansberg D, Ellinwood WE, Gliessman P, Spies HG. Pulsatile secretion of LH during the menstrual cycle of rhesus macaques. *Endocrinology* 1984; 115:261-266.
- Filicori M, Santoro N, Merriam GR, Crowley WF. Characterization of the physiological pattern of gonadotropin secretion throughout the human menstrual cycle. *J Clin Endocrinol & Metab* 1986; 62:1136-1146.
- Hauger RL, Karsch FJ, Foster DL. A new concept for control of the estrous cycle of the ewe based on the temporal relationships between LH, estradiol, and progesterone in peripheral serum and evidence that progesterone inhibits tonic LH secretion. *Endocrinology* 1977; 101:807-817.
- Moenter SM, Caraty A, Locatelli A, Karsch FJ. Pattern of gonadotropin-releasing hormone (GnRH) secretion leading up to ovulation in the ewe: existence of a preovulatory GnRH surge. *Endocrinology* 1991; 129:1175-1182.
- Legan SJ, Coon GA, Karsch FJ. Role of estrogen as initiator of daily LH surges in the ovariectomized rat. *Endocrinology* 1975; 96:50-56.
- Leung K, Kim KE, Maurer RA, Landefeld TD. Divergent changes in the concentrations of gonadotropin β -subunit messenger ribonucleic acid during the estrous cycle of sheep. *Mol Endocrinol* 1988; 2:272-276.
- Nilson JH, Nejedlik MT, Virgin JB, Crowder ME, Nett TM. Expression of alpha subunit and luteinizing hormone beta genes in the ovine anterior pituitary. Estradiol suppresses accumulation of mRNAs for both alpha subunit and luteinizing hormone beta. *J Biol Chem* 1983; 258:12087-12090.
- Mercer JE, Clements J, Funder JW, Clarke IJ. Regulation of follicle-stimulating hormone β and common α subunit messenger ribonucleic acid by gonadotropin releasing hormone and estrogen in the sheep pituitary. *Neuroendocrinology* 1989; 50:321-326.
- Adams TE, Kinder JE, Chakraborty PK, Estergreen VL, Reeves JJ. Ewe luteal function influenced by pulsatile administration of synthetic LHRH/FSHRH. *Endocrinology* 1975; 97:1460-1467.
- Wolfe MW, Stumpf TT, Roberson MS, Wolfe PL, Kittok RJ, Kinder JE. Estradiol influences on pattern of gonadotropin secretion in bovine males during the period of changed responses to estradiol feedback in age matched females. *Biol Reprod* 1989; 41:626-634.
- Acosta B, Tarnavesky GK, Platt TE, Hamernik DL, Brown JL, Schoenemann HM, Reeves JJ. Nursing enhances the negative effect of estrogen on LH release in the cow. *J Anim Sci* 1983; 57:1530-1536.
- Kojima FN, Stumpf TT, Cupp AS, Werth LA, Roberson MS, Wolfe MW, Kittok RJ, Kinder JE. Exogenous progesterone and progestins as used in estrous synchrony regimens do not mimic the corpus luteum in regulation of luteinizing hormone and 17 β -estradiol in circulation of cows. *Biol Reprod* 1992; 47:1009-1017.
- Roberson MS, Wolfe MW, Stumpf TT, Hamernik DL, Cupp AS, Werth LA, Kojima N, Kittok RJ, Grotjan HE, Kinder JE. Steady-state amounts of α - and luteinizing hormone (LH) β -subunit messenger ribonucleic acids are uncoupled from pulsatility of LH secretion during sexual maturation of the heifer. *Biol Reprod* 1992; 46:435-441.
- Roberson MS, Wolfe MW, Stumpf TT, Kittok RJ, Kinder JE. Luteinizing hormone secretion and corpus luteum function in cows receiving two levels of progesterone. *Biol Reprod* 1989; 41:997-1003.
- Kojima FN, Cupp AS, Stumpf TT, Zalesky DD, Roberson MS, Werth LA, Wolfe MW, Kittok RJ, Grotjan HE, Kinder JE. Effect of 17 β -estradiol on distribution of pituitary isoforms of luteinizing hormone and follicle-stimulating hormone during the follicular phase of the bovine estrous cycle. *Biol Reprod* 1995; 52:297-304.
- Erwin CR, Croyle ML, Donelson JE, Maurer RA. Nucleotide sequence of cloned complementary deoxyribonucleic acid for the alpha subunit of bovine pituitary glycoprotein hormones. *Biochemistry* 1983; 22:4856-4860.
- Maurer RA. Analysis of several bovine lutropin beta-subunit cDNAs reveals heterogeneity of nucleotide sequence. *J Biol Chem* 1985; 260:4684-4687.
- Sambrook J, Fritsch EF, Maniatis T. Molecular cloning. A Laboratory Manual. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press; 1989: 753.
- Kim KE, Gordon DF, Maurer RA. Nucleotide sequence of the bovine gene for follicle-stimulating hormone β -subunit. *DNA* 1988; 7:227-233.
- Feinberg A, Vogelstein B. A technique for p DNA restriction endonuclease fragments to high specific activity. *Anal Biochem* 1983; 132:6-13.
- SAS. SAS User's Guide, Statistics. Cary, NC: Statistical Analysis System Institute, Inc.; 1985.
- Steele RGD, Torrie JH. Principles and Procedures of Statistics. New York: McGraw Hill Book Co.; 1980.
- Alexander DC, Miller WL. Regulation of ovine follicle stimulating hormone β chain mRNA by 17 β -estradiol in vivo and in vitro. *J Biol Chem* 1982; 257:2282-2286.
- Landefeld TD, Kepa J, Karsch FJ. Estradiol feedback effects on the alpha subunit mRNA in sheep pituitary glands: correlation with serum and pituitary luteinizing hormone concentrations. *Proc Natl Acad Sci USA* 1984; 81:1322-1326.
- Shupnik MA, Gharib SD, Chin WW. Estrogen suppresses rat gonadotropin gene transcription in vivo. *Endocrinology* 1988; 122:1842-1846.
- Di Gregorio GB, Kile JP, Herring RH, Nett TM. More rapid restoration of pituitary content of follicle-stimulating hormone than of luteinizing hormone after depletion by oestradiol-17 β in ewes. *J Reprod Fertil* 1991; 93:347-354.
- Ying SY. Inhibins, activins and follistatins: gonadal protein modulating the secretion of follicle-stimulating hormone. *Endocr Rev* 1988; 9:267-293.
- Mann GE, Campbell BK, McNeilly AS, Baird DT. The role of inhibin and oestradiol in the control of FSH secretion in the sheep. *J Endocrinol* 1992; 133:381-391.
- Campbell BK, Mann GE, McNeilly AS, Baird DT. The pattern of ovarian inhibin, estradiol, and androstenedione secretion during the estrous cycle of the ewe. *Endocrinology* 1990; 127:227-235.

45. Findlay JK, Quigg H, Juhola P, Katsahambas S, Clarke IJ, Doughton B, Robertson DM. Immunoreactive inhibin concentrations in sheep plasma during the ovarian cycle. *Endocrinology* 1988; Abstract 563.
46. Cupp AS, Stumpf TT, Kojima FN, Werth LA, Wolfe MW, Roberson MS, Kittok RJ, Kinder JE. Secretion of gonadotrophins change during the luteal phase of the bovine oestrous cycle in the absence of corresponding changes in progesterone or 17 β -oestradiol. *Anim Reprod Sci* 1994; (in press).
47. Mercer JE, Clements JA, Funder JW, Clarke IJ. Rapid and specific lowering of pituitary FSH β mRNA levels by inhibin. *Mol Cell Endocrinol* 1987; 53:251–254.
48. Phillips CL, Lin LW, Wu JC, Guzman K, Milsted A, Miller WL. 17 β -estradiol and progesterone inhibit transcription of the genes encoding the subunit of ovine-follicle-stimulating hormone. *Mol Endocrinol* 1988; 2:641–649.
49. Weiss J, Crowley WF Jr, Halvorson LM, Jameson JL. Perfusion of rat pituitary cells with gonadotropin-releasing hormone, activin, and inhibin reveals distinct effects on gonadotropin gene expression and secretion. *Endocrinology* 1993; 132:2307–2311.
50. Dalkin AC, Haisenleder DJ, Ortolano GA, Suhr A, Marshall JC. Gonadal regulation of gonadotropin subunit gene expression: evidence for regulation of follicle-stimulating hormone- β messenger ribonucleic acid by nonsteroidal hormones in female rats. *Endocrinology* 1990; 127:798–806.