

2-2009

## *Rathayibacter iranicus* isolated from symptomless wheat seeds in Turkey

E. Postnikova  
USDA-ARS, Ft. Detrick, MD

Irina V. Agarkova  
University of Nebraska - Lincoln, iagarkova2@unl.edu

S. Altundag  
Plant Protection Research Institute, Ankara, Turkey

F. Eskandari  
USDA-ARS, Ft. Detrick, MD

A. Sechler  
USDA-ARS, Ft. Detrick, MD

*See next page for additional authors*

Follow this and additional works at: <http://digitalcommons.unl.edu/plantpathpapers>

 Part of the [Plant Pathology Commons](#)

---

Postnikova, E.; Agarkova, Irina V.; Altundag, S.; Eskandari, F.; Sechler, A.; Karahan, A.; Vidaver, Anne M.; Schneider, W.; Ozakman, M.; and Schaad, N. W., "*Rathayibacter iranicus* isolated from symptomless wheat seeds in Turkey" (2009). *Papers in Plant Pathology*. 193.

<http://digitalcommons.unl.edu/plantpathpapers/193>

This Article is brought to you for free and open access by the Plant Pathology Department at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Papers in Plant Pathology by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

---

**Authors**

E. Postnikova, Irina V. Agarkova, S. Altundag, F. Eskandari, A. Sechler, A. Karahan, Anne M. Vidaver, W. Schneider, M. Ozakman, and N. W. Schaad

## **Rathayibacter iranica** isolated from symptomless wheat seeds in Turkey

E. Postnikova<sup>a</sup>, I. Agarkova<sup>b</sup>, S. Altundag<sup>c</sup>, F. Eskandari<sup>a</sup>, A. Sechler<sup>a</sup>, A. Karahan<sup>c</sup>, A. K. Vidaver<sup>b</sup>, W. Schneider<sup>a</sup>, M. Ozakman<sup>c</sup> and N. W. Schaad<sup>1a\*</sup>

<sup>a</sup>ARS-USDA, Foreign Disease-Weed Science Research Unit, Ft. Detrick, Maryland; <sup>b</sup>Department of Plant Pathology, University of Nebraska, Lincoln, Nebraska, USA; and <sup>c</sup>Plant Protection Research Institute, P.O. Box 49, Ankara, Turkey

*Rathayibacter iranica* (Ri), originally reported in Iran in 1961 (Sharif, 1961), has not been reported outside Iran and only one strain is known to exist. Like *R. tritici* (Rt), Ri causes a gumming disease of wheat in association with the nematode *Anguina tritici* (Paruthi *et al.*, 1989). During 2003, a survey of wheat seed for *Rathayibacter* species (RS) in Turkey using samples from 799 farmers in six provinces in Central Anatolia was conducted. The samples showed neither the brown to black galls typical of *A. tritici* infection nor the yellowish galls typical of Ri and Rt. To determine the presence of RS, 120 g samples were washed in 100 mL of sterile 0.85% NaCl with 0.02% Tween 20 and plated onto 523 agar (Schaad *et al.*, 2001) modified by adding 30 mg nalidixic acid, 15 mg polymixin B sulphate and 100 mg cycloheximide per litre. After incubation at 27°C for seven days, several typical colonies of RS were cloned by streaking onto YDC agar (Schaad *et al.*, 2001). Twenty five strains were presumptively identified as RS based on a Gram positive reaction, oxidase negative reaction and yellow growth.

Three strains, TRS2 from Konya, TRS10 from Hisarkikaya, and TRS25 from Ankara, were used for further identification and speciation. The other 22 strains were archived for later studies. The type strains of Rt, (International Collection of Phytopathogenic Bacteria [ICPB] 70004<sup>1</sup>; FH-5; CT 102<sup>1</sup>) and Ri (ICPB 70005<sup>1</sup>; FH-6<sup>1</sup>; CI 148<sup>1</sup>), were included as controls. All three strains were biochemically typical of Ri and Rt by producing acid from D-mannose, inulin, galactose, and mannitol. However, the strains failed to utilize acetate, or to hydrolyze casein, failed to grow in greater than 1% sodium chloride amended NBY, or on CNS

agar, which is characteristic of Ri but not Rt (Davis & Vidaver, 2001). The new strains contained the fatty acids iso-14:0 and 16:0; anteiso 15:0 and 15:1, and 17:0, and 16:0, also typical of Ri. Identification of the three strains as Ri was further supported by 16s rDNA sequence similarities to Ri and Rt of 100 and 99.3%, respectively (GenBank Accession Nos. FJ595101 - Ri, ICPB 70005 and FJ595102 - Rt, ICPB 70004). Identification as Ri was confirmed by AFLP analysis (EcoRI + 0 and MseI + C) showing a pattern of 90% similarity to Ri but only 46% to Rt. This is the first report of Ri outside of Iran. Cultures have been deposited in the ICPB at Ft. Detrick, MD, USA, as ICPB 70146 (FH-154; TRS2), 70 154 (FH-162; TRS10), and 70 169 (FH-177; TRS25).

### References

- Davis MJ, Vidaver AK, 2001. Gram positive bacteria, coryneform plant pathogens. In: Schaad NW, Jones JB, Chun W, eds. *Laboratory Guide for Identification of Plant Pathogenic Bacteria*, 3rd edn. St. Paul, MN, USA: APS Press, 218–34.
- Paruthi IJ, Bajaj HK, Bhatti DS, 1989. Further observations on *Anguina tritici* (Nematoda) and *Corynebacterium michiganense* pv. *tritici* causing earcockle and yellow ear rot in wheat. *Nematologica* 35, 491–3.
- Schaad NW, Jones JB, Chun W, eds, 2001. *Laboratory Guide for Identification of Plant Pathogenic Bacteria*, 3rd edn. St. Paul, MN, USA: APS Press.
- Scharif G, 1961. *Corynebacterium iranica* sp. nov. on wheat (*Triticum vulgare* L.) in Iran, and a comparative study of it with *C. tritici* and *C. rathayi*. *Entomologie et Phytopathologie Appliqués* 19, 1–24.

\*E-mail: norman.schaad@ars.usda.gov. Accepted 18 February 2009 at [www.bspp.org.uk/ndr](http://www.bspp.org.uk/ndr) where figures relating to this paper can be viewed.

## **Pyrofomes demidoffii** newly reported to cause a white trunk rot of juniper (*Juniperus formosana*) in China

Y. C. Dai<sup>1a\*</sup> and X. S. He<sup>b</sup>

<sup>a</sup>Institute of Applied Ecology, Chinese Academy of Sciences, Shenyang 110016; and <sup>b</sup>Southwest University of Science and Technology, Mianyang 621010, China

*Juniperus formosana* has a wide distribution in China, and it is an important conifer tree in central and southwestern China. In the summer of 2006, a typical white trunk rot of the tree was observed at Wolong area (30°50'N, 103°10'E), Dujiangyan County, central Sichuan Province, southwest China. Twelve trees were affected from 85 examined, and showed signs of advanced internal decay of the trunk.

A perennial, unguulate and woody polypore was found on the trunk of affected trees. Fruiting bodies were 12–15 cm wide and 14 cm thick at the base. The upper surface was greyish black, the poroid surface bright yellow to orange brown with densely packed tubes (around 3–4 per mm). The inside of the fruiting bodies was cinnamon to brick-red and contained dimitic hyphae; the generative hyphae had clamp connections. Basidiospores were ellipsoid, truncate, thick-walled, pale brownish, slightly dextrinoid and cyanophilous, 6–8 × 5–6 µm. These characteristics agree with the published description of *Pyrofomes demidoffii* (Ryvarden & Gilbertson, 1994) and match specimens examined from the Herbarium of the Botanical Museum of the University of Helsinki (H). Voucher specimens have been deposited at the Institute of Applied Ecology, Chinese Academy of Sciences (IFP).

Previously *P. demidoffii* has been recorded on junipers in Central Europe, Africa and North America (Ryvarden & Gilbertson, 1994). It

grows exclusively on *Juniperus* spp. and causes a white trunk rot of living trees. The fungus causes major decay of junipers in western North America (Scharpf, 1993) and Pakistan (Zakaullah, 1978). The fungus was not found in previous surveys (Dai *et al.*, 2007) and this is the first report of *P. demidoffii* on *Juniperus formosana* from China.

### Acknowledgement

The work is supported by the Ministry of Science and Technology of China (Project No. 007AA021506).

### References

- Dai YC, Cui BK, Yuan HS, Li BD, 2007. Pathogenic wood-decaying fungi in China. *Forest Pathology* 37, 105–20.
- Ryvarden L, Gilbertson RL, 1994. European polypores 2. *Synopsis Fungorum* 7, 394–743.
- Scharpf, RF, 1993. *Diseases of Pacific Coast Conifers*. Albany, OR, USA: U.S. Forest Service, Pacific Southwest Research Station.
- Zakaullah, 1978. Decay in the Ziarat juniper forests of Baluchistan. *Pakistan Journal of Forestry* 28, 28–34.

\*E-mail: yuchengd@yahoo.com. Accepted 18 February 2009 at [www.bspp.org.uk/ndr](http://www.bspp.org.uk/ndr) where figures relating to this paper can be viewed.