

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Nebraska Bird Review

Nebraska Ornithologists' Union

12-1998

Endemicity and Regional Biodiversity in Nebraska's Breeding Avifauna

Paul A. Johnsgard

University of Nebraska-Lincoln, pajohnsgard@gmail.com

Follow this and additional works at: <http://digitalcommons.unl.edu/nebbirdrev>



Part of the [Poultry or Avian Science Commons](#), and the [Zoology Commons](#)

Johnsgard, Paul A., "Endemicity and Regional Biodiversity in Nebraska's Breeding Avifauna" (1998). *Nebraska Bird Review*. 203.
<http://digitalcommons.unl.edu/nebbirdrev/203>

This Article is brought to you for free and open access by the Nebraska Ornithologists' Union at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Nebraska Bird Review by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Endemicity and Regional Biodiversity in Nebraska's Breeding Avifauna
by Paul A. Johnsgard (School of Biological Sciences, U. of Nebraska-Lincoln)

I estimate that 215 bird species currently breed or have previously bred in Nebraska. This number compares with a total of 330 species that breed or have bred in the Great Plains region south of Canada, as I defined that region in my book on the breeding birds of the Great Plains (Johnsgard, 1979).

Grasslands or potential grasslands make up about 81 percent of the Great Plains' vegetation, and probably represented about 96 percent of Nebraska's original vegetation. Only in the Sandhills region is that vegetation type still essentially intact; the tall-grass prairie of eastern Nebraska is virtually gone. In my 1979 summary, I judged that 36 species of Great Plains birds are grassland-adapted forms, with 15 of these endemic to the Great Plains, and the rest more broadly distributed. Nine of these endemics breed in Nebraska (Greater Prairie-Chicken, Mountain Plover, Long-billed Curlew, Upland Sandpiper, Dickcissel, Lark Bunting, Clay-colored Sparrow, Cassin's Sparrow, McCown's Longspur and Chestnut-collared Longspur), and the Chihuahuan Raven also once did. None of these ten species is considered endangered nationally, although the Greater Prairie-Chicken needs to be monitored closely. Birds of the tall-grass prairie that currently breed in eastern Nebraska include the Upland Sandpiper, Dickcissel, Eastern Meadowlark, Field Sparrow and Henslow's Sparrow. Both of the latter are not, strictly speaking, prairie species, but either represent a later successional stage containing some small trees (Field Sparrow), or the presence of some brushy or weedy elements (Henslow's Sparrow). In the west the short-grass plains support several declining or rare species, including Ferruginous Hawk, Swainson's Hawk,

Prairie Falcon, Mountain Plover, Burrowing Owl, Common Poor-will, Lark Bunting, and two longspurs. Some of these also have certain additional non-vegetative needs such as cliff-side nest sites (Prairie Falcon), burrows (Burrowing Owl), nearly bare ground (Mountain Plover), or rocky substrates (Poor-will). The geographically and ecologically intermediate Sandhills prairies support many typical grassland and wet meadow breeders, such as Sharp-tailed Grouse, Long-billed Curlew, Bobolink, Lark Sparrow, Grasshopper Sparrow and Horned Lark.

Woodlands and forests comprise about 15 percent of the native vegetation of the Great Plains states, and support some 51 percent of the region's breeding avifauna (Johnsgard, 1979). I estimate that in Nebraska this vegetation type occupies three percent of the state's surface area, and supports 48 percent of its avifauna. It is thus one of the most important vegetation types for generating species diversity of our avifauna. Bald Eagles are among the rare woodland-nesting species of Nebraska, although they are also strongly water-dependent for food. Several hawks (Broad-winged, Red-shouldered, Cooper's, etc.) are other clear associates of mature woodlands. Several woodland nesting passerines that have generally large ranges elsewhere in the central or western states are limited to woodlands in extreme northwestern Nebraska's Pine Ridge region, or in the lower Missouri woodlands of extreme southeastern Nebraska (see Table 2).

Wetlands (including rivers, lakes, large reservoirs, etc.) cover only about one percent of the surface area of the Great Plains states, but support 21 percent of its breeding avifauna, thus contributing more to the region's avifauna diversity than any other habitat type relative to its area (Johnsgard, 1979). Similarly, surface water covers only about one percent of Nebraska's surface, but such areas support 27 percent of our breeding avifauna. Rare or localized species that depend on such wetlands in Nebraska include the Trumpeter Swan, Black-necked Stilt, White-faced Ibis, Least Tern and Piping Plover. All of the rails are marsh dependent and thus rather localized; but the breeding status of the Black Rail in Nebraska is still unknown. The Northern Harrier also nests in marshes, but forages over meadows and grasslands. The Sandhill Crane once bred in Nebraska's Sandhill marshes, and may again return as a breeding species, just as the Bald Eagle has done. Wetlands also provide extremely important migratory habitats in Nebraska for many species of shorebirds, waterfowl, and cranes, including the endangered Whooping Crane.

Methods

As part of the ongoing national interest in defining Important Bird Areas, I decided that an analysis of my distribution maps of the breeding ranges of Nebraska's avifauna would be of value. These maps, originally drawn for my book on the breeding birds of the Great Plains (Johnsgard, 1979) have been periodically updated; the most recent versions appear in my summary of Nebraska's avifauna (Johnsgard, 1998). Although I used my breeding-distribution maps for the Great Plains as my data source, I accepted the fact that the range limits indicated there are usually little more than "informed guesses," as we still lack highly detailed breeding range maps for most Nebraska species. When the Breeding Bird Atlas for Nebraska is finally published, we will have better mapping possibilities. My maps tend to be "generous" as to probable breeding occurrence; thus the Northern Harrier and Long-eared Owl were both mapped as occurring throughout the entire

state, even though they respectively nest only in areas having available marshy or mature woodland breeding habitats. I prepared a transparent overlay map of the state, using the 68 quadrants of latitude and longitude that were adopted in the *Nebraska Atlas and Gazetteer*, one of a series of state atlases published by DeLorme Co., of Freeport, Maine.

Using these standardized quadrants makes for easy reference by others, and might facilitate future between-state comparisons as well. This atlas series' quadrants in Nebraska measure 30.7 miles (49.4 kilometers) in east-west distance and 44 miles (70.8 kilometers) in north-south distance, or 1,351 square miles (3,497 sq. kilometers) per quadrant. Each of my range maps was then compared with the quadrant overlay, and those all quadrants falling within the species' indicated range were tallied (Figure 1). The total number of species falling within each quadrant was then determined. Many of the quadrants along the southwestern and eastern borders of the state are ones whose areas extend well beyond the state's legal boundaries (e.g., Nos. 10-13, 27, 41-45, 56, 68). A few species were tallied as "present" if they occurred within the limits of such quadrants although these species' apparent actual range limits failed to reach Nebraska's boundaries. Treated otherwise, projected species totals for these variably smaller border quadrants could not be numerically compared with numbers for those quadrants falling entirely within the state's boundaries. A total of 51 species were judged to be pandemic breeders throughout the state (Table 1), all the remaining 164 species were judged absent from one or more quadrants. At the other extreme are those species occurring in as few as a single quadrant; these are of special interest in terms of defining regional endemism. Such species, which are clearly localized in Nebraska but may not be rare elsewhere in their ranges, are listed as regional endemics in Table 2.

Results

Ranges in estimated species diversity varied from as few as 96 species (quadrant 43) to a maximum of 140 species (quadrant 41). Part of this apparent variation reflects regional differences in data bases; thus quadrant 43 is located in the southwestern Panhandle, where few ornithologists have studied local birds, whereas quadrant 41 includes the Douglas County area, with its long history of ornithological investigations. However, several trends are clearly visible. The generally highest levels of species diversity occur along the lower Missouri River

TABLE 1

Estimated Number of Breeding Species in 68 Nebraska Quadrants

1 128	2 115	3 105	4 106	5 105	6 112	7 122	8 125	9 124	10 125	11 124	12 125	13 128		
14 118	15 107	16 111	17 103	18 101	19 108	20 109	21 111	22 112	23 104	24 108	25 108	26 124	27 127	
28 109	29 104	30 112	31 109	32 111	33 112	34 114	35 113	36 108	37 104	38 113	39 112	40 117	41 140	
42 98	43 96	44 100	45 103	46 105	47 115	48 110	49 114	50 110	51 115	52 113	53 97	54 118	55 133	56 138
57 101	58 97	59 98	60 100	61 100	62 100	63 99	64 102	65 97	66 108	67 113	68 138			

valley, from the Kansas border north to the Omaha area. Numbers gradually tail off from that area up the Missouri and Niobrara valleys to about Cherry County, where there is apparently a sharp decline (quadrant 6), roughly where the mixed deciduous forest changes to a ponderosa pine forest. There is a second center of breeding biodiversity centered in Sioux County (quadrant 1), tapering off in Dawes and Scottsbluff counties (quadrants 2 & 14). The Platte Valley represents a corridor of moderate diversity, but the Republican River Valley along the lowest tier of quadrants is distinctly less diverse in its breeding avifauna than are other regions. The Sandhills region shows only slightly less diversity than the Platte Valley, probably because of the substantial number of wetland species breeding with the region.

There are some ways of comparing these hypothetical estimates of breeding diversity with actual estimates. Thus, quadrant 32 includes the Cedar Point Biological Station area, where 111 species were judged as likely breeders. The known total number of breeding species as of 1996 was 104 (Brown et al., 1996). Quadrant 19, with 108 prospective breeding species includes Valentine National Wildlife Refuge, which has at least 93 known breeders. The Lancaster County list of breeding birds includes at least 103 species (personal determination), as compared with an estimated total of 118 species for all of quadrant 54. These numbers suggest that the margin of overestimates for these projected figures may average only about ten percent, and should be less if the entire quadrant area were carefully surveyed.

Discussion

Important Bird Areas have been defined as those that (1) support threatened or endangered species, (2) support rare or endemic species, or species of special conservation concern, (3) include species associated with rare or threatened habitats, (4) represent sites supporting areas of high numerical population abundance or species diversity, or (5) those where long-term monitoring has occurred. The data presented here offers direct information on points two and four. Information on the distribution of rare and endangered species, as well as of endangered habitats, is available through state and federal agencies, the Nature Conservancy, and similar conservation groups. Long-term monitoring of bird populations has gone on only in only a few sites, most obviously some of the state's major national wildlife refuges and Cedar Point Biological Station, all of which are obvious candidates for inclusion.

Thus regions that should be examined for the identification of important bird areas on the basis of unusually high species diversity should include the lower Missouri Valley (e.g., Indian Cave State Park, Fontenelle Forest, etc.). It should also include much of the Pine Ridge's already preserved areas. The middle Niobrara River Valley is of special interest as an east-west faunal transition zone (Johnsgard, 1982), and one of the major "suture zones" of overlapping eastern and western biota. The Platte Valley has similar basic biological significance, especially the region of the middle Platte, which additionally supports major populations of threatened Least Terns and Piping Plovers, and critical habitat for the endangered Whooping Crane. The Lake McConaughy area is already known to support one of the highest areas of avian species diversity in the entire country (Brown et al., 1996), and additionally includes Cedar Point Biological Station.

Summary

An analysis of the relative species diversity of Nebraska's breeding avifauna establishes several areas of unusual species richness and endemism, these most important being the Missouri Valley and associated middle to lower Niobrara Valley, the Pine Ridge area of the northwestern Panhandle, and the entire Platte Valley.

References

- Brown, C. R., M. B. Brown, P. A. Johnsgard, J. Kren & W. C. Scharf. 1996. "Birds of the Cedar Point Biological Station Area, Keith and Garden Counties, Nebraska: Seasonal Occurrence and Breeding Data." *Transactions of the Nebraska Academy of Sciences* 29: 91-108.
- Johnsgard, P. A. 1979. *Birds of the Great Plains: Breeding Species and Their Distribution*. Lincoln: Univ. of Nebraska Press.
- _____. 1987. The Ornithogeography of the Great Plains States. *Prairie Naturalist* 10:97-112.
- _____. 1998. *The Birds of Nebraska*. Published by the author, Lincoln, Nebraska.

Table 1

Species of Birds Considered as Pandemic Breeders Across Nebraska

Pied-billed Grebe	Blue Jay
American Bittern	American Crow
Great Blue Heron	House Wren
Black-crowned Night Heron	American Robin
Canada Goose	Gray Catbird
Mallard	Loggerhead Shrike
Northern Pintail	European Starling
Blue-winged Teal	Yellow Warbler
Red-tailed Hawk	Common Yellow-throat
American Kestrel	Blue Grosbeak
Northern Harrier	Chipping Sparrow
American Coot	Lark Sparrow
Killdeer	Grasshopper Sparrow
Rock Dove	Red-winged Blackbird
Mourning Dove	Western Meadowlark
Yellow-billed Cuckoo	Yellow-headed Blackbird
Barn Owl	Common Grackle
Great Horned Owl	Brown-headed Cowbird
Long-eared Owl	American Goldfinch
Common Nighthawk	House Sparrow
Belted Kingfisher	
Downy Woodpecker	
Hairy Woodpecker	
Western Kingbird	
Eastern Kingbird	
Northern Rough-winged Swallow	
Bank Swallow	
Cliff Swallow	
Barn Swallow	

Table 2

Species of Birds Considered to be Endemic to Specific Nebraska RegionsNorthwestern Panhandle Endemics

White-throated Swift
Lewis' Woodpecker
Cordilleran Flycatcher
Plumbeus Vireo
Pinyon Jay
Violet-green Swallow
Brown Creeper
Pygmy Nuthatch
Mountain Bluebird
Townsend's Solitaire
Swainson's Thrush
Sage Thrasher
Western Tanager
Yellow-rumped Warbler
Brewer's Sparrow
Dark-eyed Junco
McCown's Longspur
Chestnut-collared Longspur

Lower Missouri Valley Endemics

Red-shouldered Hawk
Broad-winged Hawk
Chuck-wills-widow
White-eyed Vireo
Carolina Wren
Summer Tanager
Northern Parula
Cerulean Warbler
Prothonotary Warbler
Kentucky Warbler
Yellow-throated Warbler

Southwestern Panhandle Endemics

Mountain Plover
Cassin's Kingbird
Cassin's Sparrow