

1-13-2006

# Search for Neutral Higgs Bosons of the Minimal Supersymmetric Standard Model Decaying to $\tau$ Pairs in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV

A. Abulencia

*University of Illinois, Urbana, Illinois 61801, USA*

Kenneth A. Bloom

*University of Nebraska - Lincoln, kbloom2@unl.edu*

CDF Collaboration

Follow this and additional works at: <http://digitalcommons.unl.edu/physicsbloom>



Part of the [Physics Commons](#)

---

Abulencia, A.; Bloom, Kenneth A.; and Collaboration, CDF, "Search for Neutral Higgs Bosons of the Minimal Supersymmetric Standard Model Decaying to  $\tau$  Pairs in  $p\bar{p}$  Collisions at  $\sqrt{s} = 1.96$  TeV" (2006). *Kenneth Bloom Publications*. 204.  
<http://digitalcommons.unl.edu/physicsbloom/204>

This Article is brought to you for free and open access by the Research Papers in Physics and Astronomy at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Kenneth Bloom Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

## Search for Neutral Higgs Bosons of the Minimal Supersymmetric Standard Model Decaying to $\tau$ Pairs in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV

A. Abulencia,<sup>23</sup> D. Acosta,<sup>17</sup> J. Adelman,<sup>13</sup> T. Affolder,<sup>10</sup> T. Akimoto,<sup>53</sup> M. G. Albrow,<sup>16</sup> D. Ambrose,<sup>16</sup> S. Amerio,<sup>42</sup> D. Amidei,<sup>33</sup> A. Anastassov,<sup>50</sup> K. Anikeev,<sup>16</sup> A. Annovi,<sup>44</sup> J. Antos,<sup>1</sup> M. Aoki,<sup>53</sup> G. Apollinari,<sup>16</sup> J.-F. Arguin,<sup>32</sup> T. Arisawa,<sup>55</sup> A. Artikov,<sup>14</sup> W. Ashmanskas,<sup>16</sup> A. Attal,<sup>8</sup> F. Azfar,<sup>41</sup> P. Azzi-Bacchetta,<sup>42</sup> P. Azzurri,<sup>44</sup> N. Bacchetta,<sup>42</sup> H. Bachacou,<sup>28</sup> W. Badgett,<sup>16</sup> A. Barbaro-Galtieri,<sup>28</sup> V. E. Barnes,<sup>46</sup> B. A. Barnett,<sup>24</sup> S. Baroiant,<sup>7</sup> V. Bartsch,<sup>30</sup> G. Bauer,<sup>31</sup> F. Bedeschi,<sup>44</sup> S. Behari,<sup>24</sup> S. Belforte,<sup>52</sup> G. Bellettini,<sup>44</sup> J. Bellinger,<sup>57</sup> A. Belloni,<sup>31</sup> E. Ben-Haim,<sup>16</sup> D. Benjamin,<sup>15</sup> A. Beretvas,<sup>16</sup> J. Beringer,<sup>28</sup> T. Berry,<sup>29</sup> A. Bhatti,<sup>48</sup> M. Binkley,<sup>16</sup> D. Bisello,<sup>42</sup> M. Bishai,<sup>16</sup> R. E. Blair,<sup>2</sup> C. Blocker,<sup>6</sup> K. Bloom,<sup>33</sup> B. Blumenfeld,<sup>24</sup> A. Bocci,<sup>48</sup> A. Bodek,<sup>47</sup> V. Boisvert,<sup>47</sup> G. Bolla,<sup>46</sup> A. Bolshov,<sup>31</sup> D. Bortoletto,<sup>46</sup> J. Boudreau,<sup>45</sup> S. Bourov,<sup>16</sup> A. Boveia,<sup>10</sup> B. Brau,<sup>10</sup> C. Bromberg,<sup>34</sup> E. Brubaker,<sup>13</sup> J. Budagov,<sup>14</sup> H. S. Budd,<sup>47</sup> S. Budd,<sup>23</sup> K. Burkett,<sup>16</sup> G. Busetto,<sup>42</sup> P. Bussey,<sup>20</sup> K. L. Byrum,<sup>2</sup> S. Cabrera,<sup>15</sup> M. Campanelli,<sup>19</sup> M. Campbell,<sup>33</sup> F. Canelli,<sup>8</sup> A. Canepa,<sup>46</sup> D. Carlsmith,<sup>57</sup> R. Carosi,<sup>44</sup> S. Carron,<sup>15</sup> M. Casarsa,<sup>52</sup> A. Castro,<sup>5</sup> P. Catastini,<sup>44</sup> D. Cauz,<sup>52</sup> M. Cavalli-Sforza,<sup>3</sup> A. Cerri,<sup>28</sup> L. Cerrito,<sup>41</sup> S. H. Chang,<sup>27</sup> J. Chapman,<sup>33</sup> Y. C. Chen,<sup>1</sup> M. Chertok,<sup>7</sup> G. Chiarelli,<sup>44</sup> G. Chlachidze,<sup>14</sup> F. Chlebana,<sup>16</sup> I. Cho,<sup>27</sup> K. Cho,<sup>27</sup> D. Chokheli,<sup>14</sup> J. P. Chou,<sup>21</sup> P. H. Chu,<sup>23</sup> S. H. Chuang,<sup>57</sup> K. Chung,<sup>12</sup> W. H. Chung,<sup>57</sup> Y. S. Chung,<sup>47</sup> M. Ciljak,<sup>44</sup> C. I. Ciobanu,<sup>23</sup> M. A. Ciocci,<sup>44</sup> A. Clark,<sup>19</sup> D. Clark,<sup>6</sup> M. Coca,<sup>15</sup> A. Connolly,<sup>28</sup> M. E. Convery,<sup>48</sup> J. Conway,<sup>7</sup> B. Cooper,<sup>30</sup> K. Copic,<sup>33</sup> M. Cordelli,<sup>18</sup> G. Cortiana,<sup>42</sup> A. Cruz,<sup>17</sup> J. Cuevas,<sup>11</sup> R. Culbertson,<sup>16</sup> D. Cyr,<sup>57</sup> S. DaRonco,<sup>42</sup> S. D'Auria,<sup>20</sup> M. D'Onofrio,<sup>19</sup> D. Dagenhart,<sup>6</sup> P. de Barbaro,<sup>47</sup> S. De Cecco,<sup>49</sup> A. Deisher,<sup>28</sup> G. De Lentdecker,<sup>47</sup> M. Dell'Orso,<sup>44</sup> S. Demers,<sup>47</sup> L. Demortier,<sup>48</sup> J. Deng,<sup>15</sup> M. Deninno,<sup>5</sup> D. De Pedis,<sup>49</sup> P. F. Derwent,<sup>16</sup> C. Dionisi,<sup>49</sup> J. Dittmann,<sup>4</sup> P. DiTuro,<sup>50</sup> C. Dörr,<sup>25</sup> A. Dominguez,<sup>28</sup> S. Donati,<sup>44</sup> M. Donega,<sup>19</sup> P. Dong,<sup>8</sup> J. Donini,<sup>42</sup> T. Dorigo,<sup>42</sup> S. Dube,<sup>50</sup> K. Ebina,<sup>55</sup> J. Efron,<sup>38</sup> J. Ehlers,<sup>19</sup> R. Erbacher,<sup>7</sup> D. Errede,<sup>23</sup> S. Errede,<sup>23</sup> R. Eusebi,<sup>47</sup> H. C. Fang,<sup>28</sup> S. Farrington,<sup>29</sup> I. Fedorko,<sup>44</sup> W. T. Fedorko,<sup>13</sup> R. G. Feild,<sup>58</sup> M. Feindt,<sup>25</sup> J. P. Fernandez,<sup>46</sup> R. Field,<sup>17</sup> G. Flanagan,<sup>34</sup> L. R. Flores-Castillo,<sup>45</sup> A. Foland,<sup>21</sup> S. Forrester,<sup>7</sup> G. W. Foster,<sup>16</sup> M. Franklin,<sup>21</sup> J. C. Freeman,<sup>28</sup> Y. Fujii,<sup>26</sup> I. Furic,<sup>13</sup> A. Gajjar,<sup>29</sup> M. Gallinaro,<sup>48</sup> J. Galyardt,<sup>12</sup> J. E. Garcia,<sup>44</sup> M. Garcia Sciveres,<sup>28</sup> A. F. Garfinkel,<sup>46</sup> C. Gay,<sup>58</sup> H. Gerberich,<sup>23</sup> E. Gerchtein,<sup>12</sup> D. Gerdes,<sup>33</sup> S. Giagu,<sup>49</sup> P. Giannetti,<sup>44</sup> A. Gibson,<sup>28</sup> K. Gibson,<sup>12</sup> C. Ginsburg,<sup>16</sup> K. Giolo,<sup>46</sup> M. Giordani,<sup>52</sup> M. Giunta,<sup>44</sup> G. Giurgiu,<sup>12</sup> V. Glagolev,<sup>14</sup> D. Glenzinski,<sup>16</sup> M. Gold,<sup>36</sup> N. Goldschmidt,<sup>33</sup> J. Goldstein,<sup>41</sup> G. Gomez,<sup>11</sup> G. Gomez-Ceballos,<sup>11</sup> M. Goncharov,<sup>51</sup> O. González,<sup>46</sup> I. Gorelov,<sup>36</sup> A. T. Goshaw,<sup>15</sup> Y. Gotra,<sup>45</sup> K. Goulios,<sup>48</sup> A. Gresele,<sup>42</sup> M. Griffiths,<sup>29</sup> S. Grinstein,<sup>21</sup> C. Grosso-Pilcher,<sup>13</sup> U. Grundler,<sup>23</sup> J. Guimaraes da Costa,<sup>21</sup> C. Haber,<sup>28</sup> S. R. Hahn,<sup>16</sup> K. Hahn,<sup>43</sup> E. Halkiadakis,<sup>47</sup> A. Hamilton,<sup>32</sup> B.-Y. Han,<sup>47</sup> R. Handler,<sup>57</sup> F. Happacher,<sup>18</sup> K. Hara,<sup>53</sup> M. Hare,<sup>54</sup> S. Harper,<sup>41</sup> R. F. Harr,<sup>56</sup> R. M. Harris,<sup>16</sup> K. Hatakeyama,<sup>48</sup> J. Hauser,<sup>8</sup> C. Hays,<sup>15</sup> H. Hayward,<sup>29</sup> A. Heijboer,<sup>43</sup> B. Heinemann,<sup>29</sup> J. Heinrich,<sup>43</sup> M. Hennecke,<sup>25</sup> M. Herndon,<sup>57</sup> J. Heuser,<sup>25</sup> D. Hidas,<sup>15</sup> C. S. Hill,<sup>10</sup> D. Hirschbuehl,<sup>25</sup> A. Hocker,<sup>16</sup> A. Holloway,<sup>21</sup> S. Hou,<sup>1</sup> M. Houlden,<sup>29</sup> S.-C. Hsu,<sup>9</sup> B. T. Huffman,<sup>41</sup> R. E. Hughes,<sup>38</sup> J. Huston,<sup>34</sup> K. Ikado,<sup>55</sup> J. Incandela,<sup>10</sup> G. Introzzi,<sup>44</sup> M. Iori,<sup>49</sup> Y. Ishizawa,<sup>53</sup> A. Ivanov,<sup>7</sup> B. Iyutin,<sup>31</sup> E. James,<sup>16</sup> D. Jang,<sup>50</sup> B. Jayatilaka,<sup>33</sup> D. Jeans,<sup>49</sup> H. Jensen,<sup>16</sup> E. J. Jeon,<sup>27</sup> M. Jones,<sup>46</sup> K. K. Joo,<sup>27</sup> S. Y. Jun,<sup>12</sup> T. R. Junk,<sup>23</sup> T. Kamon,<sup>51</sup> J. Kang,<sup>33</sup> M. Karagöz-Unel,<sup>37</sup> P. E. Karchin,<sup>56</sup> Y. Kato,<sup>40</sup> Y. Kemp,<sup>25</sup> R. Kephart,<sup>16</sup> U. Kerzel,<sup>25</sup> V. Khotilovich,<sup>51</sup> B. Kilminster,<sup>38</sup> D. H. Kim,<sup>27</sup> H. S. Kim,<sup>27</sup> J. E. Kim,<sup>27</sup> M. J. Kim,<sup>12</sup> M. S. Kim,<sup>27</sup> S. B. Kim,<sup>27</sup> S. H. Kim,<sup>53</sup> Y. K. Kim,<sup>13</sup> M. Kirby,<sup>15</sup> L. Kirsch,<sup>6</sup> S. Klimenko,<sup>17</sup> M. Klute,<sup>31</sup> B. Knuteson,<sup>31</sup> B. R. Ko,<sup>15</sup> H. Kobayashi,<sup>53</sup> K. Kondo,<sup>55</sup> D. J. Kong,<sup>27</sup> J. Konigsberg,<sup>17</sup> K. Kordas,<sup>18</sup> A. Korytov,<sup>17</sup> A. V. Kotwal,<sup>15</sup> A. Kovalev,<sup>43</sup> J. Kraus,<sup>23</sup> I. Kravchenko,<sup>31</sup> M. Kreps,<sup>25</sup> A. Kreymer,<sup>16</sup> J. Kroll,<sup>43</sup> N. Krumnack,<sup>4</sup> M. Kruse,<sup>15</sup> V. Krutelyov,<sup>51</sup> S. E. Kuhlmann,<sup>2</sup> Y. Kusakabe,<sup>55</sup> S. Kwang,<sup>13</sup> A. T. Laasanen,<sup>46</sup> S. Lai,<sup>32</sup> S. Lami,<sup>44</sup> S. Lammel,<sup>16</sup> M. Lancaster,<sup>30</sup> R. L. Lander,<sup>7</sup> K. Lannon,<sup>38</sup> A. Lath,<sup>50</sup> G. Latino,<sup>44</sup> I. Lazzizzera,<sup>42</sup> C. Lecci,<sup>25</sup> T. LeCompte,<sup>2</sup> J. Lee,<sup>47</sup> J. Lee,<sup>27</sup> S. W. Lee,<sup>51</sup> R. Lefèvre,<sup>3</sup> N. Leonardo,<sup>31</sup> S. Leone,<sup>44</sup> S. Levy,<sup>13</sup> J. D. Lewis,<sup>16</sup> K. Li,<sup>58</sup> C. Lin,<sup>58</sup> C. S. Lin,<sup>16</sup> M. Lindgren,<sup>16</sup> E. Lipeles,<sup>9</sup> T. M. Liss,<sup>23</sup> A. Lister,<sup>19</sup> D. O. Litvintsev,<sup>16</sup> T. Liu,<sup>16</sup> Y. Liu,<sup>19</sup> N. S. Lockyer,<sup>43</sup> A. Loginov,<sup>35</sup> M. Loretì,<sup>42</sup> P. Loverre,<sup>49</sup> R.-S. Lu,<sup>1</sup> D. Lucchesi,<sup>42</sup> P. Lujan,<sup>28</sup> P. Lukens,<sup>16</sup> G. Lungu,<sup>17</sup> L. Lyons,<sup>41</sup> J. Lys,<sup>28</sup> R. Lysak,<sup>1</sup> E. Lytken,<sup>46</sup> P. Mack,<sup>25</sup> D. MacQueen,<sup>32</sup> R. Madrak,<sup>16</sup> K. Maeshima,<sup>16</sup> P. Maksimovic,<sup>24</sup> G. Manca,<sup>29</sup> F. Margaroli,<sup>5</sup> R. Marginean,<sup>16</sup> C. Marino,<sup>23</sup> A. Martin,<sup>58</sup> M. Martin,<sup>24</sup> V. Martin,<sup>37</sup> M. Martínez,<sup>3</sup> T. Maruyama,<sup>53</sup> H. Matsunaga,<sup>53</sup> M. E. Mattson,<sup>56</sup> R. Mazini,<sup>32</sup> P. Mazzanti,<sup>5</sup> K. S. McFarland,<sup>47</sup> D. McGivern,<sup>30</sup> P. McIntyre,<sup>51</sup> P. McNamara,<sup>50</sup> R. McNulty,<sup>29</sup> A. Mehta,<sup>29</sup> S. Menzemer,<sup>31</sup> A. Menzione,<sup>44</sup> P. Merkel,<sup>46</sup> C. Mesropian,<sup>48</sup> A. Messina,<sup>49</sup> M. von der Mey,<sup>8</sup> T. Miao,<sup>16</sup> N. Miladinovic,<sup>6</sup> J. Miles,<sup>31</sup> R. Miller,<sup>34</sup> J. S. Miller,<sup>33</sup> C. Mills,<sup>10</sup> M. Milnik,<sup>25</sup> R. Miquel,<sup>28</sup> S. Miscetti,<sup>18</sup> G. Mitselmakher,<sup>17</sup> A. Miyamoto,<sup>26</sup> N. Moggi,<sup>5</sup> B. Mohr,<sup>8</sup> R. Moore,<sup>16</sup> M. Morello,<sup>44</sup> P. Movilla Fernandez,<sup>28</sup> J. Mülmenstädt,<sup>28</sup>

A. Mukherjee,<sup>16</sup> M. Mulhearn,<sup>31</sup> Th. Muller,<sup>25</sup> R. Mumford,<sup>24</sup> P. Murat,<sup>16</sup> J. Nachtman,<sup>16</sup> S. Nahn,<sup>58</sup> I. Nakano,<sup>39</sup>  
 A. Napier,<sup>54</sup> D. Naumov,<sup>36</sup> V. Necula,<sup>17</sup> C. Neu,<sup>43</sup> M. S. Neubauer,<sup>9</sup> J. Nielsen,<sup>28</sup> T. Nigmanov,<sup>45</sup> L. Nodulman,<sup>2</sup>  
 O. Norniella,<sup>3</sup> T. Ogawa,<sup>55</sup> S. H. Oh,<sup>15</sup> Y. D. Oh,<sup>27</sup> T. Okusawa,<sup>40</sup> R. Oldeman,<sup>29</sup> R. Orava,<sup>22</sup> K. Osterberg,<sup>22</sup>  
 C. Pagliarone,<sup>44</sup> E. Palencia,<sup>11</sup> R. Paoletti,<sup>44</sup> V. Papadimitriou,<sup>16</sup> A. Papikonomou,<sup>25</sup> A. A. Paramonov,<sup>13</sup> B. Parks,<sup>38</sup>  
 S. Pashapour,<sup>32</sup> J. Patrick,<sup>16</sup> G. Pauletta,<sup>52</sup> M. Paulini,<sup>12</sup> C. Paus,<sup>31</sup> D. E. Pellett,<sup>7</sup> A. Penzo,<sup>52</sup> T. J. Phillips,<sup>15</sup>  
 G. Piacentino,<sup>44</sup> J. Piedra,<sup>11</sup> K. Pitts,<sup>23</sup> C. Plager,<sup>8</sup> L. Pondrom,<sup>57</sup> G. Pope,<sup>45</sup> X. Portell,<sup>3</sup> O. Poukhov,<sup>14</sup> N. Pounder,<sup>41</sup>  
 F. Prakooshyn,<sup>14</sup> A. Pronko,<sup>16</sup> J. Proudfoot,<sup>2</sup> F. Ptohos,<sup>18</sup> G. Punzi,<sup>44</sup> J. Pursley,<sup>24</sup> J. Rademacker,<sup>41</sup> A. Rahaman,<sup>45</sup>  
 A. Rakitin,<sup>31</sup> S. Rappoccio,<sup>21</sup> F. Ratnikov,<sup>50</sup> B. Reiser,<sup>16</sup> V. Rekovic,<sup>36</sup> N. van Remortel,<sup>22</sup> P. Renton,<sup>41</sup> M. Rescigno,<sup>49</sup>  
 S. Richter,<sup>25</sup> F. Rimondi,<sup>5</sup> K. Rinnert,<sup>25</sup> L. Ristori,<sup>44</sup> W. J. Robertson,<sup>15</sup> A. Robson,<sup>20</sup> T. Rodrigo,<sup>11</sup> E. Rogers,<sup>23</sup> S. Rolli,<sup>54</sup>  
 R. Roser,<sup>16</sup> M. Rossi,<sup>52</sup> R. Rossin,<sup>17</sup> C. Rott,<sup>46</sup> A. Ruiz,<sup>11</sup> J. Russ,<sup>12</sup> V. Rusu,<sup>13</sup> D. Ryan,<sup>54</sup> H. Saarikko,<sup>22</sup> S. Sabik,<sup>32</sup>  
 A. Safonov,<sup>7</sup> W. K. Sakumoto,<sup>47</sup> G. Salamanna,<sup>49</sup> O. Salto,<sup>3</sup> D. Saltzberg,<sup>8</sup> C. Sanchez,<sup>3</sup> L. Santi,<sup>52</sup> S. Sarkar,<sup>49</sup> K. Sato,<sup>53</sup>  
 P. Savard,<sup>32</sup> A. Savoy-Navarro,<sup>16</sup> T. Scheidle,<sup>25</sup> P. Schlabach,<sup>16</sup> E. E. Schmidt,<sup>16</sup> M. P. Schmidt,<sup>58</sup> M. Schmitt,<sup>37</sup>  
 T. Schwarz,<sup>33</sup> L. Scodellaro,<sup>11</sup> A. L. Scott,<sup>10</sup> A. Scribano,<sup>44</sup> F. Scuri,<sup>44</sup> A. Sedov,<sup>46</sup> S. Seidel,<sup>36</sup> Y. Seiya,<sup>40</sup> A. Semenov,<sup>14</sup>  
 F. Semeria,<sup>5</sup> L. Sexton-Kennedy,<sup>16</sup> I. Sfiligoi,<sup>18</sup> M. D. Shapiro,<sup>28</sup> T. Shears,<sup>29</sup> P. F. Shepard,<sup>45</sup> D. Sherman,<sup>21</sup>  
 M. Shimojima,<sup>53</sup> M. Shochet,<sup>13</sup> Y. Shon,<sup>57</sup> I. Shreyber,<sup>35</sup> A. Sidoti,<sup>44</sup> A. Sill,<sup>16</sup> P. Sinervo,<sup>32</sup> A. Sisakyan,<sup>14</sup> J. Sjolín,<sup>41</sup>  
 A. Skiba,<sup>25</sup> A. J. Slaughter,<sup>16</sup> K. Sliwa,<sup>54</sup> D. Smirnov,<sup>36</sup> J. R. Smith,<sup>7</sup> F. D. Snider,<sup>16</sup> R. Snihur,<sup>32</sup> M. Soderberg,<sup>33</sup> A. Soha,<sup>7</sup>  
 S. Somalwar,<sup>50</sup> V. Sorin,<sup>34</sup> J. Spalding,<sup>16</sup> F. Spinella,<sup>44</sup> P. Squillacioti,<sup>44</sup> M. Stanitzki,<sup>58</sup> A. Staveris-Polykalas,<sup>44</sup>  
 R. St. Denis,<sup>20</sup> B. Stelzer,<sup>8</sup> O. Stelzer-Chilton,<sup>32</sup> D. Stentz,<sup>37</sup> J. Strologas,<sup>36</sup> D. Stuart,<sup>10</sup> J. S. Suh,<sup>27</sup> A. Sukhanov,<sup>17</sup>  
 K. Sumorok,<sup>31</sup> H. Sun,<sup>54</sup> T. Suzuki,<sup>53</sup> A. Taffard,<sup>23</sup> R. Tafirout,<sup>32</sup> R. Takashima,<sup>39</sup> Y. Takeuchi,<sup>53</sup> K. Takikawa,<sup>53</sup>  
 M. Tanaka,<sup>2</sup> R. Tanaka,<sup>39</sup> M. Tecchio,<sup>33</sup> P. K. Teng,<sup>1</sup> K. Terashi,<sup>48</sup> S. Tether,<sup>31</sup> J. Thom,<sup>16</sup> A. S. Thompson,<sup>20</sup>  
 E. Thomson,<sup>43</sup> P. Tipton,<sup>47</sup> V. Tiwari,<sup>12</sup> S. Tkaczyk,<sup>16</sup> D. Toback,<sup>51</sup> K. Tollefson,<sup>34</sup> T. Tomura,<sup>53</sup> D. Tonelli,<sup>44</sup>  
 M. Tönnemann,<sup>34</sup> S. Torre,<sup>44</sup> D. Torretta,<sup>16</sup> S. Tourneur,<sup>16</sup> W. Trischuk,<sup>32</sup> R. Tsuchiya,<sup>55</sup> S. Tsuno,<sup>39</sup> N. Turini,<sup>44</sup>  
 F. Ukegawa,<sup>53</sup> T. Unverhau,<sup>20</sup> S. Uozumi,<sup>53</sup> D. Usynin,<sup>43</sup> L. Vacavant,<sup>28</sup> A. Vaiciulis,<sup>47</sup> S. Vallecorsa,<sup>19</sup> A. Varganov,<sup>33</sup>  
 E. Vataga,<sup>36</sup> G. Velev,<sup>16</sup> G. Veramendi,<sup>23</sup> V. Veszpremi,<sup>46</sup> T. Vickey,<sup>23</sup> R. Vidal,<sup>16</sup> I. Vila,<sup>11</sup> R. Vilar,<sup>11</sup> I. Vollrath,<sup>32</sup>  
 I. Volobouev,<sup>28</sup> F. Würthwein,<sup>9</sup> P. Wagner,<sup>51</sup> R. G. Wagner,<sup>2</sup> R. L. Wagner,<sup>16</sup> W. Wagner,<sup>25</sup> R. Wallny,<sup>8</sup> T. Walter,<sup>25</sup>  
 Z. Wan,<sup>50</sup> M. J. Wang,<sup>1</sup> S. M. Wang,<sup>17</sup> A. Warburton,<sup>32</sup> B. Ward,<sup>20</sup> S. Waschke,<sup>20</sup> D. Waters,<sup>30</sup> T. Watts,<sup>50</sup> M. Weber,<sup>28</sup>  
 W. C. Wester III,<sup>16</sup> B. Whitehouse,<sup>54</sup> D. Whiteson,<sup>43</sup> A. B. Wicklund,<sup>2</sup> E. Wicklund,<sup>16</sup> H. H. Williams,<sup>43</sup> P. Wilson,<sup>16</sup>  
 B. L. Winer,<sup>38</sup> P. Wittich,<sup>43</sup> S. Wolbers,<sup>16</sup> C. Wolfe,<sup>13</sup> S. Worm,<sup>50</sup> T. Wright,<sup>33</sup> X. Wu,<sup>19</sup> S. M. Wynne,<sup>29</sup> A. Yagil,<sup>16</sup>  
 K. Yamamoto,<sup>40</sup> J. Yamaoka,<sup>50</sup> Y. Yamashita,<sup>39</sup> C. Yang,<sup>58</sup> U. K. Yang,<sup>13</sup> W. M. Yao,<sup>28</sup> G. P. Yeh,<sup>16</sup> J. Yoh,<sup>16</sup> K. Yorita,<sup>13</sup>  
 T. Yoshida,<sup>40</sup> I. Yu,<sup>27</sup> S. S. Yu,<sup>43</sup> J. C. Yun,<sup>16</sup> L. Zanello,<sup>49</sup> A. Zanetti,<sup>52</sup> I. Zaw,<sup>21</sup> F. Zetti,<sup>44</sup> X. Zhang,<sup>23</sup>  
 J. Zhou,<sup>50</sup> and S. Zucchelli<sup>5</sup>

(CDF Collaboration)

<sup>1</sup>*Institute of Physics, Academia Sinica, Taipei, Taiwan 11529, Republic of China*<sup>2</sup>*Argonne National Laboratory, Argonne, Illinois 60439, USA*<sup>3</sup>*Institut de Física d'Altes Energies, Universitat Autònoma de Barcelona, E-08193, Bellaterra (Barcelona), Spain*<sup>4</sup>*Baylor University, Waco, Texas 76798, USA*<sup>5</sup>*Istituto Nazionale di Fisica Nucleare, University of Bologna, I-40127 Bologna, Italy*<sup>6</sup>*Brandeis University, Waltham, Massachusetts 02254, USA*<sup>7</sup>*University of California–Davis, Davis, California 95616, USA*<sup>8</sup>*University of California–Los Angeles, Los Angeles, California 90024, USA*<sup>9</sup>*University of California–San Diego, La Jolla, California 92093, USA*<sup>10</sup>*University of California–Santa Barbara, Santa Barbara, California 93106, USA*<sup>11</sup>*Instituto de Física de Cantabria, CSIC-University of Cantabria, 39005 Santander, Spain*<sup>12</sup>*Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA*<sup>13</sup>*Enrico Fermi Institute, University of Chicago, Chicago, Illinois 60637, USA*<sup>14</sup>*Joint Institute for Nuclear Research, RU-141980 Dubna, Russia*<sup>15</sup>*Duke University, Durham, North Carolina 27708, USA*<sup>16</sup>*Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA*<sup>17</sup>*University of Florida, Gainesville, Florida 32611, USA*<sup>18</sup>*Laboratori Nazionali di Frascati, Istituto Nazionale di Fisica Nucleare, I-00044 Frascati, Italy*<sup>19</sup>*University of Geneva, CH-1211 Geneva 4, Switzerland*<sup>20</sup>*Glasgow University, Glasgow G12 8QQ, United Kingdom*

- <sup>21</sup>Harvard University, Cambridge, Massachusetts 02138, USA
- <sup>22</sup>Division of High Energy Physics, Department of Physics, University of Helsinki and Helsinki Institute of Physics, FIN-00014, Helsinki, Finland
- <sup>23</sup>University of Illinois, Urbana, Illinois 61801, USA
- <sup>24</sup>The Johns Hopkins University, Baltimore, Maryland 21218, USA
- <sup>25</sup>Institut für Experimentelle Kernphysik, Universität Karlsruhe, 76128 Karlsruhe, Germany
- <sup>26</sup>High Energy Accelerator Research Organization (KEK), Tsukuba, Ibaraki 305, Japan
- <sup>27</sup>Center for High Energy Physics: Kyungpook National University, Taegu 702-701; Seoul National University, Seoul 151-742; and SungKyunKwan University, Suwon 440-746; Korea
- <sup>28</sup>Ernest Orlando Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA
- <sup>29</sup>University of Liverpool, Liverpool L69 7ZE, United Kingdom
- <sup>30</sup>University College London, London WC1E 6BT, United Kingdom
- <sup>31</sup>Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA
- <sup>32</sup>Institute of Particle Physics: McGill University, Montréal, Canada H3A 2T8; and University of Toronto, Toronto, Canada M5S 1A7
- <sup>33</sup>University of Michigan, Ann Arbor, Michigan 48109, USA
- <sup>34</sup>Michigan State University, East Lansing, Michigan 48824, USA
- <sup>35</sup>Institution for Theoretical and Experimental Physics, ITEP, Moscow 117259, Russia
- <sup>36</sup>University of New Mexico, Albuquerque, New Mexico 87131, USA
- <sup>37</sup>Northwestern University, Evanston, Illinois 60208, USA
- <sup>38</sup>The Ohio State University, Columbus, Ohio 43210, USA
- <sup>39</sup>Okayama University, Okayama 700-8530, Japan
- <sup>40</sup>Osaka City University, Osaka 588, Japan
- <sup>41</sup>University of Oxford, Oxford OX1 3RH, United Kingdom
- <sup>42</sup>University of Padova, Istituto Nazionale di Fisica Nucleare, Sezione di Padova-Trento, I-35131 Padova, Italy
- <sup>43</sup>University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA
- <sup>44</sup>Istituto Nazionale di Fisica Nucleare Pisa, Universities of Pisa, Siena and Scuola Normale Superiore, I-56127 Pisa, Italy
- <sup>45</sup>University of Pittsburgh, Pittsburgh, Pennsylvania 15260, USA
- <sup>46</sup>Purdue University, West Lafayette, Indiana 47907, USA
- <sup>47</sup>University of Rochester, Rochester, New York 14627, USA
- <sup>48</sup>The Rockefeller University, New York, New York 10021, USA
- <sup>49</sup>Istituto Nazionale di Fisica Nucleare, Sezione di Roma 1, University of Rome “La Sapienza”, I-00185 Roma, Italy
- <sup>50</sup>Rutgers University, Piscataway, New Jersey 08855, USA
- <sup>51</sup>Texas A&M University, College Station, Texas 77843, USA
- <sup>52</sup>Istituto Nazionale di Fisica Nucleare, University of Trieste/Udine, Italy
- <sup>53</sup>University of Tsukuba, Tsukuba, Ibaraki 305, Japan
- <sup>54</sup>Tufts University, Medford, Massachusetts 02155, USA
- <sup>55</sup>Waseda University, Tokyo 169, Japan
- <sup>56</sup>Wayne State University, Detroit, Michigan 48201, USA
- <sup>57</sup>University of Wisconsin, Madison, Wisconsin 53706, USA
- <sup>58</sup>Yale University, New Haven, Connecticut 06520, USA

(Received 24 August 2005; published 5 January 2006)

We present a search for neutral supersymmetric Higgs bosons decaying to  $\tau$  pairs produced in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV. The data, corresponding to  $310 \text{ pb}^{-1}$  integrated luminosity, were collected with the Collider Detector at Fermilab in run II of the Tevatron. No significant excess above the standard model backgrounds is observed. We set exclusion limits on the production cross section times branching fraction to  $\tau$  pairs for Higgs boson masses in the range from 90 to 250 GeV/ $c^2$ .

DOI: [10.1103/PhysRevLett.96.011802](https://doi.org/10.1103/PhysRevLett.96.011802)

PACS numbers: 14.80.Cp, 12.60.Fr, 12.60.Jv, 13.85.Rm

One of the outstanding questions in particle physics is the dynamics of electroweak (EW) symmetry breaking and the origin of particle masses. In the standard model (SM), EW symmetry is spontaneously broken through the Higgs mechanism [1], which predicts the existence of a massive scalar Higgs boson  $h_{\text{SM}}$ . Theoretical difficulties related to divergent radiative corrections to the  $h_{\text{SM}}$  mass have natural solutions in supersymmetric (SUSY) models [2].

The minimal supersymmetric extension of the standard model (MSSM) [3] is the simplest realistic SUSY theory. The Higgs sector in the MSSM consists of two charged and three neutral scalar bosons. Assuming  $CP$  invariance, one of the neutral bosons ( $A$ ) is  $CP$ -odd, and the other two ( $h, H$ ) are  $CP$ -even. Throughout this Letter, we use  $h$  ( $H$ ) for the lighter (heavier)  $CP$ -even neutral Higgs boson and  $\phi$  to denote any of  $h, H, A$ . At tree level, the MSSM Higgs

bosons are described by the mass of  $A$  ( $m_A$ ), and  $\tan\beta = v_u/v_d$ , where  $v_u, v_d$  are the vacuum expectation values of the neutral Higgs fields that couple to up-type and down-type fermions, respectively. The Yukawa couplings of  $A$  to down-type fermions (such as the  $b$  quark and  $\tau$ ) are enhanced by a factor of  $\tan\beta$  relative to the SM. For large  $\tan\beta$ , one of the  $CP$ -even bosons is nearly mass-degenerate with  $A$  and has similar couplings. The dominant production mechanisms of neutral MSSM Higgs bosons at hadron colliders are gluon fusion [4] and  $b\bar{b}$  fusion [5,6]. The leading decay modes of  $A$  and the corresponding mass-degenerate  $CP$ -even Higgs boson are  $\phi \rightarrow b\bar{b}$  ( $\sim 90\%$ ) and  $\phi \rightarrow \tau\tau$  ( $\sim 10\%$ ).

In this Letter, we present the results of a search for neutral MSSM Higgs bosons produced in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV. The data sample of  $310 \text{ pb}^{-1}$  integrated luminosity was collected with the upgraded Collider Detector at Fermilab (CDF II) between 2002 and 2004 in run II of the Tevatron. The search is performed in the  $\phi \rightarrow \tau\tau$  decay channel for  $90 < m_A < 250 \text{ GeV}/c^2$ . One  $\tau$  is detected in the decay to an  $e$  or  $\mu$  and neutrinos, and the other in the decay to hadrons and a neutrino. In the following, we use  $\tau_e, \tau_\mu$ , and  $\tau_{\text{had}}$  as shorthand notations for the decay modes  $\tau \rightarrow e\nu_e\nu_\tau$ ,  $\tau \rightarrow \mu\nu_\mu\nu_\tau$ , and  $\tau \rightarrow \text{hadrons}\nu_\tau$ , respectively. Previous and related searches in the di- $\tau$  channel are presented in Refs. [7,8].

CDF II [9] is a general purpose detector with tracking and calorimetry. The tracking system consists of silicon microstrip detectors and a cylindrical wire drift chamber. It is immersed in a 1.4 T magnetic field produced by a superconducting solenoid. Electromagnetic (EM) and hadronic (HAD) sampling calorimeters are located outside the solenoid and cover detector pseudorapidity  $|\eta| < 3.6$ , where  $\eta = -\ln(\tan(\theta/2))$  and  $\theta$  is the polar angle with respect to the proton beam. The calorimeters are divided into towers with projective geometry. A central electromagnetic shower maximum detector (CES) consisting of proportional chambers with anode wires parallel to the beam axis and orthogonal cathode strips is embedded in the EM calorimeter at a depth of six radiation lengths. The CES is used to determine the position of EM showers with spatial resolution of  $\sim 0.5$  cm. Muons are identified by a system of drift chambers located outside the calorimeter volume with combined coverage extending to  $|\eta| < 1.5$ . The luminosity is measured by gas Cherenkov counters located in the detector forward and backward regions ( $3.7 < |\eta| < 4.7$ ) with 6% precision [10].

The search for  $\phi \rightarrow \tau\tau$  requires detection of an  $e$  or  $\mu$  (from  $\tau_e, \tau_\mu$ ) and the reconstruction of the  $\tau_{\text{had}}$  decay products. Events are preselected with “lepton plus track” triggers [11]. The triggers require a lepton ( $e, \mu$ ) candidate and another track, both pointing to the central calorimeter ( $|\eta| \lesssim 1.0$ ) and having azimuthal separation  $\Delta\phi > 10^\circ$ . The overall trigger efficiency for signal

events passing the selection criteria described below is greater than 90%. The algorithms for  $e$  and  $\mu$  identification are described in detail in Ref. [9]. The vector sum of the transverse momenta [12] of the neutrinos from  $\tau$  decays appears as missing transverse energy ( $\cancel{E}_T$ ), determined from the imbalance of energy deposition in the calorimeter [13]. The decay products in  $\tau_{\text{had}}$  form narrow jets with low multiplicity of neutral and charged particles. The positions and energies of  $\pi^0$ 's and photons are reconstructed with the CES detector and EM calorimeter, respectively. In this search, we do not distinguish reconstructed photons and  $\pi^0$ 's, and all neutrals are assumed to be  $\pi^0$ 's. A charged track with  $p_T > 6 \text{ GeV}/c$  pointing to a cluster of six or fewer contiguous calorimeter towers serves as a seed for a  $\tau_{\text{had}}$  candidate. The direction of the track defines the axis of a signal cone of size  $\alpha_{\text{sig}}$  and an isolation annulus extending from  $\alpha_{\text{sig}}$  to  $\alpha_{\text{iso}} = 0.52$  rad. The signal cone size depends on the calorimeter cluster energy  $E^{\text{cl}}$ :  $\alpha_{\text{sig}}$  is the minimum of 0.17 and  $(5 \text{ GeV})/E^{\text{cl}}$  rad. To reduce position resolution effects, the minimum value of  $\alpha_{\text{sig}}$  is set to 0.05 (0.1) rad for tracks ( $\pi^0$ 's). The four-momentum of  $\tau_{\text{had}}$  is calculated from tracks and  $\pi^0$ 's in the signal cone. Particles in the isolation annulus are used to impose requirements that discriminate against quark and gluon jets: The scalar sum of the  $p_T$  of tracks (sum of  $E_T$  of  $\pi^0$ 's) is required to be less than  $1 \text{ GeV}/c$  (1 GeV). We select  $\tau_{\text{had}}$  candidates with one or three tracks in the signal cone ( $N_{\text{sig}}^{\text{trk}} = 1, 3$ ) with  $p_T > 1 \text{ GeV}/c$ , consistent with the dominant  $\tau$  decay modes. In the  $N_{\text{sig}}^{\text{trk}} = 3$  case, the sum of the electric charges must be equal to  $\pm 1$ . The invariant mass of the hadronic system is required to be less than  $1.8 \text{ GeV}/c^2$ . Electrons are rejected by imposing the condition  $(E^{\text{cl}}/P_{\text{sig}}^{\text{trk}})(0.95 - f) > 0.1$ , where  $f$  is the ratio of EM to HAD energy in the calorimeter cluster, and  $P_{\text{sig}}^{\text{trk}}$  is the scalar sum of track momenta in the signal cone. Muons are suppressed by requiring  $E_T^{\text{cl}} > 15 \text{ GeV}$ . The  $\tau_{\text{had}}$  identification efficiency increases from 38% at transverse momentum of the hadronic system  $p_T^{\text{had}} = 15 \text{ GeV}/c$  to  $\sim 46\%$  for  $p_T^{\text{had}} \geq 25 \text{ GeV}/c$ . The probability for misidentifying a quark or gluon jet as  $\tau_{\text{had}}$  is measured using jet data samples. It is  $\sim 1.5\%$  for jet transverse energy  $E_T^{\text{jet}} = 20 \text{ GeV}$ , dropping to  $\sim 0.1\%$  for  $E_T^{\text{jet}} = 100 \text{ GeV}$ .

The acceptances for signal and most of the backgrounds are determined from samples of Monte Carlo (MC) simulated events produced by the PYTHIA event generator [14] with CTEQ5L [15] parton distribution functions (PDF's). Tau decays are simulated by the TAUOLA package [16]. Detector response is simulated with a GEANT-based [17] model of the detector.

The dominant (and irreducible) background in the final sample of selected events is from inclusive  $Z/\gamma^*$  production with subsequent decays to  $\tau$  pairs. It is estimated using MC simulated events with normalization corresponding to

$\sigma(p\bar{p} \rightarrow Z/\gamma^* \rightarrow l\bar{l}) = 254.9$  pb in the dilepton mass region  $66 < m_{ll} < 116$  GeV/ $c^2$  [18]. The second largest background contribution comes from processes with quark or gluon jets misidentified as  $\tau_{\text{had}}$ , such as dijet and multijet,  $W$  + jets, and  $\gamma$  + jets production. These backgrounds are estimated from the data by applying jet  $\rightarrow \tau_{\text{had}}$  misidentification rates to jets in events that pass all selection criteria except for  $\tau_{\text{had}}$  identification. The validity of the predictions is verified using independent data samples representing the background processes. The third group of backgrounds includes  $Z/\gamma^* \rightarrow l\bar{l}$  ( $l = e, \mu$ ),  $WW$ ,  $WZ$ ,  $ZZ$ , and  $t\bar{t}$  production. Their contributions are determined from MC samples normalized to the theoretical cross sections.

The events in the  $\tau_e\tau_{\text{had}}$  ( $\tau_\mu\tau_{\text{had}}$ ) channel are selected by requiring one  $e$  ( $\mu$ ) candidate with  $p_T^{e(\mu)} > 10$  GeV/ $c$  and one  $\tau_{\text{had}}$  candidate with  $p_T^{\text{had}} > 15$  GeV/ $c$  and opposite electric charge. Low-energy multijet backgrounds are suppressed by rejecting events with  $|p_T^{e(\mu)}| + |p_T^{\text{had}}| + |\cancel{E}_T| < 50$  GeV. Backgrounds from  $W$  + jet events are suppressed by imposing a requirement on the relative directions of the visible  $\tau$  decay products and  $\cancel{E}_T$ . We define a unit vector  $\hat{\zeta}$  along the bisector of the angle between the directions of  $e$  ( $\mu$ ) and  $\tau_{\text{had}}$  in the transverse plane. The projections  $p_\zeta^{\text{vis}} = (\vec{p}_{e(\mu)} + \vec{p}_{\text{had}}) \cdot \hat{\zeta}$  and  $p_\zeta^{\cancel{E}_T} = \vec{\cancel{E}}_T \cdot \hat{\zeta}$  are required to satisfy  $p_\zeta^{\cancel{E}_T} > 0.6p_\zeta^{\text{vis}} - 10$  GeV/ $c$ . This condition removes  $\sim 85\%$  of the  $W$  + jet events passing the other selection criteria while retaining  $\sim 95\%$  of the signal. To suppress backgrounds from  $Z \rightarrow l\bar{l}$  decays with a misidentified lepton, we do not accept events with invariant mass of an  $e$  ( $\mu$ ) and a single-track  $\tau_{\text{had}}$  candidate within 10 GeV/ $c^2$  of the  $Z$  mass. The combined signal acceptance for a Higgs boson of mass 90 GeV/ $c^2$  (250 GeV/ $c^2$ ) in the  $\tau_e\tau_{\text{had}}$  and  $\tau_\mu\tau_{\text{had}}$  channels is 0.8% (2.0%).

The systematic uncertainties for particle identification efficiency are 3.5% ( $\tau_{\text{had}}$ ), 1.3% ( $e$ ), and 4.6% ( $\mu$ ). The uncertainties in trigger efficiency for the  $\tau_e\tau_{\text{had}}$  and  $\tau_\mu\tau_{\text{had}}$  channels are 2.1% and 1.4%, respectively. The uncertainty in the determination of backgrounds due to jet  $\rightarrow \tau$  misidentification is 20%, resulting in 3% effect on the total background estimate. The systematic uncertainty in signal acceptance from event-level cuts is less than 2%. The imprecise knowledge of the PDF's introduces an additional 5.7% uncertainty on signal acceptance [19].

Figure 1 shows the track multiplicity distribution for  $\tau_{\text{had}}$  candidates in the data, along with the background predictions. The characteristic enhancement in the one- and three-track bins clearly shows the contribution from events with  $\tau_{\text{had}}$  in the final state. The total number of expected events from SM processes after applying all selection criteria is  $N_{\text{SM}} = 496 \pm 5(\text{stat}) \pm 28(\text{syst}) \pm 25(\text{lumi})$ . The contributions from  $Z/\gamma^* \rightarrow \tau\tau$ , backgrounds with jet  $\rightarrow \tau$  misidentification, and all remaining background sources are 405, 75, and 16, respectively.




FIG. 1 (color online). Track multiplicity for hadronically decaying tau candidates before applying the opposite charge and  $N_{\text{trk}}^{\text{sig}}$  requirements.

We observe 487 events, in agreement with  $N_{\text{SM}}$ . To probe for a possible Higgs signal, we perform binned likelihood fits of the partially reconstructed mass of the di- $\tau$  system ( $m_{\text{vis}}$ ) defined as the invariant mass of the visible tau decay products and  $\cancel{E}_T$ . The backgrounds are allowed to float within limits set by Gaussian constraints corresponding to the systematic uncertainties in trigger efficiencies, particle identification, production cross sections, PDF's, event cuts, and luminosity measurement. Potential differences in  $m_{\text{vis}}$  shapes between data and the MC simulation in different channels are treated as systematic uncertainties. We create signal and background  $m_{\text{vis}}$  templates with the MC energy scales shifted from the nominal values according to the uncertainties and study the effect on hypothetical cross section measurements. The deviations from the results obtained with the nominal templates are parametrized in terms of the Higgs boson mass and input cross section. An example fit for  $m_A = 140$  GeV/ $c^2$  is shown in Fig. 2. We observe no signal evidence for  $m_A = 90$ –250 GeV/ $c^2$  and set exclusion limits at 95% C.L. on  $\sigma(p\bar{p} \rightarrow \phi + X) \times \text{BR}(\phi \rightarrow \tau\tau)$  as




FIG. 2 (color online). Example fit of the  $m_{\text{vis}}$  distribution for signal with  $m_A = 140$  GeV/ $c^2$ . Signal and background normalizations correspond to the fit results for signal exclusion at 95% C.L.




FIG. 3 (color online). Upper limits at 95% C.L. on Higgs production cross section times branching fraction to  $\tau$  pairs. The expected limits from the pseudoexperiments are also shown.

shown in Fig. 3. The sensitivity of the limit-setting procedure is determined from MC simulations assuming no signal. The  $m_{\text{vis}}$  shape uncertainty leads to 15% (5%) deterioration of the limits for the low (high) end of the considered  $m_A$  region. The observed limits range from 24.4 pb for  $m_A = 90 \text{ GeV}/c^2$ , to 9.3 pb for  $m_A = 140 \text{ GeV}/c^2$ , to 1.8 pb for  $m_A = 250 \text{ GeV}/c^2$ .

Using the theoretical predictions for the MSSM Higgs boson production and decay to  $\tau$  pairs, we interpret the limits on  $\sigma(p\bar{p} \rightarrow \phi + X) \times \text{BR}(\phi \rightarrow \tau\tau)$  as exclusions of parameter regions in the  $\tan\beta$  vs  $m_A$  plane. The cross sections are obtained from SM calculations and scaling factors  $\sigma_{\text{MSSM}}/\sigma_{\text{SM}}$  accounting for the modified Higgs couplings [20]. The cross sections for gluon fusion mediated by a  $b$ -quark loop are calculated with the HIGLU program [21]. The corresponding values for  $b\bar{b} \rightarrow \phi + X$




FIG. 4 (color online). Excluded regions in the  $\tan\beta$  vs  $m_A$  plane for the  $m_h^{\text{max}}$  and no-mixing scenarios with (a)  $\mu < 0$  and (b)  $\mu > 0$ .

are taken from Ref. [6]. The scaling factors and  $\text{BR}(\phi \rightarrow \tau\tau)$  are calculated with the FEYNHIGGS program [22]. They depend on  $m_A$ ,  $\tan\beta$ , the  $SU(2)$  gaugino mass parameter  $M_2$ , the SUSY mass scale  $M_{\text{SUSY}}$ , the squark mixing parameter  $X_t$ , the gluino mass  $m_{\tilde{g}}$ , and the Higgs mixing parameter  $\mu$ . We consider four benchmarks [23]: the  $m_h^{\text{max}}$  and no-mixing scenarios, with  $\mu > 0$  and  $\mu < 0$ . The excluded  $\tan\beta$  vs  $m_A$  regions are shown in Fig. 4.

The LEP experiments have excluded  $m_A \lesssim 93 \text{ GeV}/c^2$  and higher-mass  $A$  for small  $\tan\beta$  [24]. Our search is complementary, providing sensitivity in the large  $\tan\beta$  region. The excluded parameter space in the  $\tan\beta$  vs  $m_A$  plane for  $\mu < 0$  is similar to the D0 results obtained in the  $\phi \rightarrow b\bar{b}$  decay mode [25] and extends to higher  $m_A$ . Moreover, our results in the  $\phi \rightarrow \tau\tau$  channel allow us to set comparable exclusions for scenarios with  $\mu > 0$ , as the lower production cross sections are compensated by an increase in  $\text{BR}(\phi \rightarrow \tau\tau)$ .

We thank A. Belyaev, M. Carena, J. Gunion, T. Han, S. Heinemeyer, W. Kilgore, S. Mrenna, M. Spira, C. Wagner, G. Weiglein, and S. Willenbrock for illuminating discussions on the theory of MSSM Higgs production and decays. We thank the Fermilab staff and the technical staffs of the participating institutions for their vital contributions. This work was supported by the U.S. Department of Energy and National Science Foundation; the Italian Istituto Nazionale di Fisica Nucleare; the Ministry of Education, Culture, Sports, Science and Technology of Japan; the Natural Sciences and Engineering Research Council of Canada; the National Science Council of the Republic of China; the Swiss National Science Foundation; the A.P. Sloan Foundation; the Bundesministerium für Bildung und Forschung, Germany; the Korean Science and Engineering Foundation and the Korean Research Foundation; the Particle Physics and Astronomy Research Council and the Royal Society, U.K.; the Russian Foundation for Basic Research; the Comisión Interministerial de Ciencia y Tecnología, Spain; in part by the European Community's Human Potential Programme under Contract No. HPRN-CT-2002-00292; and the Academy of Finland.

- [1] P. W. Higgs, Phys. Lett. **12**, 132 (1964); Phys. Rev. Lett. **13**, 508 (1964); Phys. Rev. **145**, 1156 (1966).
- [2] S. P. Martin, hep-ph/9709356, and references therein.
- [3] D. J. H. Chung *et al.*, Phys. Rep. **407**, 1 (2005), and references therein.
- [4] S. Dawson, A. Djouadi, and M. Spira, Phys. Rev. Lett. **77**, 16 (1996).
- [5] F. Maltoni, Z. Sullivan, and S. Willenbrock, Phys. Rev. D **67**, 093005 (2003).
- [6] R. V. Harlander and W. B. Kilgore, Phys. Rev. D **68**, 013001 (2003).
- [7] D. Acosta *et al.* (CDF Collaboration), Phys. Rev. D **72**, 072004 (2005).

- [8] D. Acosta *et al.* (CDF Collaboration), Phys. Rev. Lett. **95**, 131801 (2005).
- [9] D. Acosta *et al.* (CDF Collaboration), Phys. Rev. D **71**, 032001 (2005).
- [10] S. Klimenko, J. Konigsberg, and T.M. Liss, Fermilab Report No. FERMILAB-FN-0741, 2003.
- [11] A. Anastassov *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **518**, 609 (2004).
- [12] We define transverse momentum ( $p_T$ ) and transverse energy ( $E_T$ ) as  $p_T = p \sin\theta$  and  $E_T = E \sin\theta$ .
- [13] D. Acosta *et al.* (CDF Collaboration), Phys. Rev. D **72**, 052003 (2005).
- [14] T. Sjöstrand *et al.*, Comput. Phys. Commun. **135**, 238 (2001). We use PYTHIA v. 6.215.
- [15] H.L. Lai *et al.* (CTEQ Collaboration), Eur. Phys. J. C **12**, 375 (2000).
- [16] Z. Was *et al.*, Nucl. Phys. B, Proc. Suppl. **98**, 96 (2001).
- [17] R. Brun and F. Carminati, CERN Programming Library Long Writeup W5013, 1993.
- [18] D. Acosta *et al.* (CDF Collaboration), Phys. Rev. Lett. **94**, 091803 (2005).
- [19] We compare the acceptances obtained with the CTEQ5 and CTEQ6 PDF sets. In the latter case, we also include the effect of the eigenvector variations.
- [20] M. Carena, S. Heinemeyer, G. Weiglein, and C.E.M. Wagner, Fermilab Report No. FERMILAB-PUB-05-370-T, 2005.
- [21] M. Spira, Nucl. Instrum. Methods Phys. Res., Sect. A **389**, 357 (1997).
- [22] S. Heinemeyer, W. Hollik, and G. Weiglein, Eur. Phys. J. C **9**, 343 (1999); Comput. Phys. Commun. **124**, 76 (2000).
- [23] M. Carena, S. Heinemeyer, C.E.M. Wagner, and G. Weiglein, hep-ph/9912223; Eur. Phys. J. C **26**, 601 (2003).
- [24] ALEPH, DELPHI, L3, and OPAL Collaborations, LHWG-Note 2004-01, 2004.
- [25] V. Abazov *et al.* (D0 Collaboration), Phys. Rev. Lett. **95**, 151801 (2005).