

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

USDA National Wildlife Research Center - Staff
Publications

U.S. Department of Agriculture: Animal and
Plant Health Inspection Service

November 2003

AMOUNT AND ECONOMIC VALUATION OF FERAL HOG DAMAGE TO A UNIQUE BASIN MARSH WETLAND IN FLORIDA

Richard M. Engeman

USDA-APHIS-Wildlife Services, s_r100@yahoo.com

Henry T. Smith

Florida Department of Environmental Protection, Florida Park Service

Robert G. Severson

Florida Department of Environmental Protection, Savannas Preserve State Park

Mary Ann M. Severson

Florida Department of Environmental Protection, Savannas Preserve State Park

Stephanie A. Shwiff

USDA/APHIS/WS National Wildlife Research Center, stephanie.a.shwiff@aphis.usda.gov

See next page for additional authors

Follow this and additional works at: https://digitalcommons.unl.edu/icwdm_usdanwrc



Part of the [Environmental Sciences Commons](#)

Engeman, Richard M.; Smith, Henry T.; Severson, Robert G.; Severson, Mary Ann M.; Shwiff, Stephanie A.; Constantin, Bernice; and Griffin, Daniel, "AMOUNT AND ECONOMIC VALUATION OF FERAL HOG DAMAGE TO A UNIQUE BASIN MARSH WETLAND IN FLORIDA" (2003). *USDA National Wildlife Research Center - Staff Publications*. 217.

https://digitalcommons.unl.edu/icwdm_usdanwrc/217

This Article is brought to you for free and open access by the U.S. Department of Agriculture: Animal and Plant Health Inspection Service at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in USDA National Wildlife Research Center - Staff Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors

Richard M. Engeman, Henry T. Smith, Robert G. Severson, Mary Ann M. Severson, Stephanie A. Shwiff, Bernice Constantin, and Daniel Griffin



AMOUNT AND ECONOMIC VALUATION OF FERAL HOG DAMAGE TO A UNIQUE BASIN MARSH WETLAND IN FLORIDA

RICHARD M. ENGEMAN^{1*}, HENRY T. SMITH², ROBERT G. SEVERSON³, MARY ANN M. SEVERSON³, STEPHANIE A. SHWIFF¹, BERNICE CONSTANTIN⁴, AND DANIEL GRIFFIN³

¹ National Wildlife Research Center, 4101 LaPorte Ave., Fort Collins, Colorado 80521-2154 USA.

² Florida Department of Environmental Protection, Florida Park Service,
13798 S.E. Federal Highway, Hobe Sound, Florida 33455 USA.

³ Florida Department of Environmental Protection, Savannas Preserve State Park,
9551 Gumbo Limbo Ln., Jensen Beach, Florida 34957 USA.

⁴ USDA Wildlife Services, 2820 East University Ave., Gainesville, Florida 32641 USA.

* Author to whom correspondence should be addressed: Richard M. Engeman,
Telephone (970) 266-6091, Fax (970) 266-6089, E-mail: richard.m.engeman@aphis.usda.gov

Abstract. Feral hogs (*Sus scrofa*) have been introduced into many natural habitats throughout the world, and they have adversely affected the environment in most of those places. Basin marshes are unique, but dwindling ecosystems in Florida that are especially vulnerable to damage by feral hogs. We estimated the amount of hog damage to the last remnant of a basin marsh system in Savannas Preserve State Park (SPSP), and to ecotones within the marsh. We also applied an economic valuation method for the hog damage that was based on the dollar amounts that wetland regulators have allowed permit applicants to spend in mitigation attempts to replace lost wetland resources. We found that hogs damaged 19% of the exposed portion of the basin marsh in our study area. Seventy percent of the sample sites showed hog damage at the shoreline ecotone and 58% showed damage at the upland ecotone. The area damaged within our study site alone was valued between \$1,238,760 and \$4,036,290. These damage valuation estimates were considered conservative, because it was impossible to incorporate values for such contingencies as hog impacts to state and federally listed endangered plants in SPSP, some of which are found nowhere else in the world. We also could not extrapolate an economic quantity to describe the threat posed by the hogs inhabiting SPSP as a reservoir for transmission of diseases to livestock and wildlife.

Key words: damage estimation; economic value; exotic species; feral hogs; feral pigs; invasive species; wetland damage.

INTRODUCTION

Feral hogs (*Sus scrofa*) can be a particularly destructive exotic species in the places where they have been introduced (U.S. Department of Agriculture 1999). They impose a variety of negative environmental impacts through habitat degradation, predation on native species, and competition with native species (Chequenot et al. 1996, Taft 1999, Engeman et al. in press^a). Florida, along with Hawaii, is one of the two states of the United States with the most severe invasive species problems (U.S. Congress 1993), and hogs were one of the first invasive exotic species to take hold in Florida. Florida is where hogs were first introduced to North America, by DeSoto in 1539 (Towne and Wentworth 1950). The species possesses the highest reproductive potential of any large mammal in North America (Wood and Barrett 1979, Hellgren 1999) and, with subsequent introductions, the range of feral hogs in the U.S. continues to expand (Gipson et al.

1997). Feral hogs currently inhabit many areas in such large numbers that they adversely impact wildland and agricultural ecosystems.

Feral hogs often are the single greatest vertebrate modifier of natural plant communities (Bratton 1977, Wood and Barrett 1979, Stone and Keith 1987, Engeman et al. in press^a). Rooting may damage population structures of plants, set back succession, and change species composition (Bratton 1977). In addition, mounting evidence indicates that hogs help spread rootrot fungus (*Phytophthora cinnamomi*), which causes dieback disease in native vegetation (Kliejunas and Ko 1976). Habitat damage by hogs is most pronounced in wet environments (e.g., Chequenot et al. 1996, Engeman et al. in press^a). The exposed portions of basin marshes are particularly susceptible, because they are large, irregular, wet basins with outlets only during high water. Their shallow waters are dominated by herbs and shrubs (Florida Natural Areas Inventory 1990), making them very attractive for



PLATE 1. Feral hogs "rooting" in basin marsh at Savannas Preserve State Park. Image by R.M. Engeman, NWRC.

foraging by hogs. Water generally stands in the basins for only about 200 days per year, and they are associated with and often grade into wet prairies or marsh lakes (Florida Natural Areas Inventory 1990).

Funding to manage feral hogs and restore habitat is finite and must be carefully managed to optimize the positive impact on the protected resources. Decisions on management actions towards destructive invasive species are based on economic constraints, but the metric for success of management actions is measured in resource quality. Therefore, a means to efficiently estimate feral hog damage to habitats, and to apply a monetary value to the damage, would permit economic analyses to help guide and evaluate management actions. Here, we quantify habitat damage by feral hogs to the exposed portions of the only significant basin marsh undisturbed by human habitat conversions in southeastern Florida. We also apply economic valuations to that damage (Engeman et al. in press⁴).

METHODS

The Basin Marsh

Savannas Preserve State Park (SPSP) protects the last remnant of extensive freshwater basin marsh systems that formerly extended for about 320 km along Florida's east coast. The SPSP basin marsh is unique in that water levels fluctuate dramatically from year to year, more so than most basin marshes (Florida Department of Environmental Protection, unpublished data). The basin marsh in SPSP occurs in the form of a band about 0.6 to 1.3 km wide immediately west of the Atlantic Coastal Ridge for the length of the property (approximately 15.5 km). Within the 570 ha of basin marsh, a mixture of grasses and sedges in shallow, open water dominates approximately 75% of the marsh, and sawgrass stands (*Cladium jamaicense*) cover the remaining 25%. Because of its ephemeral nature, the SPSP basin marsh aquatic




PLATE 2. Close-up of feral hog "rooting" damage in basin marsh at Savannas Preserve State Park. Image by R.M. Engeman, NWRC.

plant and animal communities are extremely sensitive to environmental disturbances. The fluctuating nature and gentle slope of the marsh also result in a wide, wet margin vegetated with forbs and grasses. This exposed zone is extremely attractive to hogs for foraging and it is highly vulnerable to damage.

SPSP supports at least 516 species of vascular plants, 192 species of vertebrates, and 300 species of aquatic invertebrates (Florida Department of Environmental Protection, unpublished data). Thirty-five of these species are listed by the U.S. Fish and Wildlife Service, Florida Fish and Wildlife Conservation Commission, or the Florida Department of Agriculture and Consumer Services as endangered, threatened, or of special concern (Florida Department of Environmental Protection 2003).

Study Site and Damage Measurements

During January 2003, damage was sampled at 100 m intervals for 5.9 km along the periphery of the basin

marsh. At each sample site, a tape measure transect was placed along the perpendicular distance from the ecotone defined by the water's edge to the ecotone between the marsh and upland vegetation of the surrounding mesic flatwoods community (Hartman 1978, Kautz 1987). Both ecotones that defined the limits of the exposed portion of basin marsh were abrupt and easy to define. After placing the tape measure, three damage measurements were made. The first was the total distance under the tape that was damaged by hogs. This amount could represent a single patch of damage or the combined distances under the tape of multiple patches. Damage not lying directly under the tape was not recorded. Hog damage was defined as ground overturned during foraging (rooting) activity. Tracks verified the species responsible. Armadillos (*Dasypus novemcinctus*) were the only other species in the park that could produce superficially similar (small) patches of damage, which were easily distinguished from hog damage by examining tracks and whether the ground was overturned, or dug by forefoot. The other two damage measures were binary and aimed at evaluating the damage precisely at the two ecotones. The point at which the tape contacted the water's edge was recorded as damaged or undamaged, as was the point at which the tape contacted the upland ecotone.

Data Analyses

The percent damaged at each sample site was calculated as the ratio of the distance along the tape that damage patches comprised, divided by the total distance between the two ecotones. The percent damaged along the basin marsh was calculated as the mean percent damaged among the sample sites ($n = 60$). The total exposed area of the study area was calculated as the mean distance between ecotones across the sample sites multiplied by the 5.9 km of shoreline. The damage rate along the shoreline ecotone was calculated as the percent of sample sites with damage at the water's edge. The damage rate at the upland ecotone was calculated similarly. Cochran's Q test for single-factor repeated measures designs with dichotomous data (e.g., Winer 1971) was used to examine whether the rate of damage was different at the two ecotones.

Damage Valuation

Determination of monetary values for protected habitats is not a straight-forward nor precise process. A means of applying a monetary value on a unit-area basis to damaged native habitats was needed to estimate the unit (per ha) and total cost of hog damage (Engeman et al. in press^a). Engeman et al. (in press^b) also discussed a variety of ways to apply monetary values to threatened and endangered animal species. Analogies to these methodologies were considered for application to habitat values, as well as other avenues specific to habitat issues. One simplistic consideration for valuation of habitat is to appraise the

land on the basis of market value. However, special habitats such as wetlands have limited "market value", and if such habitat is selectively protected, the market value diminishes even further (King 1998). The use of contingent valuation surveys for special habitats, analogous to those applied to endangered animals (Engeman et al. in press^b), tend to be even more abstract appraisals of value (King 1998). Estimated costs for restoring habitat to pristine condition (replacement costs) frequently produce values well in excess of the public's "willingness-to-pay", and therefore also do not represent a realistic valuation. The most defensible, logical, and applicable valuation for the damaged habitat characteristic of our study site was to use expenditure data for permitted wetland mitigation projects in the United States (Engeman et al. in press^a). Such data represent an empirical demonstration of willingness-to-pay value. King (1998) presented the dollar amounts per unit-area spent in efforts to restore a spectrum of wetland habitat types. The numbers represent the dollar amounts that environmental regulators have allowed permit applicants to spend in attempts to replace lost wetland services and values (King 1998). We identified the dollar value for the appropriate wetland habitat category from each of the two studies in King (1998). For these two studies, the habitats from our study areas would be classified as "freshwater emergent" or "open-water emergent", with respective 1997 empirical willingness-to-pay dollar values of \$207,480/ha and \$676,039/ha (King 1998). The 2003 values for each of these willingness-to-pay dollar amounts after adjusting for a 3% annual rate of inflation (Zerbe and Dively 1994) were \$247,742/ha and \$807,226/ha, respectively. The cost of the total area in our study site damaged by hogs was calculated by multiplying the above values by the estimated area of hog damage (Engeman et al. in press^a).

RESULTS

Damage Estimates

The mean width of the exposed portion of the basin marsh was 44.7 m (SE = 4.9 m). Multiplying this value by the 5.9 km distance of marsh periphery sampled resulted in an estimated total area of exposed basin marsh in our study site of 26.4 ha. The mean percent width of exposed marsh that was damaged by feral hogs was 19.0%, producing an estimated 5.0 ha of exposed basin marsh damaged by hogs. The total value of the damaged area was \$1,238,710 when using the "freshwater emergent" valuation and \$4,036,130 when using the "open-water emergent" valuation.

Seventy percent of the sample sites showed damage at the water's edge ecotone, versus 58% at the upland shrub ecotone. However, this difference in damage rates was not detectable statistically (χ^2 , 1 df = 1.96, $p = 0.16$).

DISCUSSION

In Florida, a premium is placed on sanctuaries for protection and preservation of habitats and species, especially because much of the natural habitat in Florida already has been lost to development. We found a substantial proportion (19%) of the exposed portion of the basin marsh to be damaged by feral hogs. In estimating the monetary values of the hog damage to the habitat we assumed standard costs for restoration. The total value of the damaged area just within our study site ranged from \$1.2 million to \$4.0 million for the two wetland valuation classifications. The periphery of the entire basin marsh would be about five times our study site. These values are substantially underestimated if the potential economic spillover effects of feral hog management are extended to incorporate endangered and threatened species, water quality impacts, suburban development areas, agricultural lands, and the transmission of diseases. These spillover effects, or secondary multiplier effects, represent the indirect benefits to society of feral hog management through additional benefits of management that were not necessarily intended directly as an outcome of the project (Boardman et al. 1996).

To expand on the above effects, consider that SPSP is home to a myriad of threatened and endangered plant (and animal) species, as well as their unique habitats (e.g., Hartman 1978, Gann et al. 2002), some of which are found nowhere else in the world (we do not identify the species or the specific areas in which these species are found to protect them from illegal collection). Damage to, or loss of, such a species is much more difficult to ascribe a monetary value to (Engeman et al. in press^b). In fact, consider that values of endangered or threatened species have been deemed "incalculable" in U.S. Supreme Court case law (Tennessee Valley Authority vs. Hill, 1978). The opinion goes so far as to say "it would be difficult for a court to balance the loss of a sum certain - even \$100 million - against a congressionally declared 'incalculable' value, even assuming we had the power to engage in such a weighing process, which we emphatically do not."

A further consideration of the magnitude of potential damage costs by hogs is that a further level of magnification occurs if the definition is broadened to include the potential for transmission of diseases to livestock and wildlife. Feral hogs can harbor a number of diseases transmissible to livestock and/or humans (e.g., Becker et al. 1978, Choquenot et al. 1996, Conger et al. 1999, Romero and Meade 1999, Taft 1999), and SPSP is situated at interfaces with rural lands, native habitats, and suburban development. In particular, the commercial hog industry in the USA has nearly eradicated swine brucellosis and pseudorabies, but feral hogs serve as a potential reservoir from which these diseases can be transmitted back to domestic stock (Taft 1999, Taylor 1999). The

hogs damaging the SPSP basin marsh are geographically a potential reservoir for diseases that could be transmitted to domestic livestock in the area.

Thus, while we have provided monetary estimates on the value of hog damage to the exposed portion of the basin marsh, these values should be viewed as a lower bound on the value of potential ecological damage. As our study concluded, SPSP had commenced a contract with the U.S. Department of Agriculture/Wildlife Services for intensive hog removal throughout the park. The cost of this contract was \$7,500, and represents only a minor fraction of the value of the hog damage to an average single ha of the exposed basin marsh, let alone to the synergistic value of the hog damage.

ACKNOWLEDGEMENTS

K. Fagerstone, N. Lance, and J.R. Mason provided insightful reviews of earlier drafts of this manuscript.

LITERATURE CITED

- Becker, H.N., R.C. Belden, T. Brevault, M.J. Burrige, W.B. Frankenberger, and P. Nicoletti P. 1978. Brucellosis in feral swine in Florida. *Journal of the American Veterinary Medical Association* 173:1181-1182.
- Boardman, A.E., D.H. Greenberg, A.R. Vining, and D.L. Weimer. 1996. *Cost-Benefit Analysis: Concepts and Practice*. Prentice Hall, Upper Saddle River, New Jersey. 493 pp.
- Bratton, S.P. 1977. Wild hogs in the United States - origin and nomenclature. Pp. 1-4 in *Research and Management of Wild Hog Populations* (G.W. Wood, ed.). The Belle W. Baruch Forest Science Institute of Clemson University, Georgetown, South Carolina.
- Choquenot, D., J. Mellroy, and T. Korn. 1996. *Managing Vertebrate Pests: Feral Pigs*. Canberra, ACT: Bureau of Resource Sciences, Australian Government Publishing Service. 163 pp.
- Conger, T.H., E. Young, and R.A. Heckmann. 1999. *Brucella suis* in feral swine. Pp. 98-107 in *Proceedings of the 1999 National Feral Swine Symposium*, Austin, Texas: Texas Animal Health Commission. 149 pp.
- Engeman, R.M., H.T. Smith, S.A. Shwiff, B. Constantin, J. Wollard, M. Nelson, and D. Griffin. In press^a. Estimating the prevalence and value of feral swine damage to native habitat in three Florida state parks. *Environmental Conservation* 30(4).
- Engeman, R.M., S.A. Shwiff, H.T. Smith, and B. Constantin. In press^b. Monetary valuation methods for economic analysis of the benefit-costs of protecting rare wildlife species from predators. *Integrated Pest Management Reviews*.
- Florida Department of Environmental Protection. 2003. *Savannas Preserve State Park Management Plan*.

- Florida Department of Environmental Protection, Florida Park Service, Tallahassee, Florida. 34 pp.
- Florida Natural Areas Inventory. 1990. Guide to the natural communities of Florida. Florida Natural Areas Inventory and Florida Department of Natural Resources, Tallahassee, Florida. 111 pp.
- Gann, G.D., K.A. Bradley, and S.W. Woodmansec. 2002. Rare Plants of South Florida: Their History, Conservation, and Restoration. Institute for Regional Conservation, Miami, Florida. 1056 pp.
- Gipson, P.S., B. Hlavachick, T. Berger, and C.D. Lee. 1997. ~~Explanations for recent range expansions~~ by wild hogs into midwestern states. Great Plains Wildlife Damage Control Workshop 13:148-150.
- Hartman, B.J. 1978. Description of major terrestrial and wetland habitats of Florida. Pp. xvi-xix in *Rare and Endangered Biota of Florida*, Vol. 2: Birds (H.W. Kale, ed.). University Presses of Florida, Gainesville, Florida. 121 pp.
- Hellgren, E. 1999. Reproduction in feral swine. Pp. 67-68 in *Proceedings of the 1999 National Feral Swine Symposium*, Austin, Texas: Texas Animal Health Commission. 149 pp.
- Kautz, R. 1987. Preliminary cover types for Landsat inventory. Florida Game and Freshwater Fish Commission, Office of Environmental Services, Nongame Wildlife Section.
- King, D. 1998. The dollar value of wetlands: trap set, bait taken, don't swallow. *National Wetlands Newsletter*. July-August: 7-11.
- Kliejunas, J.T., and W.H. Ko. 1976. Dispersal of *Phytophthora cinnamomi* on the island of Hawaii. *Phytopathology* 66:457-460.
- Romero, C.H., and P.N. Meade. 1999. Pseudorabies virus in feral swine: a research update. Pp. 73-74 in *Proceedings of the 1999 National Feral Swine Symposium*, Austin, Texas: Texas Animal Health Commission. 149 pp.
- Stone, C.P., and J.O. Keith. 1987. Control of feral ungulates and small mammals in Hawaii's National Parks: research and management strategies. Pp. 277-287 in *Control of Mammal Pests* (C.G.H. Richards and T.Y. Ku, eds.). Taylor and Francis, London.
- Taft, A.C. 1999. Feral swine - national concerns. Pp. 25-26 in *Proceedings of the 1999 National Feral Swine Symposium*, Austin, Texas: Texas Animal Health Commission. 149 pp.
- Taylor, R.B. 1999. Seasonal diets and food habits of feral swine. Pp. 58-66 in *Proceedings of the 1999 National Feral Swine Symposium*, Austin, Texas: Texas Animal Health Commission. 149 pp.
- Tennessee Valley Authority vs. Hill. (1978) 437 U.S. 153.
- Towne, C.W., and E.N. Wentworth. 1950. Pigs from Cave to Corbelt. Norman, Oklahoma, University of Oklahoma Press. 305 pp.
- U.S. Congress. 1993. Harmful Non-indigenous Species in the United States. Office of Technology Assessment, OTA-F-565, Government Printing Office, Washington, D.C. 391 pp.
- U.S. Department of Agriculture. 1999. Wild pigs hidden danger for farmers and hunters. *Animal and Plant Health Inspection Services Information Bulletin* 620. Washington, D.C. 7 pp.
- Winer, B.J. 1971. *Statistical Principles in Experimental Design*. McGraw-Hill, NY. 907 pp.
- Wood, G.W., and R.H. Barrett. 1979. Status of wild pigs in the United States. *Wildlife Society Bulletin* 7:237-246.
- Zerbe, R.O., and D.D. Dively. 1994. Benefit-Cost Analysis: In Theory and Practice. HarperCollins College Publishers, New York, NY. 557 pp.

SUGGESTED CITATION FOR THIS REPORT

- Engeman, R.M., H.T. Smith, R.G. Severson, M.A.M. Severson, S.A. Shwiff, B. Constantin, and D. Griffin. 2003. Amount and Economic Valuation of Feral Hog Damage to a Unique Basin Marsh Wetland in Florida. Florida Park Service "Parknership" Technical Report. 5 pp.