University of Nebraska - Lincoln Digital Commons@University of Nebraska - Lincoln

Papers in Natural Resources

Natural Resources, School of

12-1-2008

Measuring and modeling CO_2 and H_2O fluxes in complex terrain

Diego Andrés Riveros-Iregui University of Nebraska - Lincoln, driveros2@unl.edu

Brian L. McGlynn Montana State University - Bozeman

Follow this and additional works at: http://digitalcommons.unl.edu/natrespapers



Part of the Natural Resources and Conservation Commons

Riveros-Iregui, Diego Andrés and McGlynn, Brian L., "Measuring and modeling CO2 and H2O fluxes in complex terrain" (2008). Papers in Natural Resources. Paper 212.

http://digitalcommons.unl.edu/natrespapers/212

This Article is brought to you for free and open access by the Natural Resources, School of at Digital Commons@University of Nebraska - Lincoln. It has been accepted for inclusion in Papers in Natural Resources by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.



Measuring and modeling CO₂ and H₂O fluxes in complex terrain

Diego A. Riveros-Iregui and Brian L. McGlynn

The feedbacks between the water and the carbon cycles are of critical importance to global carbon balances. Forests and forest soils in northern latitudes are important carbon pools because of their potential as sinks for atmospheric carbon. However there are significant unknowns related to the effects of hydrologic variability, mountainous terrain, and landscape heterogeneity in controlling soil carbon dioxide (CO₂) efflux. Mountainous terrain imposes large spatial heterogeneity in the biophysical controls of soil CO2 production and efflux, including soil temperature, soil water content, vegetation, substrate, and soil physical properties. Further complications are introduced by the superimposed temporal heterogeneity (i.e., the asynchronous response of each variable to changes in environmental conditions). As a result, extrapolating from single- or multiple-point measurements to larger areas requires understanding of the emerging patterns controlled by the underlying spatiotemporal nature of biophysical drivers.

At the Tenderfoot Creek Experimental Forest (TCEF) in central Montana, two factors that we use to our advantage in the understanding of watershedscale soil CO2 efflux are I) the temporal seasonality imposed by snowmelt and 2) the spatial redistribution of soil water imposed by topography. Snowmelt controls the timing of the most dramatic increase in water content in the soil, while landscape morphology redistributes that moisture down slope to lower areas of the landscape. Our site selection (62 soil respiration plots) targeted those areas of the landscape that offered natural biophysical gradients (Figure 1). Two eddy covariance systems were installed over the




Figure 1: Tenderfoot Creek Experimental Forest (TCEF) in the Little Belt Mountains of Central Montana. Soil respiration is measured at 62 sites distributed across the catchment.



Photo: Riparian meadow

canopy of the two most important systems: a riparian meadow and a lodgepole pine forest Additionally, we (Figure 2). investigate surface water groundwater interactions and subsurface flowpaths between different landscape positions and the stream using an array of 80+ groundwater wells and piezometers (~I-2 m depth). Our coupled water-carbon approach is based on the concept of topographic similarity [Beven and Kirkby, 1979], which hydrologists and biogeochemists have used to transfer process and response understanding to topographically and thus hydrologically and biogeochemically similar areas [e.g., Creed et al., 1996; Boyer et al., 1997; McGlynn and McDonnell, 2003]. This idea is conceptually intuitive because I) many biogeochemical processes are mediated by temperature, water content, radiation, and energy balance, variables that often vary predictably with topographic position; and 2) this form of heterogeneity also depends on other abiotic factors (e.g., slope,

soil type, upslope accumulated area), which can be considered static over relevant times scales [Moorcroft, 2006]. As such, topographic similarity can help scale soil CO2 efflux rates from single- or multiple-point measurements to watershed scales or larger areas. Determining the minimum set of watershed measurements or variables needed to characterize soil CO2 efflux both spatially and temporally is not trivial. However, terrain analysis techniques can help link spatial watershed patterns with biogeochemical processes, aid in transfer and interpolation, and indicate where additional field observations are needed. process knowledge gained from such observations can help characterize the landscape, discretely or continuously, as an arrangement of response characteristics and thresholds.

Current work at the TCEF based on multiple-point measurements has demonstrated that the spatial variability of soil CO₂ efflux across the 62 sites is higher than previously thought (Figure 3). Riveros-Iregui et al.

cont. on page 13



Measuring and modeling CO₂ and H₂O fluxes

cont. from page 12




Figure 2: LIDAR-derived image of vegetation cover at the TCEF.

[in press] demonstrated that more persistent, high soil water content was the major control on the spatial difference of soil CO2 efflux and nighttime ecosystem respiration between riparian-hillslope sites especially after snowmelt or rainfall. Pacific et al. [2008] analyzed soil CO₂ efflux measurements across riparian-hillslopes transitions based on soil gas wells and discrete chamber measurements. Their results show that soil CO₂ efflux rates differ, both in magnitude and timing, across riparianhillslope transitions. While early in the growing season soil CO2

efflux is higher in hillslopes than in riparian areas, later in the season soil CO2 efflux from riparian areas becomes higher than in hillslopes. Our preliminary results demonstrate how landscape discretization and the concept of topographic similarity can help extend understanding and measurements of soil CO2 efflux based on benchmark measurements to larger areas of the landscape. There remain great challenges in dealing with complex topography, especially transferring knowledge acquired at point/plot scales to larger spatial scales, and reconciling C




Figure 3: Comparison of cumulative CO_2 fluxes for the 2006 growing season at 62 sites (gray lines) and ecosystem respiration (R_E) measured with the EC technique over a riparian meadow (blue line) and the lodgepole pine forest (red line). Range of soil respiration varies by us much as ~600% across the catchment. Terrain analysis as well as process understanding of the organized heterogeneity are critical in up-scaling plot measurements of soil respiration to spatial scales comparable to towers.

fluxes measured at different levels of the ecosystem. However, based on initial results of coupled water-carbon studies in the subalpine forests of the TCEF, we recommend implementation and further investigation of the concept of "organized heterogeneity" as a responsible agent in the emergent patterns of these fluxes from the plot to the watershed to regional scales.

contact: Diego Riveros-Iregui University of Colorado, Boulder, riveros@Colorado.edu

Brian McGlynn

Montana State University-Bozeman

bmcglynn@montana.edu

Literature

Beven, K. J., and M. J. Kirkby (1979), A physically based, variable contributing area model of basin hydrology. *Hydrological Sciences Bulletin*, 24(1), 43-69.

Boyer, E. W., et al. (1997), Response characteristics of DOC flushing in an alpine catchment, *Hydrological Processes*, 11(12), 1635-1647.

Creed, I. F., et al. (1996), Regulation of nitrate-N release from temperate forests: A test of the N flushing hypothesis, *Water Resources Research*, 32(11), 3337-3354.

McGlynn, B. L., and J. J. McDonnell (2003), Role of discrete landscape units in controlling catchment dissolved organic carbon dynamics, *Water Resources Research*, 39(4), 1090,

Moorcroft, P. R. (2006), How close are we to a predictive science of the biosphere?, *Trends in Ecology & Evolution*, 21 (7), 400-407.


Pacific, V. J., B. L. McGlynn, D. A. Riveros-Iregui, D. L. Welsch, and H. E. Epstein (2008), Variability in soil respiration across riparian-hillslope transitions, *Biogeochemistry*, 91:51-70.

Riveros-Iregui, D. A., B. L. McGlynn, H. E. Epstein, and D. L. Welsch (in press), Interpretation and evaluation of combined measurement techniques for soil CO₂ efflux: surface chambers and soil CO₂ concentration probes, *Journal of Geophysical Research-Biogeosciences*

FluxLetter The Newsletter of FLUXNET

Vol.1 No.4 December, 2008

FluxLetter is produced quarterly at the FLUXNET Office with support from the National Science Foundation.



This issue of FluxLetter was edited, designed and produced by:

Rodrigo Vargas Dennis Baldocchi

FLUXNET Office, 137 Mulford Hall, University of California, Berkeley, CA 94720 ph: 1-(510)-642-2421 Fax: 1-(510)-643-5098

We plan to make the FLUXNET newsletter a powerful information, networking, and communication resource for the community. If you want to contribute to any section or propose a new one please contact the FLUXNET Office. THANKS!!