

8-3-2007

Direct Observation of the Strange b Baryon Ξ_b^-

V.M. Abazov

Joint Institute for Nuclear Research, Dubna, Russia

Kenneth A. Bloom

University of Nebraska - Lincoln, kbloom2@unl.edu

Gregory R. Snow

University of Nebraska-Lincoln, gsnow1@unl.edu

D0 Collaboration

Follow this and additional works at: <http://digitalcommons.unl.edu/physicsbloom>



Part of the [Physics Commons](#)

Abazov, V.M.; Bloom, Kenneth A.; Snow, Gregory R.; and Collaboration, D0, "Direct Observation of the Strange b Baryon Ξ_b^- " (2007). *Kenneth Bloom Publications*. 218.

<http://digitalcommons.unl.edu/physicsbloom/218>

This Article is brought to you for free and open access by the Research Papers in Physics and Astronomy at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Kenneth Bloom Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.



Direct Observation of the Strange b Baryon Ξ_b^-

V. M. Abazov,³⁵ B. Abbott,⁷⁵ M. Abolins,⁶⁵ B. S. Acharya,²⁸ M. Adams,⁵¹ T. Adams,⁴⁹ E. Aguilo,⁵ S. H. Ahn,³⁰ M. Ahsan,⁵⁹ G. D. Alexeev,³⁵ G. Alkhalaf,³⁹ A. Alton,^{64,*} G. Alverson,⁶³ G. A. Alves,² M. Anastasoae,³⁴ L. S. Ancu,³⁴ T. Andeen,⁵³ S. Anderson,⁴⁵ B. Andrieu,¹⁶ M. S. Anzels,⁵³ Y. Arnoud,¹³ M. Arov,⁶⁰ M. Arthaud,¹⁷ A. Askew,⁴⁹ B. Åsman,⁴⁰ A. C. S. Assis Jesus,³ O. Atramentov,⁴⁹ C. Autermann,²⁰ C. Avila,⁷ C. Ay,²³ F. Badaud,¹² A. Baden,⁶¹ L. Bagby,⁵² B. Baldin,⁵⁰ D. V. Bandurin,⁵⁹ S. Banerjee,²⁸ P. Banerjee,²⁸ E. Barberis,⁶³ A.-F. Barfuss,¹⁴ P. Bargassa,⁸⁰ P. Baringer,⁵⁸ J. Barreto,² J. F. Bartlett,⁵⁰ U. Bassler,¹⁶ D. Bauer,⁴³ S. Beale,⁵ A. Bean,⁵⁸ M. Begalli,³ M. Begel,⁷¹ C. Belanger-Champagne,⁴⁰ L. Bellantoni,⁵⁰ A. Bellavance,⁵⁰ J. A. Benitez,⁶⁵ S. B. Beri,²⁶ G. Bernardi,¹⁶ R. Bernhard,²² L. Berntzon,¹⁴ I. Bertram,⁴² M. Bessaon,¹⁷ R. Beuselinck,⁴³ V. A. Bezzubov,³⁸ P. C. Bhat,⁵⁰ V. Bhatnagar,²⁶ C. Biscarat,¹⁹ G. Blazey,⁵² F. Blekman,⁴³ S. Blessing,⁴⁹ D. Bloch,¹⁸ K. Bloom,⁶⁷ A. Boehnlein,⁵⁰ D. Boline,⁶² T. A. Bolton,⁵⁹ G. Borissov,⁴² K. Bos,³³ T. Bose,⁷⁷ A. Brandt,⁷⁸ R. Brock,⁶⁵ G. Brooijmans,⁷⁰ A. Bross,⁵⁰ D. Brown,⁷⁸ N. J. Buchanan,⁴⁹ D. Buchholz,⁵³ M. Buehler,⁸¹ V. Buescher,²¹ S. Burdin,^{42,†} S. Burke,⁴⁵ T. H. Burnett,⁸² C. P. Buszello,⁴³ J. M. Butler,⁶² P. Calfayan,²⁴ S. Calvet,¹⁴ J. Cammin,⁷¹ S. Caron,³³ W. Carvalho,³ B. C. K. Casey,⁷⁷ N. M. Cason,⁵⁵ H. Castilla-Valdez,³² S. Chakrabarti,¹⁷ D. Chakraborty,⁵² K. M. Chan,⁵⁵ K. Chan,⁵ A. Chandra,⁴⁸ F. Charles,¹⁸ E. Cheu,⁴⁵ F. Chevallier,¹³ D. K. Cho,⁶² S. Choi,³¹ B. Choudhary,²⁷ L. Christofek,⁷⁷ T. Christoudias,⁴³ S. Cihangir,⁵⁰ D. Claes,⁶⁷ C. Clément,⁴⁰ B. Clément,¹⁸ Y. Coadou,⁵ M. Cooke,⁸⁰ W. E. Cooper,⁵⁰ M. Corcoran,⁸⁰ F. Couderc,¹⁷ M.-C. Cousinou,¹⁴ S. Crépe-Renaudin,¹³ D. Cutts,⁷⁷ M. Ćwiok,²⁹ H. da Motta,² A. Das,⁶² G. Davies,⁴³ K. De,⁷⁸ S. J. de Jong,³⁴ P. de Jong,³³ E. De La Cruz-Burelo,⁶⁴ C. De Oliveira Martins,³ J. D. Degenhardt,⁶⁴ F. Déliot,¹⁷ M. Demarteau,⁵⁰ R. Demina,⁷¹ D. Denisov,⁵⁰ S. P. Denisov,³⁸ S. Desai,⁵⁰ H. T. Diehl,⁵⁰ M. Diesburg,⁵⁰ A. Dominguez,⁶⁷ H. Dong,⁷² L. V. Dudko,³⁷ L. Duflot,¹⁵ S. R. Dugad,²⁸ D. Duggan,⁴⁹ A. Duperrin,¹⁴ J. Dyer,⁶⁵ A. Dyshkant,⁵² M. Eads,⁶⁷ D. Edmunds,⁶⁵ J. Ellison,⁴⁸ V. D. Elvira,⁵⁰ Y. Enari,⁷⁷ S. Eno,⁶¹ P. Ermolov,³⁷ H. Evans,⁵⁴ A. Evdokimov,⁷³ V. N. Evdokimov,³⁸ A. V. Ferapontov,⁵⁹ T. Ferbel,⁷¹ F. Fiedler,²⁴ F. Filthaut,³⁴ W. Fisher,⁵⁰ H. E. Fisk,⁵⁰ M. Ford,⁴⁴ M. Fortner,⁵² H. Fox,²² S. Fu,⁵⁰ S. Fuess,⁵⁰ T. Gadfort,⁸² C. F. Galea,³⁴ E. Gallas,⁵⁰ E. Galyaev,⁵⁵ C. Garcia,⁷¹ A. Garcia-Bellido,⁸² V. Gavrilov,³⁶ P. Gay,¹² W. Geist,¹⁸ D. Gelé,¹⁸ C. E. Gerber,⁵¹ Y. Gershtein,⁴⁹ D. Gillberg,⁵ G. Ginter,⁷¹ N. Gollub,⁴⁰ B. Gómez,⁷ A. Goussiou,⁵⁵ P. D. Grannis,⁷² H. Greenlee,⁵⁰ Z. D. Greenwood,⁶⁰ E. M. Gregores,⁴ G. Grenier,¹⁹ Ph. Gris,¹² J.-F. Grivaz,¹⁵ A. Grohsjean,²⁴ S. Grünendahl,⁵⁰ M. W. Grünwald,²⁹ J. Guo,⁷² F. Guo,⁷² P. Gutierrez,⁷⁵ G. Gutierrez,⁵⁰ A. Haas,⁷⁰ N. J. Hadley,⁶¹ P. Haefner,²⁴ S. Hagopian,⁴⁹ J. Haley,⁶⁸ I. Hall,⁷⁵ R. E. Hall,⁴⁷ L. Han,⁶ K. Hanagaki,⁵⁰ P. Hansson,⁴⁰ K. Harder,⁴⁴ A. Harel,⁷¹ R. Harrington,⁶³ J. M. Hauptman,⁵⁷ R. Hauser,⁶⁵ J. Hays,⁴³ T. Hebbeker,²⁰ D. Hedin,⁵² J. G. Hegeman,³³ J. M. Heinmiller,⁵¹ A. P. Heinson,⁴⁸ U. Heintz,⁶² C. Hensel,⁵⁸ K. Herner,⁷² G. Hesketh,⁶³ M. D. Hildreth,⁵⁵ R. Hirsch,⁸¹ J. D. Hobbs,⁷² B. Hoeneisen,¹¹ H. Hoeth,²⁵ M. Hohlheid,²¹ S. J. Hong,³⁰ R. Hooper,⁷⁷ S. Hossain,⁷⁵ P. Houben,³³ Y. Hu,⁷² Z. Hubacek,⁹ V. Hynek,⁸ I. Iashvili,⁶⁹ R. Illingworth,⁵⁰ A. S. Ito,⁵⁰ S. Jabeen,⁶² M. Jaffré,¹⁵ S. Jain,⁷⁵ K. Jakobs,²² C. Jarvis,⁶¹ R. Jesik,⁴³ K. Johns,⁴⁵ C. Johnson,⁷⁰ M. Johnson,⁵⁰ A. Jonckheere,⁵⁰ P. Jonsson,⁴³ A. Juste,⁵⁰ D. Käfer,²⁰ S. Kahn,⁷³ E. Kajfasz,¹⁴ A. M. Kalinin,³⁵ J. R. Kalk,⁶⁵ J. M. Kalk,⁶⁰ S. Kappler,²⁰ D. Karmanov,³⁷ J. Kasper,⁶² P. Kasper,⁵⁰ I. Katsanos,⁷⁰ D. Kau,⁴⁹ R. Kaur,²⁶ V. Kaushik,⁷⁸ R. Kehoe,⁷⁹ S. Kermiche,¹⁴ N. Khalatyan,³⁸ A. Khanov,⁷⁶ A. Kharchilava,⁶⁹ Y. M. Kharzhev,³⁵ D. Khatidze,⁷⁰ H. Kim,³¹ T. J. Kim,³⁰ M. H. Kirby,³⁴ M. Kirsch,²⁰ B. Klima,⁵⁰ J. M. Kohli,²⁶ J.-P. Konrath,²² M. Kopal,⁷⁵ V. M. Korablev,³⁸ B. Kothari,⁷⁰ A. V. Kozelov,³⁸ D. Krop,⁵⁴ A. Kryemadhi,⁸¹ T. Kuhl,²³ A. Kumar,⁶⁹ S. Kunori,⁶¹ A. Kupco,¹⁰ T. Kurča,¹⁹ J. Kvita,⁸ F. Lacroix,¹² D. Lam,⁵⁵ S. Lammers,⁷⁰ G. Landsberg,⁷⁷ J. Lazoflores,⁴⁹ P. Lebrun,¹⁹ W. M. Lee,⁵⁰ A. Leflat,³⁷ F. Lehner,⁴¹ J. Lellouch,¹⁶ V. Lesne,¹² J. Leveque,⁴⁵ P. Lewis,⁴³ J. Li,⁷⁸ Q. Z. Li,⁵⁰ L. Li,⁴⁸ S. M. Lietti,⁴ J. G. R. Lima,⁵² D. Lincoln,⁵⁰ J. Linnemann,⁶⁵ V. V. Lipaev,³⁸ R. Lipton,⁵⁰ Y. Liu,⁶ Z. Liu,⁵ L. Lobo,⁴³ A. Lobodenko,³⁹ M. Lokajicek,¹⁰ A. Lounis,¹⁸ P. Love,⁴² H. J. Lubatti,⁸² A. L. Lyon,⁵⁰ A. K. A. Maciel,² D. Mackin,⁸⁰ R. J. Madaras,⁴⁶ P. Mättig,²⁵ C. Magass,²⁰ A. Magerkurth,⁶⁴ N. Makovec,¹⁵ P. K. Mal,⁵⁵ H. B. Malbouisson,³ S. Malik,⁶⁷ V. L. Malyshev,³⁵ H. S. Mao,⁵⁰ Y. Maravin,⁵⁹ B. Martin,¹³ R. McCarthy,⁷² A. Melnitchouk,⁶⁶ A. Mendes,¹⁴ L. Mendoza,⁷ P. G. Mercadante,⁴ Y. P. Merekov,³⁵ M. Merkin,³⁷ K. W. Merritt,⁵⁰ J. Meyer,²¹ A. Meyer,²⁰ M. Michaut,¹⁷ T. Millet,¹⁹ J. Mitrevski,⁷⁰ J. Molina,³ R. K. Mommsen,⁴⁴ N. K. Mondal,²⁸ R. W. Moore,⁵ T. Moulik,⁵⁸ G. S. Muanza,¹⁹ M. Mulders,⁵⁰ M. Mulhearn,⁷⁰ O. Mundal,²¹ L. Mundim,³ E. Nagy,¹⁴ M. Naimuddin,⁵⁰ M. Narain,⁷⁷ N. A. Naumann,³⁴ H. A. Neal,⁶⁴ J. P. Negret,⁷ P. Neustroev,³⁹ H. Nilsen,²² A. Nomerotski,⁵⁰ S. F. Novaes,⁴ T. Nunnemann,²⁴ V. O'Dell,⁵⁰ D. C. O'Neil,⁵ G. Odrant,³⁹ C. Ochando,¹⁵ D. Onoprienko,⁵⁹ N. Oshima,⁵⁰ J. Osta,⁵⁵ R. Otec,⁹ G. J. Otero y Garzón,⁵¹ M. Owen,⁴⁴ P. Padley,⁸⁰ M. Pangilinan,⁷⁷ G. Panov,³⁵ N. Parashar,⁵⁶ S.-J. Park,⁷¹ S. K. Park,³⁰ J. Parsons,⁷⁰ R. Partridge,⁷⁷ N. Parua,⁵⁴ A. Patwa,⁷³ G. Pawloski,⁸⁰

B. Penning,²² P. M. Perea,⁴⁸ K. Peters,⁴⁴ Y. Peters,²⁵ P. Pétroff,¹⁵ M. Petteni,⁴³ R. Piegaiia,¹ J. Piper,⁶⁵ M.-A. Pleier,²¹ P. L. M. Podesta-Lerma,^{32,‡} V. M. Podstavkov,⁵⁰ Y. Pogorelov,⁵⁵ M.-E. Pol,² P. Polozov,³⁶ A. Pompoš,⁷⁵ B. G. Pope,⁶⁵ A. V. Popov,³⁸ C. Potter,⁵ W. L. Prado da Silva,³ H. B. Prosper,⁴⁹ S. Protopopescu,⁷³ J. Qian,⁶⁴ A. Quadt,²¹ B. Quinn,⁶⁶ A. Rakitine,⁴² M. S. Rangel,² K. J. Rani,²⁸ K. Ranjan,²⁷ P. N. Ratoff,⁴² P. Renkel,⁷⁹ S. Reucroft,⁶³ P. Rich,⁴⁴ M. Rijssenbeek,⁷² I. Ripp-Baudot,¹⁸ F. Rizatdinova,⁷⁶ S. Robinson,⁴³ R. F. Rodrigues,³ C. Royon,¹⁷ A. Rozhdestvenski,³⁵ P. Rubinov,⁵⁰ R. Ruchti,⁵⁵ G. Safronov,³⁶ G. Sajot,¹³ A. Sánchez-Hernández,³² M. P. Sanders,¹⁶ A. Santoro,³ G. Savage,⁵⁰ L. Sawyer,⁶⁰ T. Scanlon,⁴³ D. Schaile,²⁴ R. D. Schamberger,⁷² Y. Scheglov,³⁹ H. Schellman,⁵³ P. Schieferdecker,²⁴ T. Schliephake,²⁵ C. Schmitt,²⁵ C. Schwanenberger,⁴⁴ A. Schwartzman,⁶⁸ R. Schwienhorst,⁶⁵ J. Sekaric,⁴⁹ S. Sengupta,⁴⁹ H. Severini,⁷⁵ E. Shabalina,⁵¹ M. Shamim,⁵⁹ V. Shary,¹⁷ A. A. Shchukin,³⁸ R. K. Shivpuri,²⁷ D. Shpakov,⁵⁰ V. Siccaldi,¹⁸ V. Simak,⁹ V. Sirotenko,⁵⁰ P. Skubic,⁷⁵ P. Slattery,⁷¹ D. Smirnov,⁵⁵ R. P. Smith,⁵⁰ J. Snow,⁷⁴ G. R. Snow,⁶⁷ S. Snyder,⁷³ S. Söldner-Rembold,⁴⁴ L. Sonnenschein,¹⁶ A. Sopczak,⁴² M. Sosebee,⁷⁸ K. Soustruznik,⁸ M. Souza,² B. Spurlock,⁷⁸ J. Stark,¹³ J. Steele,⁶⁰ V. Stolin,³⁶ A. Stone,⁵¹ D. A. Stoyanova,³⁸ J. Strandberg,⁶⁴ S. Strandberg,⁴⁰ M. A. Strang,⁶⁹ M. Strauss,⁷⁵ E. Strauss,⁷² R. Ströhmer,²⁴ D. Strom,⁵³ M. Strovink,⁴⁶ L. Stutte,⁵⁰ S. Sumowidagdo,⁴⁹ P. Svoisky,⁵⁵ A. Sznajder,³ M. Talby,¹⁴ P. Tamburello,⁴⁵ A. Tanasijczuk,¹ W. Taylor,⁵ P. Telford,⁴⁴ J. Temple,⁴⁵ B. Tiller,²⁴ F. Tissandier,¹² M. Titov,¹⁷ V. V. Tokmenin,³⁵ M. Tomoto,⁵⁰ T. Toole,⁶¹ I. Torchiani,²² T. Trefzger,²³ D. Tsybychev,⁷² B. Tuchming,¹⁷ C. Tully,⁶⁸ P. M. Tuts,⁷⁰ R. Unalan,⁶⁵ S. Uvarov,³⁹ L. Uvarov,³⁹ S. Uzunyan,⁵² B. Vachon,⁵ P. J. van den Berg,³³ B. van Eijk,³³ R. Van Kooten,⁵⁴ W. M. van Leeuwen,³³ N. Varelas,⁵¹ E. W. Varnes,⁴⁵ A. Vartapetian,⁷⁸ I. A. Vasilyev,³⁸ M. Vaupel,²⁵ P. Verdier,¹⁹ L. S. Vertogradov,³⁵ Y. Vertogradova,³⁵ M. Verzocchi,⁵⁰ F. Villeneuve-Seguier,⁴³ P. Vint,⁴³ P. Vokac,⁹ E. Von Toerne,⁵⁹ M. Voutilainen,^{67,§} M. Vreeswijk,³³ R. Wagner,⁶⁸ H. D. Wahl,⁴⁹ L. Wang,⁶¹ M. H. L. S Wang,⁵⁰ J. Warchol,⁵⁵ G. Watts,⁸² M. Wayne,⁵⁵ M. Weber,⁵⁰ G. Weber,²³ H. Weerts,⁶⁵ A. Wenger,^{22,||} N. Wermes,²¹ M. Wetstein,⁶¹ A. White,⁷⁸ D. Wicke,²⁵ G. W. Wilson,⁵⁸ S. J. Wimpenny,⁴⁸ M. Wobisch,⁶⁰ D. R. Wood,⁶³ T. R. Wyatt,⁴⁴ Y. Xie,⁷⁷ S. Yacoub,⁵³ R. Yamada,⁵⁰ M. Yan,⁶¹ T. Yasuda,⁵⁰ Y. A. Yatsunenkov,³⁵ K. Yip,⁷³ H. D. Yoo,⁷⁷ S. W. Youn,⁵³ J. Yu,⁷⁸ C. Yu,¹³ A. Yurkewicz,⁷² A. Zatserklyaniy,⁵² C. Zeitnitz,²⁵ D. Zhang,⁵⁰ T. Zhao,⁸² B. Zhou,⁶⁴ J. Zhu,⁷² M. Zielinski,⁷¹ D. Zieminska,⁵⁴ A. Zieminski,⁵⁴ L. Zivkovic,⁷⁰ V. Zutshi,⁵² and E. G. Zverev³⁷

(D0 Collaboration)

¹Universidad de Buenos Aires, Buenos Aires, Argentina²LAFEX, Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil³Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil⁴Instituto de Física Teórica, Universidade Estadual Paulista, São Paulo, Brazil⁵University of Alberta, Edmonton, Alberta, Canada, Simon Fraser University, Burnaby, British Columbia, Canada, York University, Toronto, Ontario, Canada,

and McGill University, Montreal, Quebec, Canada

⁶University of Science and Technology of China, Hefei, People's Republic of China⁷Universidad de los Andes, Bogotá, Colombia⁸Center for Particle Physics, Charles University, Prague, Czech Republic⁹Czech Technical University, Prague, Czech Republic¹⁰Center for Particle Physics, Institute of Physics, Academy of Sciences of the Czech Republic, Prague, Czech Republic¹¹Universidad San Francisco de Quito, Quito, Ecuador¹²Laboratoire de Physique Corpusculaire, IN2P3-CNRS, Université Blaise Pascal, Clermont-Ferrand, France¹³Laboratoire de Physique Subatomique et de Cosmologie, IN2P3-CNRS, Université de Grenoble I, Grenoble, France¹⁴CPPM, IN2P3-CNRS, Université de la Méditerranée, Marseille, France¹⁵Laboratoire de l'Accélérateur Linéaire, IN2P3-CNRS et Université Paris-Sud, Orsay, France¹⁶LPNHE, IN2P3-CNRS, Universités Paris VI and VII, Paris, France¹⁷DAPNIA/Service de Physique des Particules, CEA, Saclay, France¹⁸IPHC, Université Louis Pasteur et Université de Haute Alsace, CNRS, IN2P3, Strasbourg, France¹⁹IPNL, Université Lyon I, CNRS/IN2P3, Villeurbanne, France

and Université de Lyon, Lyon, France

²⁰III. Physikalisches Institut A, RWTH Aachen, Aachen, Germany²¹Physikalisches Institut, Universität Bonn, Bonn, Germany²²Physikalisches Institut, Universität Freiburg, Freiburg, Germany²³Institut für Physik, Universität Mainz, Mainz, Germany²⁴Ludwig-Maximilians-Universität München, München, Germany²⁵Fachbereich Physik, University of Wuppertal, Wuppertal, Germany²⁶Panjab University, Chandigarh, India

- ²⁷Delhi University, Delhi, India
²⁸Tata Institute of Fundamental Research, Mumbai, India
²⁹University College Dublin, Dublin, Ireland
³⁰Korea Detector Laboratory, Korea University, Seoul, Korea
³¹SungKyunKwan University, Suwon, Korea
³²CINVESTAV, Mexico City, Mexico
³³FOM-Institute NIKHEF and University of Amsterdam/NIKHEF, Amsterdam, The Netherlands
³⁴Radboud University Nijmegen/NIKHEF, Nijmegen, The Netherlands
³⁵Joint Institute for Nuclear Research, Dubna, Russia
³⁶Institute for Theoretical and Experimental Physics, Moscow, Russia
³⁷Moscow State University, Moscow, Russia
³⁸Institute for High Energy Physics, Protvino, Russia
³⁹Petersburg Nuclear Physics Institute, St. Petersburg, Russia
⁴⁰Lund University, Lund, Sweden, Royal Institute of Technology and Stockholm University, Stockholm, Sweden, and Uppsala University, Uppsala, Sweden
⁴¹Physik Institut der Universität Zürich, Zürich, Switzerland
⁴²Lancaster University, Lancaster, United Kingdom
⁴³Imperial College, London, United Kingdom
⁴⁴University of Manchester, Manchester, United Kingdom
⁴⁵University of Arizona, Tucson, Arizona 85721, USA
⁴⁶Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA
⁴⁷California State University, Fresno, California 93740, USA
⁴⁸University of California, Riverside, California 92521, USA
⁴⁹Florida State University, Tallahassee, Florida 32306, USA
⁵⁰Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA
⁵¹University of Illinois at Chicago, Chicago, Illinois 60607, USA
⁵²Northern Illinois University, DeKalb, Illinois 60115, USA
⁵³Northwestern University, Evanston, Illinois 60208, USA
⁵⁴Indiana University, Bloomington, Indiana 47405, USA
⁵⁵University of Notre Dame, Notre Dame, Indiana 46556, USA
⁵⁶Purdue University Calumet, Hammond, Indiana 46323, USA
⁵⁷Iowa State University, Ames, Iowa 50011, USA
⁵⁸University of Kansas, Lawrence, Kansas 66045, USA
⁵⁹Kansas State University, Manhattan, Kansas 66506, USA
⁶⁰Louisiana Tech University, Ruston, Louisiana 71272, USA
⁶¹University of Maryland, College Park, Maryland 20742, USA
⁶²Boston University, Boston, Massachusetts 02215, USA
⁶³Northeastern University, Boston, Massachusetts 02115, USA
⁶⁴University of Michigan, Ann Arbor, Michigan 48109, USA
⁶⁵Michigan State University, East Lansing, Michigan 48824, USA
⁶⁶University of Mississippi, University, Mississippi 38677, USA
⁶⁷University of Nebraska, Lincoln, Nebraska 68588, USA
⁶⁸Princeton University, Princeton, New Jersey 08544, USA
⁶⁹State University of New York, Buffalo, New York 14260, USA
⁷⁰Columbia University, New York, New York 10027, USA
⁷¹University of Rochester, Rochester, New York 14627, USA
⁷²State University of New York, Stony Brook, New York 11794, USA
⁷³Brookhaven National Laboratory, Upton, New York 11973, USA
⁷⁴Langston University, Langston, Oklahoma 73050, USA
⁷⁵University of Oklahoma, Norman, Oklahoma 73019, USA
⁷⁶Oklahoma State University, Stillwater, Oklahoma 74078, USA
⁷⁷Brown University, Providence, Rhode Island 02912, USA
⁷⁸University of Texas, Arlington, Texas 76019, USA
⁷⁹Southern Methodist University, Dallas, Texas 75275, USA
⁸⁰Rice University, Houston, Texas 77005, USA
⁸¹University of Virginia, Charlottesville, Virginia 22901, USA
⁸²University of Washington, Seattle, Washington 98195, USA

(Received 12 June 2007; published 3 August 2007)

We report the first direct observation of the strange b baryon Ξ_b^- (Ξ_b^+). We reconstruct the decay $\Xi_b^- \rightarrow J/\psi \Xi^-$, with $J/\psi \rightarrow \mu^+ \mu^-$, and $\Xi^- \rightarrow \Lambda \pi^- \rightarrow p \pi^- \pi^-$ in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV. Using

1.3 fb⁻¹ of data collected by the D0 detector, we observe $15.2 \pm 4.4(\text{stat})_{-0.4}^{+1.9}(\text{syst}) \Xi_b^-$ candidates at a mass of $5.774 \pm 0.011(\text{stat}) \pm 0.015(\text{syst})$ GeV. The significance of the observed signal is 5.5σ , equivalent to a probability of 3.3×10^{-8} of it arising from a background fluctuation. Normalizing to the decay $\Lambda_b \rightarrow J/\psi\Lambda$, we measure the relative rate $\frac{\sigma(\Xi_b^-) \times \mathcal{B}(\Xi_b^- \rightarrow J/\psi \Xi^-)}{\sigma(\Lambda_b) \times \mathcal{B}(\Lambda_b \rightarrow J/\psi \Lambda)} = 0.28 \pm 0.09(\text{stat})_{-0.08}^{+0.09}(\text{syst})$.

DOI: 10.1103/PhysRevLett.99.052001

PACS numbers: 14.20.Mr, 13.30.Eg, 13.85.Ni

The quark model of hadrons [1] predicts the existence of a number of baryons containing b quarks, with a hierarchical structure similar to that of charmed baryons. Despite significant progress in studying b hadrons over the last decade, only the $\Lambda_b(udb)$ b baryon has been directly observed. The $\Xi_b^-(dsb)$ (charge conjugate states are assumed throughout this Letter) is a strange b baryon made of valence quarks from all three known generations of fermions and is expected to decay through the weak interaction. Theoretical calculations of heavy quark effective theory [2] and nonrelativistic QCD [3] predict the Ξ_b^- mass in the range 5.7–5.8 GeV [4].

Experiments at the CERN LEP e^+e^- collider have reported indirect evidence of the Ξ_b^- baryon based on an excess of same-sign $\Xi^- \ell^-$ events in jets [5]. Interpreting the excess as the semi-inclusive $\Xi_b^- \rightarrow \Xi^- \ell^- \bar{\nu}_\ell X$ decay, the average lifetime of the Ξ_b^- is $1.42_{-0.24}^{+0.28}$ ps [6]. In this Letter, we report the first direct observation of the Ξ_b^- baryon, fully reconstructed in an exclusive decay. We observe the decay $\Xi_b^- \rightarrow J/\psi \Xi^-$, with $J/\psi \rightarrow \mu^+ \mu^-$, $\Xi^- \rightarrow \Lambda \pi^-$, and $\Lambda \rightarrow p \pi^-$. The analysis is based on a data sample of 1.3 fb⁻¹ integrated luminosity collected in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV with the D0 detector at the Fermilab Tevatron collider during 2002–2006.

The D0 detector is described in detail elsewhere [7]. The components most relevant to this analysis are the central tracking system and the muon spectrometer. The central tracking system consists of a silicon microstrip tracker (SMT) and a central fiber tracker (CFT) that are surrounded by a 2 T superconducting solenoid. The SMT is optimized for tracking and vertexing for the pseudorapidity region $|\eta| < 3$ ($\eta = -\ln[\tan(\theta/2)]$ and θ is the polar angle) while the CFT has coverage for $|\eta| < 2$. Liquid-argon and uranium calorimeters in a central and two end-cap cryostats cover the pseudorapidity region $|\eta| < 4.2$. The muon spectrometer is located outside the calorimeter and covers the pseudorapidity region $|\eta| < 2$. It comprises a layer of drift tubes and scintillator trigger counters in front of 1.8 T iron toroids followed by two similar layers behind the toroids.

The topology of $\Xi_b^- \rightarrow J/\psi \Xi^- \rightarrow J/\psi \Lambda \pi^-$ decay (see Fig. 1) is similar to that of the $\Lambda_b \rightarrow J/\psi \Lambda$ decay; therefore, the reconstruction of the J/ψ and Λ and their selection discussed below are guided by the strategies applied to the Λ_b lifetime measurement in D0 [8]. They are then validated with simulated Monte Carlo (MC) Ξ_b^- events. The PYTHIA MC program [9] is used to generate Ξ_b^- signal events while the EVTGEN program [10] is used to simulate

Ξ_b^- decays. The Ξ_b^- mass and lifetime are set to be 5.840 GeV and 1.33 ps, respectively, as their default values in these programs. The generated events are subjected to the same reconstruction and selection programs as the data after passing through the D0 detector simulation based on the GEANT package [11]. MC events are reweighted using the weights determined by matching transverse momentum (p_T) distributions of J/ψ , proton and pion from the $\Lambda_b \rightarrow J/\psi \Lambda \rightarrow J/\psi p \pi^-$ decays in MC calculations to those observed in the data.

$J/\psi \rightarrow \mu^+ \mu^-$ decays are reconstructed from two oppositely charged muons that have a common vertex. Muons are identified by matching tracks reconstructed in the central tracking system with either track segments in the muon spectrometer or calorimeter energies consistent with the muon trajectory. They are required to have $p_T > 1.5$ GeV and at least one of them must be reconstructed in each of the three muon drift tube layers. The dimuon invariant mass $M(\mu^+ \mu^-)$ is required to be in the range 2.5–3.6 GeV. In addition, events must have at least one reconstructed primary vertex of the $p\bar{p}$ interaction. If two or more vertices are reconstructed, the one closest to the reconstructed Ξ_b^- vertex (see below) is used. Events containing a J/ψ candidate are reprocessed with a version of the track reconstruction algorithm that improves the efficiency for tracks with low p_T and high impact parameters. Consequently, the efficiencies for K_S^0 , Λ , and Ξ^- reconstruction are significantly increased. Figure 2(a) shows the




FIG. 1 (color online). Schematic of the $\Xi_b^- \rightarrow J/\psi \Xi^- \rightarrow J/\psi \Lambda \pi^- \rightarrow (\mu^+ \mu^-)(p \pi^-) \pi^-$ decay topology. The Λ and Ξ^- baryons have decay lengths of the order of cm; the Ξ_b^- has an expected decay length of the order of mm.




FIG. 2. Invariant mass distributions of the $\Lambda\pi$ pair before the Ξ_b^- reconstruction for (a) the right-sign $\Lambda\pi^-$ combinations before and after reprocessing and (b) the right-sign $\Lambda\pi^-$ and the wrong-sign $\Lambda\pi^+$ combinations after reprocessing. The reprocessing significantly increases the Ξ^- yield. Fits to the post-re-processing distributions of the right-sign combination with a Gaussian signal and a first-order polynomial background yield 603 ± 34 Ξ^- 's and 548 ± 31 Ξ^+ 's.

invariant mass distributions of the reconstructed Ξ^- candidates (see below) before and after the reprocessing. The reprocessing increases the Ξ^- yield by approximately a factor of 5.5. For further analysis, $J/\psi \rightarrow \mu^+\mu^-$ candidates are required to have mass $2.80 < M(\mu^+\mu^-) < 3.35$ GeV and $p_T > 5$ GeV. The mass windows here and below are chosen to be approximately $\pm 5\sigma$ and the p_T requirement ensures that the selected J/ψ candidates are above the sharp turn-on of the detector and trigger acceptances.

$\Lambda \rightarrow p\pi^-$ candidates are formed from two oppositely charged tracks that originate from a common vertex. The track with the higher p_T is assumed to be the proton. MC studies show that this assignment gives nearly 100% correct combination. The invariant mass of the $p\pi^-$ pair must have a mass between 1.105 and 1.125 GeV. The two tracks are required to have a total of no more than two hits in the tracking detector before the reconstructed $p\pi^-$ vertex. Furthermore, the impact parameter significance (the impact parameter with respect to the event vertex divided by its uncertainty) must exceed three for both tracks and exceed four for at least one of them. These selection cuts are the same as those in Ref. [8].

The Λ candidates are then combined with negatively charged tracks (assumed to be pions) to form $\Xi^- \rightarrow \Lambda\pi^-$ decay candidates. The pion must have an impact parameter significance greater than three. The Λ and the pion are required to have a common vertex. For both Λ and Ξ^- candidates, the distance between the event vertex and its decay vertex is required to exceed 4 times its uncertainty. Moreover, the uncertainty of the distance between the production vertex and its decay vertex (decay length) in the transverse plane (the plane perpendicular to the beam direction) must be less than 0.5 cm. These two requirements reduce combinatoric and track mismeasurement backgrounds.

The two pions from $\Xi^- \rightarrow \Lambda\pi^- \rightarrow (p\pi^-)\pi^-$ decays (right sign) have the same charge. Consequently, the combination $\Lambda\pi^+$ (wrong sign) events form an ideal control sample for background studies. Figure 2(b) compares mass distributions of the right-sign $\Lambda\pi^-$ and the wrong-sign $\Lambda\pi^+$ combinations. The Ξ^- mass peak is evident in the distribution of the right-sign events. A $\Lambda\pi^-$ pair is considered to be a Ξ^- candidate if its mass is within the range $1.305 < M(\Lambda\pi^-) < 1.340$ GeV.

$\Xi_b^- \rightarrow J/\psi\Xi^-$ decay candidates are formed from J/ψ and Ξ^- pairs that originate from a common vertex and have an opening angle in the transverse plane less than $\pi/2$ rad. The uncertainty of the proper decay length of the $J/\psi\Xi^-$ vertex must be less than 0.05 cm in the transverse plane. A total of 2308 events remains after this preselection. The wrong-sign events are subjected to the same preselection as the right-sign events. A total of 1124 wrong-sign events is selected as the control sample.

Several distinctive features of the $\Xi_b^- \rightarrow J/\psi\Xi^- \rightarrow J/\psi\Lambda\pi^- \rightarrow (\mu^+\mu^-)(p\pi^-)\pi^-$ decay are utilized to further suppress backgrounds. The wrong-sign background events from the data and MC signal Ξ_b^- events are used for studying additional event selection criteria. Protons and pions from the Ξ^- decays of the Ξ_b^- events are expected to have higher momenta than those from most of the background processes. Therefore, protons are required to have $p_T > 0.7$ GeV. Similarly, minimum p_T requirements of 0.3 and 0.2 GeV are imposed on pions from Λ and Ξ^- decays, respectively. These requirements remove 91.6% of the wrong-sign background events while keeping 68.7% of the MC Ξ_b^- signal events. Backgrounds from combinatorics and other b hadrons are reduced by using topological decay information. Contamination from decays such as $B^- \rightarrow J/\psi K^{*-} \rightarrow J/\psi K_S^0 \pi^-$ and $B^0 \rightarrow J/\psi K^{*-} \pi^+ \rightarrow J/\psi (K_S^0 \pi^-) \pi^+$ are suppressed by requiring the Ξ^- candidates to have decay lengths greater than 0.5 cm and $\cos(\theta_{\text{col}}) > 0.99$, as the Ξ^- baryons in MC calculations have an average decay length of 4.8 cm. Here θ_{col} is the angle between the Ξ^- direction and the direction from the Ξ^- production vertex to its decay vertex in the transverse plane. These two requirements on the Ξ^- reduce the background by an additional 56.4%, while removing only 1.7% of the MC signal events. The contribution from the Ω_b^- baryon is estimated to be negligible. Finally, Ξ_b^- baryons are expected to have a sizable lifetime. To reduce prompt backgrounds, the transverse proper decay length significance of the Ξ_b^- candidates is required to be greater than two. This final criterion retains 83.1% of the MC signal events but only 43.9% of the remaining background events.

In the data, 51 events with the Ξ_b^- candidate mass between 5.2 and 7.0 GeV pass all selection criteria. The mass range is chosen to be wide enough to encompass masses of all known b hadrons as well as the predicted mass of the Ξ_b^- baryon. The candidate mass, $M(\Xi_b^-)$,

is calculated as $M(\Xi_b^-) = M(J/\psi \Xi^-) - M(\mu^+ \mu^-) - M(\Lambda \pi^-) + M_{\text{PDG}}(J/\psi) + M_{\text{PDG}}(\Xi^-)$ to improve the resolution. Here $M(J/\psi \Xi^-)$, $M(\mu^+ \mu^-)$, and $M(\Lambda \pi^-)$ are the reconstructed masses while $M_{\text{PDG}}(J/\psi)$ and $M_{\text{PDG}}(\Xi^-)$ are taken from Ref. [1]. The distribution of $M(\Xi_b^-)$ is shown in Fig. 3(a). A mass peak near 5.8 GeV is apparent. A number of cross checks are performed to ensure the observed peak is not due to artifacts of the analysis: (i) The $J/\psi \Lambda \pi^+$ mass distribution of the wrong-sign events, shown in Fig. 3(b), is consistent with a flat background. (ii) The event selection is applied to the sideband events of the Ξ^- mass peak, requiring $1.28 < M(\Lambda \pi^-) < 1.36$ GeV but excluding the Ξ^- mass window. Similarly, the selection is applied to the J/ψ sideband events with $2.5 < M(\mu^+ \mu^-) < 2.7$ GeV. The high-mass sideband is not considered due to potential contamination from ψ' events. As shown in Figs. 3(c) and 3(d), no evidence of a mass peak is present for either $(\mu^+ \mu^-)(p \pi^-) \pi^-$ distribution. (iii) The possibility of a fake signal due to the residual b hadron background is investigated by applying the final Ξ_b^- selection to high statistics MC samples of $B^- \rightarrow J/\psi K^{*-} \rightarrow J/\psi K_S^0 \pi^-$, $B^0 \rightarrow J/\psi K_S^0$, and $\Lambda_b \rightarrow J/\psi \Lambda$. No indication of a mass peak is observed in the reconstructed $J/\psi \Xi^-$ mass distributions. (iv) The mass distributions of J/ψ , Ξ^- , and Λ are investigated by relaxing the mass requirements on these particles one at a time for events both in the Ξ_b^- signal region and the sidebands. The numbers of these particles determined by fitting their respective mass distribution are fully consistent with the quoted numbers of signal events plus background contributions. (v) The robustness of the observed mass peak is tested by varying selection criteria




FIG. 3 (color online). (a) The $M(\Xi_b^-)$ distribution of the Ξ_b^- candidates after all selection criteria. The dotted curve is an unbinned likelihood fit to the model of a constant background plus a Gaussian signal. The $(\mu^+ \mu^-) \Lambda \pi$ mass distributions for (b) the wrong-sign background, (c) the Ξ^- sideband, and (d) the J/ψ sideband events. The mass $M(J/\psi \Lambda \pi) - M(\mu^+ \mu^-) + M_{\text{PDG}}(J/\psi)$ is plotted for (b) and (c) while the mass $M(\mu^+ \mu^- \Xi^-) - M(\Lambda \pi^-) + M_{\text{PDG}}(\Xi^-)$ is plotted for (d).

within reasonable ranges. All studies confirm the existence of the peak at the same mass.

Interpreting the peak as Ξ_b^- production, candidate masses are fitted with the hypothesis of a signal plus background model using an unbinned likelihood method. The signal and background shapes are assumed to be Gaussian and flat, respectively. The fit results in a Ξ_b^- mass of 5.774 ± 0.011 GeV with a width of 0.037 ± 0.008 GeV and a yield of 15.2 ± 4.4 events. Unless specified, all uncertainties are statistical. Following the same procedure, a fit to the MC Ξ_b^- events yields a mass of 5.839 ± 0.003 GeV, in good agreement with the 5.840 GeV input mass. The fitted width of the MC mass distribution is 0.035 ± 0.002 GeV, consistent with the 0.037 GeV obtained from the data. Since the intrinsic decay width of the Ξ_b^- baryon in the MC calculations is negligible, the width of the mass distribution is thus dominated by the detector resolution. To assess the significance of the signal, the likelihood, \mathcal{L}_{s+b} , of the signal plus background fit above is first determined. The fit is then repeated using the background-only model, and a new likelihood \mathcal{L}_b is found. The logarithmic likelihood ratio $\sqrt{2 \ln(\mathcal{L}_{s+b}/\mathcal{L}_b)}$ indicates a statistical significance of 5.5σ , corresponding to a probability of 3.3×10^{-8} from background fluctuation for observing a signal that is equal to or more significant than what is seen in the data. Including systematic effects from the mass range, signal and background models, and the track momentum scale results in a minimum significance of 5.3σ and a Ξ_b^- yield of $15.2 \pm 4.4(\text{stat})_{-0.4}^{+1.9}(\text{syst})$. The significance can also be estimated from the numbers of candidate events and estimated background events. In the mass region of 2.5 times the fitted width centered on the fitted mass, 19 candidate events (8 $J/\psi \Xi^-$ and 11 $J/\psi \Xi^+$) are observed while $14.8 \pm 4.3(\text{stat})_{-0.4}^{+1.9}(\text{syst})$ signal and $3.6 \pm 0.6(\text{stat})_{-1.9}^{+0.4}(\text{syst})$ background events are estimated from the fit. The probability of backgrounds fluctuating to 19 or more events is 2.2×10^{-7} , equivalent to a Gaussian significance of 5.2σ .

Figure 4 shows distributions of the proper decay length for the 19 candidate events, the Ξ_b^- sideband events, and the MC Ξ_b^- signal events plus estimated background events. The distribution of the candidate events agrees well with that expected from the Ξ_b^- signal while the sideband events have a lower mean proper decay length. Because of the use of lifetime information in the event selection, a Ξ_b^- lifetime measurement is not made in this Letter.

Potential systematic biases on the measured Ξ_b^- mass are studied for the event selection, signal and background models, and the track momentum scale. Varying cut values and using a multivariate technique of different variables for event selection leads to a maximum change of 0.020 GeV in the Ξ_b^- mass. Subtracting an estimated statistical contribution to the change, a conservative ± 0.015 GeV systematic uncertainty is assigned due to the event selection.




FIG. 4 (color online). The distribution of the proper decay length in the transverse plane of the 19 candidate events in the $\pm 2.5\sigma$ signal mass window along with that of the events in the sidebands, defined to be 5σ away from the fitted mass. Also shown is the expected distribution from 14.8 MC Ξ_b^- signal events plus 3.6 background events. The distribution of the sideband events is scaled to the number of events in the signal mass window. Kolmogorov-Smirnov tests indicate that the distribution of the signal events is favored over that of the sideband events with respect to the MC expectation by a ratio of five to one.

Using double Gaussians for the signal model, a first-order polynomial for the background model, or fixing the mass resolution to that obtained from the MC Ξ_b^- events all lead to negligible changes in the mass. The mass, calculated using the world average values [1] of intermediate particle masses above, is found to have a weak dependence on the track momentum scale. This has been verified using the $\Lambda_b \rightarrow J/\psi\Lambda$ and $B^0 \rightarrow J/\psi K_S^0$ events observed in the data. A systematic uncertainty of ± 0.002 GeV is assigned, corresponding to the mass difference between our measurement and the world average [1] for the Λ_b and B^0 hadrons. Adding in quadrature, a total systematic uncertainty of ± 0.015 GeV is obtained to yield the measured Ξ_b^- mass: $5.774 \pm 0.011(\text{stat}) \pm 0.015(\text{syst})$ GeV.

The Ξ_b^- $\sigma \times \mathcal{B}$ relative to that of the Λ_b baryon is calculated using

$$\frac{\sigma(\Xi_b^-) \times \mathcal{B}(\Xi_b^- \rightarrow J/\psi \Xi^-)}{\sigma(\Lambda_b) \times \mathcal{B}(\Lambda_b \rightarrow J/\psi \Lambda)} = \frac{\epsilon(\Lambda_b \rightarrow J/\psi \Lambda)}{\epsilon(\Xi_b^- \rightarrow J/\psi \Xi^-)} \frac{N_{\Xi_b^-}}{N_{\Lambda_b}},$$

where $N_{\Xi_b^-}$ and N_{Λ_b} are the numbers of Ξ_b^- and Λ_b events reconstructed in data. Analyzing the same data and using the similar event selection criteria and fitting procedure as the Ξ_b^- analysis, a yield of $240 \pm 30(\text{stat}) \pm 12(\text{syst})$ Λ_b baryons is determined. The efficiencies to reconstruct the decays, $\epsilon(\Xi_b^-)$ and $\epsilon(\Lambda_b)$, are determined by MC simulation, and the efficiency ratio, $\epsilon(\Lambda_b)/\epsilon(\Xi_b^-)$, is found to be 4.4 ± 1.3 . The uncertainty on $\epsilon(\Lambda_b)/\epsilon(\Xi_b^-)$ arises from MC modeling (27%), MC statistics (10%), the reconstruction of the additional pion in the Ξ_b^- decay (7%), and the

Ξ_b^- mass difference between data and MC calculations (5%). The largest component, MC modeling uncertainty, is due to the difference in the efficiency ratio with and without MC reweighting. The efficiency ratio is found to be insensitive to changes in Λ_b and Ξ_b^- production models. Many other systematic uncertainties on the efficiencies themselves tend to cancel in the ratio of the efficiencies. We find a relative production ratio of $0.28 \pm 0.09(\text{stat})^{+0.09}_{-0.08}(\text{syst})$.

In summary, in 1.3 fb^{-1} of data collected by the D0 experiment in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV at the Fermilab Tevatron collider, we have made the first direct observation of the strange b baryon Ξ_b^- with a statistical significance of 5.5σ . We observe the decay mode $\Xi_b^- \rightarrow J/\psi \Xi^-$ with $J/\psi \rightarrow \mu^+ \mu^-$, $\Xi^- \rightarrow \Lambda \pi^- \rightarrow p \pi^- \pi^-$. We measure the Ξ_b^- mass to be $5.774 \pm 0.011(\text{stat}) \pm 0.015(\text{syst})$ GeV and determine its $\sigma \times \mathcal{B}$ relative to that of the Λ_b to be $0.28 \pm 0.09(\text{stat})^{+0.09}_{-0.08}(\text{syst})$.

We thank the staffs at Fermilab and collaborating institutions, and acknowledge support from the DOE and NSF (USA); CEA and CNRS/IN2P3 (France); FASI, Rosatom, and RFBR (Russia); CAPES, CNPq, FAPERJ, FAPESP, and FUNDUNESP (Brazil); DAE and DST (India); Colciencias (Colombia); CONACyT (Mexico); KRF and KOSEF (Korea); CONICET and UBACyT (Argentina); FOM (The Netherlands); Science and Technology Facilities Council (United Kingdom); MSMT and GACR (Czech Republic); CRC Program, CFI, NSERC, and WestGrid Project (Canada); BMBF and DFG (Germany); SFI (Ireland); The Swedish Research Council (Sweden); CAS and CNSF (China); Alexander von Humboldt Foundation; and the Marie Curie Program.

*Visiting scientist from Augustana College, Sioux Falls, SD, USA.

†Visiting scientist from The University of Liverpool, Liverpool, United Kingdom.

‡Visiting scientist from ICN-UNAM, Mexico City, Mexico.

§Visiting scientist from Helsinki Institute of Physics, Helsinki, Finland.

||Visiting scientist from Universität Zürich, Zürich, Switzerland.

- [1] W.-M. Yao *et al.*, J. Phys. G **33**, 1 (2006).
- [2] N. Isgur and M. B. Wise, Phys. Rev. Lett. **66**, 1130 (1991).
- [3] G. T. Bodwin, E. Braaten, G. P. Lepage, Phys. Rev. D **51**, 1125 (1995); **55**, E5853 (1997).
- [4] E. Jenkins, Phys. Rev. D **55**, R10 (1997); **54**, 4515 (1996); N. Mathur, R. Lewis, and R. M. Woloshyn, Phys. Rev. D **66**, 014502 (2002).
- [5] J. Abdallah *et al.* (DELPHI Collaboration), Eur. Phys. J. **C44**, 299 (2005); D. Buskulic *et al.* (ALEPH Collaboration), Phys. Lett. B **384**, 449 (1996).
- [6] E. Barberio *et al.* (Heavy Flavor Averaging Group Collaboration), arXiv:0704.3575.
- [7] V. M. Abazov *et al.* (D0 Collaboration), Nucl. Instrum.

- Methods Phys. Res., Sect. A **565**, 463 (2006).
- [8] V.M. Abazov *et al.* (D0 Collaboration), arXiv:0704.3909.
- [9] T. Sjöstrand *et al.*, Comput. Phys. Commun. **135**, 238 (2001).
- [10] D.J. Lange, Nucl. Instrum. Methods Phys. Res., Sect. A **462**, 152 (2001).
- [11] R. Brun and F. Carminati, CERN Program Library Writeup W5013, 1993 (unpublished).