

2016

Double peak searches for scalar and pseudoscalar resonances at the LHC

Marcela Carena
University of Chicago

Peisi Huang
University of Chicago

Ahmed Ismail
The University of Illinois at Chicago

Ian Low
Northwestern University

Nausheen R. Shah
Wayne State University

See next page for additional authors

Follow this and additional works at: <http://digitalcommons.unl.edu/physicsfacpub>

Carena, Marcela; Huang, Peisi; Ismail, Ahmed; Low, Ian; Shah, Nausheen R.; and Wagner, Carlos E. M., "Double peak searches for scalar and pseudoscalar resonances at the LHC" (2016). *Faculty Publications, Department of Physics and Astronomy*. 220.
<http://digitalcommons.unl.edu/physicsfacpub/220>

This Article is brought to you for free and open access by the Research Papers in Physics and Astronomy at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Publications, Department of Physics and Astronomy by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors

Marcela Carena, Peisi Huang, Ahmed Ismail, Ian Low, Nausheen R. Shah, and Carlos E. M. Wagner

Double peak searches for scalar and pseudoscalar resonances at the LHCMarcela Carena,^{1,2,3} Peisi Huang,^{2,4} Ahmed Ismail,^{4,5,6} Ian Low,^{4,7} Nausheen R. Shah,⁸ and Carlos E. M. Wagner^{2,3,4}¹*Fermi National Accelerator Laboratory, P.O. Box 500, Batavia, Illinois 60510, USA*²*Enrico Fermi Institute, University of Chicago, Chicago, Illinois 60637, USA*³*Kavli Institute for Cosmological Physics, University of Chicago, Chicago, Illinois 60637, USA*⁴*High Energy Physics Division, Argonne National Laboratory, Argonne, Illinois 60439, USA*⁵*Department of Physics, University of Illinois, Chicago, Illinois 60607, USA*⁶*Kavli Institute for Theoretical Physics, University of California, Santa Barbara, California 93106, USA*⁷*Department of Physics and Astronomy, Northwestern University, Evanston, Illinois 60208, USA*⁸*Department of Physics and Astronomy, Wayne State University, Detroit, Michigan 48201, USA*

(Received 4 October 2016; published 1 December 2016)

Many new physics models contain a neutral scalar resonance that can be predominantly produced via gluon fusion through loops. In such a case, there could be important effects of additional particles, that in turn may hadronize before decaying and form bound states. This interesting possibility may lead to novel signatures with double peaks that can be searched for at the LHC. We study the phenomenology of double peak searches in diboson final states from loop-induced production and decay of a new neutral spin-0 resonance at the LHC. The loop-induced couplings should be mediated by particles carrying color and electroweak charge that after forming bound states will induce a second peak in the diboson invariant mass spectrum near twice their mass. A second peak could be present via loop-induced couplings into gg (dijet), $\gamma\gamma$ and $Z\gamma$ final states as well as in the WW and ZZ channels for the case of a pseudoscalar resonance or for scalars with suppressed tree-level coupling to gauge bosons.

DOI: 10.1103/PhysRevD.94.115001

I. INTRODUCTION

Diboson resonance searches have enjoyed renewed attention in the first years of the LHC. At run 1, the diphoton channel played a central role in cementing the position of the newly discovered Higgs boson [1,2]. Furthermore, ongoing searches for TeV-scale resonances decaying into W and Z bosons are fundamental to understanding the role of the Higgs in unitarizing vector boson scattering. The LHC experiments observed small excesses in diboson searches at an invariant mass of about 1.8–2 TeV [3,4] which became less significant with the inclusion of run 2 data [5,6]. A heavy diboson resonance can have many different origins, and observing the relative signal strengths in different diboson channels can help discriminate between candidate models of physics at the TeV scale.

For instance, composite Higgs models often predict resonances in diboson channels involving the W , Z gauge bosons and the SM Higgs h . By contrast, if a resonance is first observed in the diphoton channel, the natural explanation is associated with a new neutral spin-0 resonance, which is resonantly produced through gluon fusion, and decays into two photons (see, for example, Refs. [7–11] and references therein for recent studies of scalars decaying to photons). A sizable diboson rate demands a large decay branching ratio of the new scalar and pseudoscalar to a pair of bosons, and therefore, its tree-level couplings to Standard Model (SM) fermions must be somewhat suppressed. In the case of a resonance appearing first in $\gamma\gamma$, new particles that carry color and electric charge are

expected to be present, in order to mediate the loop-induced couplings of the new heavy scalar and pseudoscalar to SM gauge bosons.

In this work, we concentrate on the possible appearance of two peaks, associated with a new neutral spin-0 resonance with negligible couplings to SM fermions. We shall assume the presence of loop-induced couplings of the new spin-0 resonance to gluons and electroweak gauge bosons, mediated by new colored and charged particles. Gluon fusion production yields a resonant peak at the spin-0 resonance mass in all possible diboson channels. The relative strength of the resonant peak in different diboson channels depends on the quantum numbers of the loop particles. We demonstrate the potential existence of two peaks, rather than one, in the diboson invariant mass spectrum, focusing on the $\gamma\gamma$ signature. The double peak signature would indicate that the loop particles that contribute to the production and decay processes hadronize before decaying, forming new QCD bound states slightly below the pair production threshold.¹ Such a bound state would then give rise to a second peak in the diboson invariant mass spectrum, in addition to the resonance due to the production and decay of the new scalar and

¹If the new loop particles decay before hadronizing, there could still be a “shoulder” in the diboson invariant mass spectrum due to the amplitude developing an imaginary part at the kinematic threshold, such as in the case of double Higgs production [12]. Such a shoulder, however, may be too small to be observed for reasonable choices of parameters [13].

pseudoscalar. Moreover, the loop particles are expected to have a mass that is larger than half the mass of the neutral spin-0 resonance, in order to avoid tree-level decays of the neutral scalar and pseudoscalar into the loop particles. In this case, the second peak would show up at an invariant mass that is larger than the mass of the neutral spin-0 resonance.

The conditions for the presence of two peaks are therefore (1) loop-induced couplings of the neutral scalar and pseudoscalar with SM gauge bosons and (2) mediation of the loop-induced couplings by particles which form bound states. The first condition is satisfied for any neutral scalar and pseudoscalar coupling to massless gauge bosons. For massive gauge bosons such as W and Z bosons, the couplings to a pseudoscalar boson are also loop induced. To fulfill the second condition, the loop particles should have a decay width that is smaller than the QCD hadronization scale. Our considerations apply to any model containing a heavy neutral scalar and pseudoscalar coupling to gluons and electroweak gauge bosons via new colored and charged particles in the loop. Therefore, the results shown here may have implications for future searches of new physics in diboson channels at the LHC.

The remainder of this paper is organized as follows. In Sec. II, we concentrate on final states containing two photons, and calculate the diphoton production cross section associated with the bound state of the new colored particles. The production cross section of other gauge boson channels may be obtained from the diphoton one by simple group theoretical factors associated with the quantum numbers of the loop particles [9]. In Sec. III, we discuss the phenomenology of the new colored particles. We reserve Sec. IV for a discussion of our scenario and our conclusions.

II. THE BOUND STATE PEAK

We shall assume the existence of a neutral pseudoscalar A , which couples to gluons and electroweak gauge bosons via loops of new vectorlike quarks (VLQ) Ψ , and that the width of A is dominated by the resulting loop-induced decays. We also assume that the VLQs hadronize before they decay, so that a $J = 0$ quarkonium state is formed. Let us denote the relevant Yukawa coupling λ ,

$$\mathcal{L} \supset -i\lambda A \bar{\Psi} \gamma^5 \Psi. \quad (1)$$

Furthermore, we will assume that there are N degenerate copies of Ψ , each with electric charge Q transforming in a given color representation \mathcal{R} of dimension $D_{\mathcal{R}}$.

Then, assuming $m_{\Psi} > m_A/2$ so that mixing effects between A and the bound states can be neglected, the following property holds:

$$\Gamma(A \rightarrow gg) \propto \lambda^2 N^2 C_{\mathcal{R}}^2 \alpha_s^2, \quad (2)$$




FIG. 1. Examples of a bound state peak accompanying a 1 TeV pseudoscalar resonance in the diphoton invariant mass spectrum with 50 fb^{-1} of integrated luminosity at 13 TeV. An 800 GeV charge $5/3$ VLQ is assumed, leading to a bound state peak (blue) near $m_{\gamma\gamma} \sim 1.6$ TeV. The signal strength of the pseudoscalar resonance is fixed by its Yukawa coupling to the VLQ, which is 0.3 (0.1) in the magenta solid (dashed) curve. We assume a 1% resolution on each peak. The red curve represents an estimate of the diphoton background, scaled from LHC data [16].

where $C_{\mathcal{R}}$ is the quadratic Casimir of \mathcal{R} . Similarly,

$$\Gamma(A \rightarrow \gamma\gamma) \propto \lambda^2 N^2 D_{\mathcal{R}}^2 Q^4 \alpha^2. \quad (3)$$

The total width is naturally dominated by $\Gamma(A \rightarrow gg)$ and, therefore,

$$\Gamma_{\text{total}} \propto \lambda^2 N^2 C_{\mathcal{R}}^2 \alpha_s^2. \quad (4)$$

Under the above conditions,

$$\begin{aligned} \sigma(pp \rightarrow A \rightarrow \gamma\gamma) &\propto \Gamma(A \rightarrow gg) \frac{\Gamma(A \rightarrow \gamma\gamma)}{\Gamma_{\text{total}}} \\ &\propto \lambda^2 N^2 D_{\mathcal{R}}^2 Q^4 \alpha^2, \end{aligned} \quad (5)$$

and the diphoton invariant mass spectrum exhibits a resonant peak at $m_{\gamma\gamma} \sim m_A$.

On the other hand, the second peak is induced by the quarkonium bound state $\bar{\Psi}\Psi$ and the strength is proportional to [14]

$$\sigma(pp \rightarrow \bar{\Psi}\Psi \rightarrow \gamma\gamma) \propto N^2 C_{\mathcal{R}}^3 D_{\mathcal{R}} Q^4 \alpha_s^3 \alpha^2, \quad (6)$$

which gives rise to a peak in the diphoton spectrum at $m_{\gamma\gamma} \sim 2m_{\Psi} > m_A$. We see then that the strength of the peak at m_A relative to the one at $2m_{\Psi}$ is governed by λ and depends on the color representation of Ψ , but is independent of the electric charge or multiplicity of Ψ .²

²If the loop-induced couplings are generated by more than one colored particle, there may be multiple additional peaks and their strengths will depend on the particular lifetimes, multiplicities and quantum numbers of each of these particles.




FIG. 2. Top panel: Diphoton signals from $\bar{\Psi}\Psi$, QCD bound states of VLQs, at $\sqrt{s} = 13$ TeV as a function of the lowest bound state mass M for the representation $(3, 1, 5/3)$. The signal strength is computed from the Coulomb potential calculations in Ref. [14]. The lower curve of the orange band corresponds to choosing $N = 1$, while the top curve corresponds to $N = 2$. Bottom panel: Diphoton resonant cross section at 13 TeV from production through a loop of VLQ in the representation $(3, 1, 5/3)$ with mass 1 TeV, as a function of the resonance mass, for $\lambda = 0.3$ (magenta, solid line) and $\lambda = 1$ (red, dashed line). The gray (blue) curve in both panels is the current diphoton cross section limit from ATLAS (CMS).

As an illustration of the above relations, in Fig. 1 we present examples of two peaks in the diphoton invariant mass spectrum assuming a luminosity of 50 fb^{-1} , where the first peak is chosen to be at 1 TeV. We choose a single charge $5/3$ weak singlet VLQ with $m_\Psi = 800$ GeV and $\lambda \approx 0.1, 0.3$. The signal strengths of the peaks induced by the quarkonia bound states were computed using the Coulomb potential approximation [15]. For $\lambda = 0.3$, we expect to see the diphoton signal from the resonance first while for $\lambda = 0.1$, we expect the diphoton signal from the bound state to first reach a sizable significance. Eventually with more data, two peaks would be seen in both cases.

In Fig. 2 we show the diphoton cross section for the bound state and the resonance. In the top panel, we show the cross section associated with the formation of quarkonium bound states $\bar{\Psi}\Psi$ at $m_{\gamma\gamma} \sim 2m_\Psi$ for a weak singlet VLQ with charge $5/3$. The bottom curve of the band

TABLE I. Possible loop VLQs that can produce a 0.4 fb diphoton signal at 1 TeV, which is somewhat below the current cross section bound and consistent with results obtained for $\lambda = 0.3$ in Fig 1. The VLQ mass is assumed to be 800 GeV. In the second column, the vectorlike fermions are assumed to couple to a pseudoscalar resonance, while in the third column, the resonance is a scalar. The diphoton signal rate depends on the product of λN and hence the value of N can be scaled to integer values by choosing different values of λ .

$SU(3) \times SU(2) \times U(1)$	λN , pseudoscalar	λN , scalar
$(3, 1, 5/3)$	0.3	0.5
$(3, 1, 4/3)$	0.5	0.8
$(3, 2, 7/6)$	0.3	0.4
$(3, 2, 5/6)$	0.5	0.7
$(3, 1, 2/3)$	2.0	3.0
$(3, 2, 1/6)$	1.6	2.4

corresponds to $N = 1$. The top curve results from the choice $N = 2$, or alternatively $N = 1$ with a bound state wave function at the origin that is two times larger than the value obtained using the Coulomb potential approximation, as suggested by a recent lattice calculation [17]. We also show the present bounds on the diphoton cross section from ATLAS [18] and CMS [19], obtained at 13 TeV with an integrated luminosity of 15.4 fb^{-1} and 12.9 fb^{-1} , respectively. Bound states of VLQs Ψ with masses up to $m_\Psi \sim 700$ GeV are already excluded. In the bottom panel, we show the diphoton cross section from a resonance assuming the existence of a single charge $5/3$ weak singlet VLQ, with mass chosen to be 1 TeV and Yukawa couplings $\lambda = 0.3$ (solid, magenta line) and $\lambda = 1$ (dashed, red line). As can be seen from this panel, Yukawa coupling values between about 0.4 and 1 are currently probed by the LHC, with smaller Yukawa couplings generically probed for smaller values of the resonance mass.

Regarding the future LHC sensitivity, scaling the ATLAS diphoton background fit [16] to higher integrated luminosity, we estimate that the LHC with 300 fb^{-1} of data will be able to probe bound states from a single $(3, 1, 5/3)$ VLQ with a mass lower than roughly 1030 GeV, which is very close to the mass assumed in our benchmark scenario. This implies that for this value of the VLQ mass the LHC is already probing Yukawa couplings in the $\lambda \approx 0.4-1$ range. Values of the Yukawa couplings a factor of about 2.5 smaller than the currently probed ones will be probed at these higher luminosities. These are the minimal values that could be probed in the absence of a LHC diphoton peak (larger values of λ would still be allowed for larger values of the VLQ mass). Let us stress that although we have taken the example of a single $(3, 1, 5/3)$ VLQ, it is straightforward to use similar methods to compute the bounds on the mass and couplings for other values of the VLQ charge, representations and multiplicities, as the ones shown in Table I.

III. PHENOMENOLOGICAL CONSTRAINTS

The prediction of two peaks in the diphoton invariant mass spectrum applies to a wide class of models. Using a sample 1 TeV spin-0 resonance as a benchmark, in Table I we summarize possible new VLQs that could explain a potential signal, assuming a production cross section 0.4 fb, which is somewhat below the current bound on the cross section for this resonance mass and consistent with results obtained for $\lambda = 0.3$ in Fig 1. We show results for both pseudoscalar and scalar resonances.³ Observe that λN is approximately the same for certain $SU(2)$ doublet and singlet cases, since the main contribution to the diphoton rate comes from the particle in the multiplet with the largest electric charge.

Since quarkonia states would only develop if the loop particles hadronize before decaying, it is relevant to discuss the possible decays of these particles. Among the VLQ quantum numbers listed in Table I, all but the $(3, 1, 5/3)$ and $(3, 1, 4/3)$ cases may decay into Standard Model particles, via Yukawa couplings that mix these exotic states with SM states. The values of these Yukawa couplings should be chosen so that the new colored particles hadronize before decaying, something that naturally occurs for Yukawa couplings smaller than about a few times 10^{-2} . Limits on the mass of the VLQs depend on the decay branching fraction into various final states⁴ and are typically $\gtrsim 800$ GeV for decays to SM quarks and vector bosons [23–25]. For multiple copies of the VLQs, as may be required for some of the representations in Table I, these bounds will be even stronger.

The colored particles also hadronize before decaying when they lie in compressed spectra, e.g. as motivated by coannihilation scenarios where the colored particles are expected to be only slightly heavier than some new dark matter state [26]. For instance, one can add a new scalar $\phi(1, 2, -1/2)$ and impose a new Z_2 parity at the TeV scale under which the SM is even and all new particles are odd. Such a new parity has several advantages and distinct collider phenomenology [27,28], similar to cases in which possible modifications of the SM Higgs diphoton rate were studied [29,30]. Since ϕ has the same quantum numbers as the Higgs field, again a Yukawa coupling among ϕ , Ψ and a SM quark is possible for all but the $(3, 1, 5/3)$ and $(3, 1, 4/3)$ states. This new Yukawa coupling leads to the decay

$$\Psi \rightarrow q + \phi^0, \quad (7)$$

where ϕ^0 is expected to be the lightest odd particle and therefore can be a dark matter candidate. Then, for

³For CP -even resonances, loop scalars may be allowed in place of the VLQs, but for resonance couplings to a pair of scalars of about the scalar mass, the resonant cross section is significantly lower than in the $\lambda = 1$ VLQ case [20–22].

⁴For a scan over decay branching fractions, see “additional materials” on the website in Ref. [23].

sufficiently heavy ϕ^0 , the decay products will have soft kinematic spectra. Consequently, Ψ pair production above threshold will not be subject to the existing strong bounds on particles decaying to SM quarks and vector gauge bosons. In the limit that ϕ and Ψ are very close in mass, coannihilation with Ψ may help to set the proper relic density of ϕ and provide a natural motivation for the long Ψ lifetime.

On the other hand, for the exotic singlet VLQs in SM representations $(3, 1, 5/3)$ and $(3, 1, 4/3)$, there is no renormalizable, gauge invariant operator to mediate their decays into ϕ (or SM particles). Additional particle content is thus required. As an example, when the loop particle Ψ is in the $(3, 1, 5/3)$ representation, one could consider the existence of an additional isospin doublet VLQ Ψ' in the $(3, 2, 7/6)$ representation. All new particles, Ψ , Ψ' and ϕ , are odd under the new Z_2 parity. If $m_{\Psi'} > m_{\Psi} > m_{\phi}$ and the Yukawas of the Ψ and Ψ' are similar, the extra VLQ does not contribute appreciably to the diphoton spectrum. Then, with the Lagrangian

$$\mathcal{L} \supset -y_{\Psi} H \bar{\Psi} \Psi' - \lambda_{\phi} \phi \bar{l}_R \Psi', \quad (8)$$

Ψ decays proceed through off-shell Ψ' ,

$$\Psi \rightarrow t + V/h + \phi^0. \quad (9)$$

Once again, the decay products of Ψ will be soft if ϕ^0 is sufficiently heavy, implying weaker collider bounds and therefore allowing lower VLQ masses. Similar considerations apply to the case in which the loop particle is in the $(3, 1, 4/3)$ representation, with the only difference being that Ψ' should be in the $(3, 2, 5/6)$ representation. The lifetime of the exotic singlet VLQs is naturally large, ensuring that these VLQs hadronize before decaying, leading to the formation of bound states.

The addition of the new VLQs in Table I to the SM modifies the renormalization group evolution of the gauge couplings and one can inquire about the perturbative consistency of the theory. For $\lambda \sim 1$, among the possibilities in Table I, all but the exotic singlet VLQs have been studied in Ref. [31] and the perturbativity of the theory is maintained up to very high energy scales. We checked that this remains the case for the exotic singlet VLQs. However, if extra fermions were included to induce the VLQs' decays, as in Eq. (8), the hypercharge gauge coupling would become nonperturbative at scales of the order of 10^{11} GeV, demanding the appearance of new physics below that energy scale.

IV. CONCLUSIONS

In this work we consider a wide class of models containing a neutral scalar and pseudoscalar coupling to gluons and electroweak gauge bosons via loop-induced couplings of new particles carrying color and electroweak charges. We showed that, if the new loop particles hadronize before decaying, there will be a second peak

in the diboson invariant mass spectrum at around twice the mass of the new loop particles. This additional peak is generically at a larger mass than the peak due to the resonant production of the neutral scalar and pseudoscalar. This new peak, on the other hand, is due to the new colored particles forming QCD bound states, which subsequently decay back into gauge boson pairs.

The existence of a second peak is quite generic for a heavy diboson resonance arising from a new neutral spin-0 resonance produced in the gluon fusion channel. The only requirement is that the new colored particle has a decay width that is smaller than the QCD hadronization scale. The signal from resonant scalar and pseudoscalar production and the bound state peak are not restricted to the $\gamma\gamma$ final state that we have considered in this work, but are generically expected to appear in the gg , $Z\gamma$ and ZZ channels as well. If the loop particle is charged under $SU(2)$, a WW signature is also possible. While searches involving photons and leptonically decaying Z bosons tend to be the cleanest experimentally, ultimately the relative strengths of these channels depend on the particular gauge quantum numbers of the loop particle. These quantum numbers affect the decays of both the bound state and the resonance simultaneously, and consequently an interesting test of our hypothesis is that two diboson peaks are seen with proportional strengths across different channels.

In conclusion, any model that attempts to predict a diboson excess by means of loops of colored and electro-weak charged particles should be subject to not only the

constraints imposed by direct searches for these loop particles, but also to constraints coming from searches for new diboson peaks associated with the formation of bound states, if the loop particles hadronize before they decay. Indeed, the positions of the additional peaks could provide a better determination of the mass of the new particles than the one provided by direct searches.

ACKNOWLEDGMENTS

Fermilab is operated by Fermi Research Alliance, LLC under Contract No. DE-AC02-07CH11359 with the U.S. Department of Energy. Work at University of Chicago is supported in part by U.S. Department of Energy Grant No. DE-FG02-13ER41958. Work at Argonne National Laboratory (ANL) is supported in part by the U.S. Department of Energy under Contract No. DE-AC02-06CH11357. P.H. is partially supported by U.S. Department of Energy Grant No. DE-FG02-04ER41286. A. I. thanks the Kavli Institute for Theoretical Physics for its hospitality, and is supported in part by the U.S. Department of Energy under Grant No. DE-FG02-12ER41811 and the National Science Foundation under Grant No. NSF PHY11-25915. I.L. is supported in part by the U.S. Department of Energy under Contract No. DE-SC0010143. N.R.S. is supported in part by the Wayne State University Start-up package. C.W. would like to thank G. Bodwin, H. S. Chung, A. Joglekar and A. Katz for helpful discussions on this subject.

-
- [1] ATLAS Collaboration, Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC, *Phys. Lett. B* **716**, 1 (2012).
 - [2] CMS Collaboration, Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC, *Phys. Lett. B* **716**, 30 (2012).
 - [3] ATLAS Collaboration, Search for high-mass diboson resonances with boson-tagged jets in proton-proton collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector, *J. High Energy Phys.* **12** (2015) 055.
 - [4] CMS Collaboration, Search for massive resonances in dijet systems containing jets tagged as W or Z boson decays in pp collisions at $\sqrt{s} = 8$ TeV, *J. High Energy Phys.* **08** (2014) 173.
 - [5] CMS Collaboration, CERN Technical Report No. CMS-PAS-EXO-15-002, 2015, <http://cds.cern.ch/record/2117062>.
 - [6] ATLAS Collaboration, CERN Technical Report No. ATLAS-CONF-2015-073, 2015, <http://cds.cern.ch/record/2114845>.
 - [7] R. Franceschini, G. F. Giudice, J. F. Kamenik, M. McCullough, A. Pomarol, R. Rattazzi, M. Redi, F. Riva, A. Strumia, and R. Torre, What is the gamma gamma resonance at 750 GeV?, *J. High Energy Phys.* **03** (2016) 144.
 - [8] J. Ellis, S. A. R. Ellis, J. Quevillon, V. Sanz, and T. You, On the interpretation of a possible ~ 750 GeV particle decaying into $\gamma\gamma$, *J. High Energy Phys.* **03** (2016) 176.
 - [9] I. Low and J. Lykken, Implications of gauge invariance on a heavy diphoton resonance, [arXiv:1512.09089](https://arxiv.org/abs/1512.09089).
 - [10] W. Altmannshofer, J. Galloway, S. Gori, A. L. Kagan, A. Martin, and J. Zupan, On the 750 GeV di-photon excess, *Phys. Rev. D* **93**, 095015 (2016).
 - [11] A. Strumia, Interpreting the 750 GeV digamma excess: A review, [arXiv:1605.09401](https://arxiv.org/abs/1605.09401).
 - [12] S. Dawson, A. Ismail, and I. Low, What's in the loop? The anatomy of double Higgs production, *Phys. Rev. D* **91**, 115008 (2015).
 - [13] S. Jain, F. Margaroli, S. Moretti, and L. Panizzi, The 750 GeV threshold to a new particle world, [arXiv:1605.08741](https://arxiv.org/abs/1605.08741).
 - [14] Y. Kats and M. J. Strassler, Probing colored particles with photons, leptons, and jets, *J. High Energy Phys.* **11** (2012) 097.
 - [15] Y. Kats and M. J. Strassler, Resonances from QCD bound states and the 750 GeV diphoton excess, *J. High Energy Phys.* **05** (2016) 092.

- [16] ATLAS Collaboration, CERN Technical Report No. ATLAS-CONF-2015-081, 2015, <http://cds.cern.ch/record/2114853>.
- [17] S. Kim, Nonrelativistic lattice study of stoponium, *Phys. Rev. D* **92**, 094505 (2015).
- [18] ATLAS Collaboration, CERN Technical Report No. ATLAS-CONF-2016-059, 2016, <http://cds.cern.ch/record/2206154>.
- [19] CMS Collaboration, CERN Technical Report No. CMS-PAS-EXO-16-027, 2016, <http://cds.cern.ch/record/2205245>.
- [20] D. Choudhury and K. Ghosh, The LHC diphoton excess at 750 GeV in the framework of the constrained minimal supersymmetric Standard Model, [arXiv:1605.00013](https://arxiv.org/abs/1605.00013).
- [21] A. Djouadi and A. Pilaftsis, The 750 GeV diphoton resonance in the MSSM, [arXiv:1605.01040](https://arxiv.org/abs/1605.01040).
- [22] G. T. Bodwin, H. S. Chung, and C. E. M. Wagner, Higgs-stoponium mixing near the stop-antistop threshold, [arXiv:1609.04831](https://arxiv.org/abs/1609.04831).
- [23] CMS Collaboration, CERN Technical Report No. CMS-PAS-B2G-12-017, 2014, https://twiki.cern.ch/twiki/bin/view/CMSPublic/PhysicsResultsB2G12017#Additional_Material.
- [24] CMS Collaboration, CERN Technical Report No. CMS-PAS-B2G-16-002, 2016, <http://cds.cern.ch/record/2141070>.
- [25] CMS Collaboration, CERN Technical Report No. CMS-PAS-B2G-15-006, 2016, <http://cds.cern.ch/record/2114805>.
- [26] K. Griest and D. Seckel, Three exceptions in the calculation of relic abundances, *Phys. Rev. D* **43**, 3191 (1991).
- [27] H.-C. Cheng and I. Low, TeV symmetry and the little hierarchy problem, *J. High Energy Phys.* **09** (2003) 051.
- [28] H.-C. Cheng, I. Low, and L.-T. Wang, Top partners in little Higgs theories with T-parity, *Phys. Rev. D* **74**, 055001 (2006).
- [29] A. Joglekar, P. Schwaller, and C. E. M. Wagner, Dark matter and enhanced Higgs to di-photon rate from vector-like leptons, *J. High Energy Phys.* **12** (2012) 064.
- [30] N. Arkani-Hamed, K. Blum, R. T. D'Agnolo, and J. Fan, 2:1 for naturalness at the LHC?, *J. High Energy Phys.* **01** (2013) 149.
- [31] K. J. Bae, M. Endo, K. Hamaguchi, and T. Moroi, Diphoton excess and running couplings, *Phys. Lett. B* **757**, 493 (2016).