

2012

# Interleukin-6 control of early Theiler's Murine Encephalomyelitis Virus replication in macrophages occurs in conjunction with STAT1 activation and nitric oxide production.

Tyler C. Moore

*University of Nebraska-Lincoln*, tylermooreunl@huskers.unl.edu

Katherine L. Bush

*University of Nebraska Medical Center*

Deborah Brown

*University of Nebraska - Lincoln*, dbrown7@unl.edu

Thomas M. Petro

*University of Nebraska Medical Center*, tpetro@unmc.edu

Follow this and additional works at: <http://digitalcommons.unl.edu/virologypub>



Part of the [Virology Commons](#)

---

Moore, Tyler C.; Bush, Katherine L.; Brown, Deborah; and Petro, Thomas M., "Interleukin-6 control of early Theiler's Murine Encephalomyelitis Virus replication in macrophages occurs in conjunction with STAT1 activation and nitric oxide production." (2012). *Virology Papers*. 228.

<http://digitalcommons.unl.edu/virologypub/228>

This Article is brought to you for free and open access by the Virology, Nebraska Center for at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Virology Papers by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

1 **Interleukin-6 control of early Theiler's Murine**  
2 **Encephalomyelitis Virus replication in**  
3 **macrophages occurs in conjunction with STAT1**  
4 **activation and nitric oxide production.**

5  
6  
7 **Tyler C. Moore<sup>a</sup>, Katherine L. Bush<sup>c</sup>, Liz Cody<sup>a,b</sup>, Deborah M. Brown<sup>a,b</sup>, and**  
8 **Thomas M. Petro<sup>b,c,#</sup>.**

9 <sup>a</sup>School of Biological Sciences and <sup>†</sup>Nebraska Center for Virology,  
10 University of Nebraska Lincoln

11 <sup>c</sup>Department of Oral Biology, University of Nebraska Medical Center

12 #Address correspondence to:

13 Thomas M. Petro, PhD

14 Dept. of Oral Biology

15 Univ. of Nebraska Med. Ctr.

16 40th and Holdrege St.

17 Lincoln, NE 68583-0740

18 phone: 402-472-1327

19 fax: 402-472-2551

20 email: [tpetro@unmc.edu](mailto:tpetro@unmc.edu)

21

22

23 **Short Running Title:**

24 Interleukin-6 and control of early TMEV infection

25

26 **Word Count:** 5,615

27 241 Words in Abstract

28

29

30

31

32

33

34

35

36

37

38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74

**Abbreviations:**

IFN=interferon

IRF=interferon response factor

TMEV=Theiler's murine encephalomyelitis virus

MAPK=mitogen-activated protein kinase

ERK= extracellular signal-regulated kinase

EAE = experimental autoimmune encephalomyelitis

75 **ABSTRACT**

76

77 During Theiler's virus (TMEV) infection of macrophages it is thought that high IL-6 levels  
78 contribute to demyelinating disease found in chronically infected SJL/J mice but absent  
79 in B10.S mice capable of clearing the infection. Therefore, IL-6 expression was  
80 measured in TMEV-susceptible SJL/J and TMEV-resistant B10.S macrophages during  
81 their infection with TMEV DA strain or responses to LPS or poly I:C. Unexpectedly, IL-6  
82 production was greater in B10.S macrophages than SJL/J macrophages during the first  
83 24 h after stimulation with TMEV, LPS or poly I:C. Further experiments showed that in  
84 B10.S, SJL/J, and RAW264.7 macrophage cells, IL-6 expression was dependent on  
85 extracellular signal-regulated kinase (ERK) mitogen-activated protein kinase (MAPK)  
86 and enhanced by exogenous IL-12. In SJL/J and RAW264.7 macrophages, exogenous  
87 IL-6 resulted in decreased TMEV replication, earlier activation of STAT1 and STAT3,  
88 production of nitric oxide, and earlier up-regulation of several anti-viral genes  
89 downstream of STAT1. However, neither inhibition of IL-6-induced nitric oxide nor  
90 knockdown of STAT1 diminished the early anti-viral effect of exogenous IL-6. In  
91 addition, neutralization of endogenous IL-6 from SJL/J macrophages with Fab  
92 antibodies did not exacerbate early TMEV infection. Therefore, endogenous IL-6  
93 expression after TMEV infection is dependent on ERK MAPK, enhanced by IL-12, but  
94 too slow to decrease viral replication during early infection. In contrast, exogenous IL-6  
95 enhances macrophage control of TMEV infection through preemptive anti-viral nitric  
96 oxide production and anti-viral STAT1 activation. These results indicate that immediate-  
97 early production of IL-6 could protect macrophages from TMEV infection.

98

99 **INTRODUCTION**

100 Interleukin (IL)-6 is a pleiotropic cytokine expressed by many cell types that is induced  
101 by microbes and other cytokines. IL-6 can play a beneficial role during immune  
102 response by contributing to neutrophil activity, nitric oxide production and development  
103 of the Th17 CD4 T cell subset, and a beneficial role in the brain (43) and neuronal  
104 health (9, 11). In contrast, persistent IL-6 induction during infection can contribute to  
105 detrimental effects on surrounding neurons or bone and poor control of cancer or  
106 autoimmune responses (24). During microbial induction of IL-6, activation of p38  
107 mitogen-activated protein kinase (MAPK) and extracellular signal-regulated kinase  
108 (ERK) MAPK signaling pathways contribute to IL-6 expression (21, 49) and stabilization  
109 of IL-6 mRNA (2) for protein translation.

110         Macrophages express IL-6 when they encounter microbes such as bacteria,  
111 fungi, or viruses (44). While the neutrophil- and Th17-promoting activity of IL-6 is critical  
112 in controlling certain bacterial or fungal infections, less is known about the role of IL-6 in  
113 viral infections. Recently, an antiviral role for IL-6 has been shown in mouse poxvirus  
114 infections, where early IL-6 production is required to control a potentially lethal infection  
115 (30), and in lymphocytic choriomeningitis virus infection, where late IL-6 production was  
116 shown to be critical for clearance of persistent viruses (12). However, failure to clear  
117 viruses from infected macrophages could result in viral spread to other tissues due to  
118 macrophage chemotaxis and could result in persistent IL-6 expression in affected tissue  
119 causing inflammatory pathologies and autoimmune diseases (16). Similarly, persistent  
120 elevation of IL-6 is associated with the development of experimental autoimmune

121 encephalomyelitis (EAE) in mice immunized peripherally with myelin peptides, whereas  
122 mice deficient in IL-6 are resistant to development of EAE (32, 40). Such data suggests  
123 that IL-6 expression may be essential to control early viral infection, yet contribute to  
124 pathology when expressed persistently.

125         Theiler's murine encephalomyelitis virus (TMEV) causes an acute infection that is  
126 cleared through innate and adaptive immune responses in most mouse strains yet  
127 produces debilitating sequelae in susceptible mice (33). C57Bl/6 and B10.S mice are  
128 prototypical TMEV-resistant strains whereas SJL/J mice fail to clear TMEV from  
129 macrophages and dendritic cells, resulting in persistent infection (25). Subsequent  
130 infection of macrophages in the central nervous system (CNS) of SJL/J mice leads to a  
131 demyelinating disease similar to human multiple sclerosis(5). Several reports have  
132 shown that a week after infection with TMEV, CNS macrophages, microglial cells, and  
133 astrocytes from SJL/J mice express IL-6 to a greater extent than C57Bl/6 mice (14, 17,  
134 34). A proposed theory is that chronic production of IL-6 in the CNS of persistently  
135 infected SJL/J mice contributes to demyelinating disease in response to TMEV(17).

136         We propose that the dichotomy of TMEV persistence seen in resistant and  
137 susceptible macrophages depends on early differences in the cytokine production. We  
138 have previously shown that B10.S macrophages produce more IL-12 p70 in response to  
139 TMEV infection than SJL/J macrophages *in vitro* (37). Significantly, when SJL/J  
140 macrophages are pretreated with IL-12, TMEV replication is reduced to levels  
141 comparable with B10.S macrophages. Paradoxically, SJL/J macrophages express  
142 more interferon (IFN) $\beta$  than B10.S macrophages before and during infection with

143 TMEV(37), yet this enhanced endogenous IFN $\beta$  expression is insufficient to control  
144 TMEV replication. However, pretreatment with additional exogenous recombinant IFN $\beta$   
145 decreases TMEV replication in SJL/J macrophages to amounts seen in B10.S  
146 macrophages (37). We found that p38 ERK MAPK pathways are activated in response  
147 to TMEV infection of B10.S or SJL/J macrophages and play a role in IL-12 and IFN $\beta$   
148 expression (37). Therefore, differences in MAPKs, IL-12, and IFN $\beta$  may be related to  
149 disparities in susceptibility of macrophages to persistent infection with TMEV.

150 In order to further understand the differences between SJL/J and B10.S mice that  
151 could account for differences in TMEV persistence, we examined IL-6 expression in  
152 macrophages during the early response to TMEV. Unexpectedly, we found that during  
153 early response to infection with TMEV, B10.S macrophages expressed significantly  
154 more IL-6 than SJL/J macrophages. Moreover, pre-treatment of macrophages with  
155 exogenous IL-6 reduced TMEV replication but neutralization of endogenous TMEV-  
156 induced IL-6 with Fab antibodies failed to affect TMEV replication in SJL/J  
157 macrophages. Additionally, we show that TMEV-induced IL-6 secretion in  
158 macrophages depends on ERK-MAPK and is enhanced by IL-12. However, neither IL-  
159 6-induced nitric oxide production nor STAT1 activation are essential to IL-6 reducing  
160 TMEV infection.

## 161 **MATERIALS AND METHODS**

162 **Mice, virus, cell lines, and reagents.** The eight to twelve week old female B10.S and  
163 SJL/J mice were obtained from Jackson Laboratories (Bar Harbor, Maine). RAW264.7  
164 cells were obtained from the American Type Culture Collection (Rockville, MD) and

165 maintained in DMEM with 10% FBS with 50 µg/ml gentamycin. The p38 MAP-kinase  
166 inhibitor SB203580 and ERK MAPK inhibitor U0126 were obtained from Promega  
167 Corporation (Madison, WI), *E. coli* LPS O127:B8 was obtained from Sigma Chemical  
168 Co.(St. Louis, MO), and poly I:C was obtained from InvivoGen (San Diego, CA). The  
169 inducible nitric oxide synthase inhibitor, L-NIL, was obtained from TOCRIS Bioscience  
170 (Bristol, UK) and incubated with macrophages where indicated at 10 µM. The nitric  
171 oxide donor spermine NONOate was obtained from Invitrogen (Carlsbad, CA) and  
172 incubated with macrophages where indicated at 1 mM. Affinity purified neutralizing rat  
173 antibody to mouse IL-6 (clone MP5-20F3) (48) was obtained from Invitrogen. Fab  
174 fragments of neutralizing anti-IL-6 were obtained after papain digestion using the Fab  
175 preparation kit of Pierce (Thermo Fisher Scientific, Rockford, IL). The initial stock of the  
176 DA strain of TMEV was obtained from Dr. Kristen Drescher, Department of Medical  
177 Microbiology and Immunology, Creighton University, Omaha, Nebraska. TMEV was  
178 grown in BHK-21 cells. The titer of stock cultures of TMEV was  $2.5 \times 10^6$  PFU/ml and  
179 macrophages cultures were infected with  $2.5 \times 10^5$  PFU of TMEV unless otherwise  
180 stated.

181 **Macrophage preparations.** Macrophages were elicited by intraperitoneal  
182 injection of 2 ml thioglycollate broth into mice. Four days later, the peritoneal cavities  
183 were each flushed with 2 ml DMEM and the cells were incubated at  $1 \times 10^6$  cells/2 ml  
184 of DMEM cell culture medium (Invitrogen, Carlsbad, CA) containing 10% fetal bovine  
185 serum (FBS) (Invitrogen), and 50 µg/ml gentamycin (Invitrogen). After 24 h, non-  
186 adherent cells were removed and 1 ml of culture medium added. Adherent cells were



187 greater than 90% Mac-1<sup>+</sup> as determined by FACS analysis. These macrophages were  
188 either untreated or pretreated for 30 min with 20  $\mu$ M of SB203580, 40  $\mu$ M U0126, 1  
189 ng/ml IL-12 p70 (p35/p40; BD-Pharmingen, San Diego, CA), 10 ng/ml IFN $\beta$   
190 (Minneapolis, MN) or 10 ng/ml IL-12 p40 (BD-Pharmingen). Untreated or pretreated  
191 macrophages were uninfected, infected with 100  $\mu$ l of the TMEV stock ( $2.5 \times 10^5$  PFU),  
192 stimulated with 1  $\mu$ g/ml LPS, or stimulated with 50 $\mu$ g poly I:C or left unstimulated. After  
193 3, 8, or 24 h of infection or stimulation, supernatants were collected for ELISA and cell  
194 extracts were collected for RNA preparation and quantitative RT-PCR.

195 **RNA interference.** Validated inhibitory siRNA targeting mouse STAT1 and  
196 STAT3 was purchased from Cell Signaling, Inc. (Danvers, MA) and transfected into  
197 RAW264.7 cells according to manufacturer's specifications using the nucleofection kit of  
198 Amaxa 36 h prior to challenge with TMEV.

199 **RNA preparation and quantitative RT-PCR.** RNA was extracted from cells  
200 using the RNeasy kit of Qiagen (Valencia, CA), PerfectPure kit from 5Prime  
201 (Gaithersburg, MD), or the Purelink kit from Ambion/Invitrogen (Carlsbad, CA) according  
202 to the manufacturer's specifications. One-hundred ng to one  $\mu$ g of RNA was reverse  
203 transcribed in 0.5 mM each of dATP, dGTP, dTTP, and dCTP, 20 U of RNase inhibitor  
204 with Superscript II reverse transcriptase (Invitrogen) at 42°C for 1.5 h followed by 94°C  
205 for 5 min. One twenty-fifth of the cDNA sample was incubated with 0.4  $\mu$ M of the  
206 following primer pairs (Invitrogen): IFN $\beta$  sense 5' ATGAACAACAG GTGGATCCTCC 3'  
207 and anti-sense 5' AGGAGCTCCTGACATTTCCGAA 3'; IL-6 sense 5' ATGAAGTTCCT  
208 CTCTGCAAGAGACT 3' and antisense 5' CACTAGGTTTGCC GAGTAGATCTC 3';

209 IRF1 sense 5' ATGCCAATCACTCGAATGCGGA 3' and antisense 5' GGCTGC  
 210 CACTCAGACTGTTCAA 3'; IRF-7 sense 5' CCAGCGAGTGC TGTTTGGAGAC 3' and  
 211 antisense 5' TTCCCTATTTCCGTGGCTGGG 3'; IRF-9 sense 5' ATGGCCTCA  
 212 GGCAAAGTACGCT 3' and antisense 5' TTCGCTTGCATGG TGATTTCTG 3'; TMEV  
 213 sense 5' CTTCCCATTC TACTGCAATG 3' and antisense 5' GTGTTCTGG  
 214 TTTACAGTAG3'; or GAPDH sense 5'-TTGTCAGCAA TGCATCCTGCAC-3' and  
 215 antisense 5'-ACAGCTTTCCA GAGGGGCCATC-3'. Quantitative PCR reactions were  
 216 run on an ABI Prism 7000 thermal cycler at 50 °C for 2 min, 95 °C for 10 min, 45 cycles  
 217 of 95 °C for 15 s/60 °C for 30 s. Cycle thresholds (Ct) of sample were normalized to Ct  
 218 of GAPDH for that sample ( $\Delta$ Ct), and then normalized to the average  $\Delta$ Ct of the control  
 219 samples ( $\Delta\Delta$ Ct) after which data are expressed as relative levels of mRNA using  $2^{-\Delta\Delta$ Ct}.  
 220 When macrophages are pretreated, TMEV RNA data are reported as percentage of  
 221 TMEV RNA in non-treated but infected macrophages.

222 **ELISAs.** ELISA plates were coated with antibodies to mouse IL-6 (MP5-  
 223 20F3;BD-Pharmingen), the plates were blocked with PBS/10% FBS. After washes, cell  
 224 culture supernatants or serial dilutions of recombinant IL-6 (BD-Pharmingen) were  
 225 added to wells. After 2 h, biotinylated antibody to mouse IL-6 (MP5-32C11;BD-  
 226 Pharmingen) was added to each well. After 1 h, streptavidin horseradish peroxidase  
 227 (1:1000; BD-Pharmingen) was added for 30 min and then 3,3', 5, 5'  
 228 Tetramethylbenzidine substrate/hydrogen peroxide solution (BD-Pharmingen) was  
 229 added to each well. IL-6 was measured by determining optical densities at OD 450 nm  
 230 wavelength with reference OD 570 nm using an ELISA spectrophotometric plate reader.

231           **PAGE and western blot analysis.** Cell lysates were obtained from RAW cells  
232 challenged with TMEV with or without treatment with 10 ng/ml recombinant murine IL-6  
233 (BD-Pharmingen, San Diego, CA) or murine recombinant IFN $\beta$  (Interferon Source,  
234 Piscataway, NJ). Twenty  $\mu$ l of each sample containing 20  $\mu$ g of protein in sample buffer  
235 with bromophenol blue was run on a 10% SDS, Tris-glycine- polyacrylamide gel and  
236 transferred to a nitrocellulose membrane. The membrane was treated with blocking  
237 buffer for 1 h at room temperature, followed by incubation in 1:500 dilution of rabbit IgG  
238 anti-phospho-STAT1(Invitrogen, Camarillo, CA), 1:1000 dilution of anti-phospho-STAT3  
239 (Cell Signalling, Beverly, MA), 1:500 dilution of anti-mouse STAT1(Invitrogen), 1:1000  
240 dilution of anti-mouse STAT3 (Invitrogen), or 1:500 mouse anti-tubulin E7  
241 (Developmental Studies Hybridoma Bank, University of Iowa) and then 1:5000 dilution  
242 of IRDye $\text{\textcircled{R}}$  800CWGoat Anti-Rabbit IgG (Rockland Immunochemicals, Inc., Gilbertsville,  
243 PA) or Alexa Fluor680-labeled anti-Mouse IgG (Rockland Immunochemicals). The  
244 washed membrane was scanned with a LICOR Odyssey $\text{\textcircled{R}}$  Infrared Imaging System and  
245 densitometric analysis done with LICOR imaging software.

246           **Nitric Oxide Assay.** Induction of nitric oxide (NO) was assayed in culture  
247 supernatants by measuring nitrite using the Greiss reagent kit of Invitrogen. Briefly, 20  
248  $\mu$ L of Griess Reagent was mixed with 150  $\mu$ L of supernatant plus 130  $\mu$ L of deionized  
249 water and incubated for 30 min at room temperature. Color development at 570 nm,  
250 which is proportional to nitric oxide in supernatants, was measured with a  
251 spectrophotometer.

252           **Statistical analysis.** The Student's *t* test of the GraphPad Prism Software was  
253 used to determine the significance of differences between means;  $p < 0.05$  was  
254 considered significant.

255

## 256   **RESULTS**

### 257   **TMEV-induced IL- 6 production is greater in TMEV-resistant B10.S macrophages.**

258 IL-6 contributes to both antiviral immunity and virus-induced pathology (3, 41). TMEV is  
259 cleared from macrophages of TMEV-resistant mice but persists in (CNS) macrophages  
260 in susceptible SJL/J mice (5). Therefore, IL-6 mRNA and protein from TMEV-  
261 susceptible SJL/J and TMEV-resistant B10.S macrophages were determined following  
262 challenge with TMEV. IL-6 mRNA was detected at 3 and 8 h after TMEV challenge  
263 (Fig. 1 A) in both B10.S and SJL/J macrophages, however B10.S macrophages  
264 produced more IL-6 mRNA than SJL/J macrophages at 8 h after infection. Similarly,  
265 B10.S macrophages produced significantly more IL-6 protein 8 h after TMEV challenge  
266 compared with SJL/J macrophages in response to TMEV (Fig. 1 B). Because TMEV  
267 RNA replication is significantly higher in SJL/J macrophages compared with B10.S  
268 macrophages (37), the data here suggest that enhanced early IL-6 production may  
269 contribute to better control of TMEV replication.

270           To determine if exogenous IL-6 could impede TMEV replication in SJL/J  
271 macrophages during early infection, recombinant IL-6 was added to SJL/J macrophages  
272 30 min prior to TMEV infection and either left in the media during infection or washed

273 out by changing the media just prior to infection. Treatment with 10 ng/ml exogenous  
274 IL-6 significantly reduced TMEV replication at 8h post infection (PI) in SJL/J  
275 macrophages (Fig. 1C). In addition, treatment with as little as 0.1 ng/ml of exogenous  
276 IL-6 significantly reduced TMEV RNA at 24 h PI in SJL/J macrophages (Fig. 1D).  
277 Equally significant, IL-6 preemptively triggered anti-viral activity in SJL macrophages  
278 even when exogenous IL-6 was added for only 30 min and washed out prior to infection  
279 (Fig. 1D). These findings suggest that early differences in the amount of IL-6 present  
280 during the course of infection play an important role in the establishment of TMEV in  
281 SJL/J macrophages.

282 To examine whether TMEV-induced endogenous IL-6 contributes to control of  
283 TMEV replication in SJL/J mice, neutralizing anti-IL-6 (IgG; clone MP5-20F3) antibody  
284 was added to SJL/J macrophages at the time of TMEV infection (48). Surprisingly,  
285 addition of either whole anti-IL-6 antibody or isotype control antibody both resulted in a  
286 significant reduction in TMEV replication (see Fig. S1A and S1B). These findings do not  
287 preclude an antiviral role for IL-6, since interactions between the IgG antibody and Fc $\gamma$   
288 receptors may also induce anti-viral immunity in macrophages. Indeed addition of  
289 isotype antibody had a similar effect on TMEV infection of macrophages. Therefore we  
290 treated the whole antibody with papain and removed the Fc fragments to generate Fab  
291 anti-IL-6. Addition of Fab anti-IL-6 to SJL/J macrophages at the time of infection did not  
292 decrease but also did not increase TMEV replication in infected SJL/J macrophages  
293 (Fig. 1E).

294 To determine the time frame at addition in which exogenous IL-6 is most effective  
295 at reducing TMEV RNA we added 10 ng/ml recombinant IL-6 at 1, 3, 6, and 7 h after or  
296 exogenous IFN $\beta$  7h PI of SJL/J macrophages with TMEV. Addition of exogenous IL-6  
297 at 1 and 3 h PI significantly reduced TMEV RNA compared with untreated SJL/J  
298 macrophages that were infected with TMEV (Fig. 1F). Addition of IL-6 at 6 and 7 h PI  
299 failed to significantly reduce TMEV RNA in SJL/J macrophages compared to untreated  
300 macrophages. In contrast addition of IFN $\beta$  at 7 h PI significantly reduced TMEV RNA in  
301 SJL/J macrophages. Thus, while small amounts of added IL-6 were capable of  
302 controlling TMEV replication in SJL/J macrophages in vitro, it appears that a sufficient  
303 quantity of endogenous IL-6 is not produced quickly enough to provide protection  
304 because the beneficial effects of IL-6 are only evident prior to 6 h PI. These results  
305 confirm that the antiviral effect of IL-6 is most effective very early during the course of  
306 TMEV infection of macrophages. These findings warrant further investigation into strain  
307 differences in early TMEV-induced IL-6 expression.

308 **TMEV-induced IL-6 expression in macrophages is dependent on ERK**  
309 **MAPK.** The cell signaling pathway activated by TMEV infection leading to IL-6  
310 expression in macrophages is not well understood. We previously showed that TMEV  
311 infection of macrophages from both B10.S and SJL/J mice strongly activates ERK  
312 MAPK and weakly activates p38 MAPK (37). Therefore, macrophages from B10.S and  
313 SJL/J mice were pretreated with ERK and p38 MAPK inhibitors 30 min prior to TMEV  
314 infection. Twenty-four h after infection, the amount of TMEV-induced IL-6 protein  
315 secreted by B10.S macrophages was significantly greater than that secreted by SJL/J

316 macrophages (Fig. 2 B). In contrast the 24 h TMEV-induced IL-6 mRNA was  
317 significantly higher in SJL/J macrophages than B10.S macrophages, suggesting  
318 translational control of IL-6. Moreover, TMEV-induced IL-6 mRNA expression was  
319 increased in macrophages treated with the p38 MAPK inhibitor, SB203580, (Fig. 2 A)  
320 but IL-6 protein production was unchanged (Fig. 2 B). Pretreatment of macrophages  
321 with the ERK MAPK inhibitor, U0126, significantly decreased TMEV induction of IL-6  
322 mRNA (Fig. 2 A) and protein (Fig. 2 B) in both SJL/J and B10.S macrophages  
323 responding to TMEV. These results indicate that ERK MAPK activation is required for  
324 IL-6 expression in response to TMEV infection and suggest that TMEV-induced IL-6  
325 expression is under translational control.

326 We have also shown that the TLR3 pathway is involved in macrophage cytokine  
327 responses to TMEV infection by inducing IFN $\beta$ , IL-12, and IL-23 (1). Therefore, we  
328 compared the response of macrophages from each strain when treated with the TLR3  
329 agonist, polyI:C or the TLR4 agonist, LPS. B10.S macrophages expressed significantly  
330 more IL-6 mRNA and protein than SJL/J macrophages treated with LPS (Fig. 3 A, B) or  
331 polyI:C (Fig. 3 C, D), but in macrophages from both strains the level of IL-6 expression  
332 in response to poly I:C was substantially less than that in response to LPS or TMEV.  
333 Because ERK MAPK is activated by both TLR3 agonists (see Fig. S2) and TLR4  
334 agonists (37), macrophages were also pretreated with U0126. Inhibition of ERK MAPK  
335 activation with U0126 significantly reduced IL-6 expression to the same extent in both  
336 B10.S and SJL/J macrophages responding to LPS and significantly decreased IL-6  
337 production from SJL/J macrophages responding to polyI:C. Therefore, ERK MAPK

338 activation is required for IL-6 expression that occurs from activation of TLR3 and TLR4  
339 pathways in SJL/J macrophages.

340 **IL-6 expression by macrophages in response to TMEV is enhanced by IL-**  
341 **12.** Previously, we showed that SJL/J macrophages responding to TMEV express  
342 significantly more IFN $\beta$  and IL-12 p40 (p40/p40) but significantly less IL-12 p70  
343 (p35/p40) than B10.S macrophages. In addition we showed that addition of IFN $\beta$ , IL-12  
344 p40, or p70 decreases TMEV replication in SJL/J macrophages (37). To see if these  
345 cytokines could affect TMEV-induced IL-6, B10.S and SJL/J macrophages were  
346 pretreated with the IL-12 p40 homodimer, bioactive IL-12 p70, or IFN $\beta$  30 min prior to  
347 and during TMEV challenge. Treatment with IL-12 p70 or p40 significantly enhanced IL-  
348 6 expression in response to TMEV in both B10.S and SJL/J macrophages at 24 and/or  
349 48 h post TMEV infection (Fig. 4 A, B), however the modest increase in IL-6 expression  
350 upon pretreatment with IFN $\beta$  was not significant (Fig. 4 ). As we have seen before, IL-  
351 12 and IFN $\beta$  treatment reduced TMEV replication in both B10.S and SJL/J  
352 macrophages (Fig. 4C). Therefore deficient production of IL-12 during response to  
353 TMEV could contribute to diminished IL-6 expression by SJL/J macrophages.

354 To gain further insight into the role of IL-12 in IL-6 expression, we used the  
355 RAW264.7 macrophage cell line which is permissive for TMEV replication, expresses  
356 IL-6 well, expresses IL-12 poorly, and activates ERK MAPK following TMEV infection  
357 (26, 37). To confirm the effects of IL-12 and the ERK MAPK inhibitor on IL-6 expression,  
358 RAW264.7 cells were treated with IL-12 with or without U0126 during TMEV infection.  
359 Pretreatment with IL-12 enhanced IL-6 mRNA (Fig. 5 A) and protein (Fig. 5 B)



360 expression following TMEV challenge of RAW264.7 cells, while pretreatment with  
361 U0126 repressed the IL-12 enhancement of IL-6 (Fig. 5 A, B). Therefore IL-12 is  
362 involved in IL-6 expression during TMEV infection of macrophages in an ERK MAPK  
363 dependent manner.

364 **IL-6 has direct antiviral activity against TMEV replication in RAW264.7 cells.**

365 We next set out to determine if, when, and at what quantity IL-6 represses TMEV  
366 infection in RAW264.7 cells. First, RAW264.7 cells were treated with 10 ng/ml  
367 exogenous recombinant IL-6 at 30 min prior to and during challenge with TMEV. Like  
368 SJL/J macrophages, treatment of RAW264.7 cells with IL-6 significantly decreased  
369 TMEV RNA at 24 h PI (Fig. 6 A). Interestingly, as little as 0.1 ng/ml significantly  
370 reduced TMEV RNA 24 h PI, (Fig. 6B). Furthermore, IL-6 treatment enhanced IL-6  
371 expression (Fig. 6C) at 24 h PI, but IL-6 treatment did not decrease TMEV RNA at 3  
372 and 8 h PI in RAW264.7 cells (Fig. 6D), suggesting the time period in which TMEV  
373 induced events occur in RAW264.7 are slightly different than that seen in SJL/J  
374 macrophages.

375 **IL-6 antiviral activity against TMEV replication is associated with activation**

376 **of STAT1 related anti-viral activity.** Like IFN $\beta$ , treatment of macrophages with IL-6  
377 leads to activation of STAT3 by phosphorylation at its tyrosine 705 and activation of  
378 STAT1 at its tyrosine 701 (13). IL-6 activation of STAT1 could contribute to control of  
379 virus replication (28, 46) by inducing expression of IRF1(22), IRF7(23), and IRF9 that  
380 enhance IFN $\beta$  expression. To determine if IL-6 activates STAT1 and induces  
381 expression of IRFs that are downstream of STAT1 during TMEV infection of

382 macrophages, RAW264.7 cells were treated with IL-6, IFN $\beta$ , or TMEV alone or in  
383 combination and phospho-STAT1, as well as phospho-STAT3 were evaluated by  
384 western blot. TMEV infection alone failed to activate STAT1 by 30 min PI but did so by  
385 6 h PI (Fig. 7A). In contrast, treatment of RAW264.7 cells with either IL-6 or IFN $\beta$   
386 activated STAT1 and STAT3 as early as 30 min after TMEV challenge. To determine if  
387 STAT1 activation is involved in control of TMEV replication, RAW264.7 cells were  
388 transfected with small inhibitory RNA that has been verified to reduce STAT1 (siSTAT1)  
389 or siSTAT3. Knockdown of STAT1 resulted in significantly increased TMEV RNA while  
390 knockdown of STAT3 failed to affect TMEV RNA replication in RAW264.7 cells (Fig.  
391 7B). However, pretreatment of STAT1-knocked-down RAW264.7 cells with exogenous  
392 IL-6 resulted in a reduction of TMEV infection back to a level seen in RAW264.7 cells  
393 infected with TMEV. Moreover, addition of exogenous IL-6 to STAT3-knocked down  
394 and STAT1/STAT3 double-knocked down RAW264.7 cells significantly reduced TMEV  
395 RNA replication compared with untreated RAW264.7 cells infected with TMEV (Fig. 7B).  
396 Therefore, STAT1 activation contributes to control of TMEV replication and the  
397 activation of STAT3 may negatively impact the antiviral effect of STAT1. Consistent  
398 with STAT1 activation, treatment of RAW264.7 cells with IL-6 resulted in significant  
399 enhancements of TMEV-induced IRF1 (Fig. 7C), IRF7 (Fig. 7D), IRF9 (Fig. 7E), and  
400 IFN $\beta$  (Fig. 7F). These results confirm that IL-6 can directly limit viral replication in  
401 macrophages early after infection with TMEV by initiating a STAT1 anti-viral program.

402 **IL-6 induces nitric oxide in RAW264.7 cells.** IL-6 (42) and activated STAT1  
403 (29) also induce expression of nitric oxide synthase and nitric oxide production, which is

404 a potent anti-viral factor (41). Therefore, another possibility is that IL-6 controls TMEV  
405 replication in macrophages by inducing nitric oxide. To explore this possibility, TMEV-  
406 challenged RAW264.7 cells were treated with 10 ng/ml IL-6 in the presence or absence  
407 of L-NIL, an inducible nitric oxide synthase inhibitor (27), and production of nitric oxide  
408 was measured. TMEV infection of RAW264.7 cells did not result in significant nitric  
409 oxide production at 24 h PI (see Fig. S3A). However, addition of exogenous IL-6 to  
410 TMEV infected RAW264.7 cell resulted in a significant enhancement of nitric oxide  
411 secretion, which was prevented by L-NIL. IL-6 or NONOate, a nitric oxide donor,  
412 repressed TMEV replication in RAW264.7 cells (see Fig. S3B). However, L-NIL did not  
413 reverse the IL-6 repression of TMEV replication in RAW264.7 cells. Thus, while IL-6  
414 induces nitric oxide and nitric oxide is anti-viral, it does not play the decisive role in the  
415 IL-6-induced antiviral effect observed in macrophages in vitro.

416

## 417 **DISCUSSION**

418 The data herein show that IL-6 protects macrophages early after infection with TMEV by  
419 decreasing virus replication in macrophages. These results are somewhat surprising  
420 because research on chronic TMEV infection has supported the theory that enhanced  
421 IL-6 may play a detrimental role in the immunopathology of TMEV in susceptible mice  
422 (14). Indeed, a recent report has shown that SJL/J mice infected with TMEV exhibit  
423 greater levels of IL-6 in the CNS than TMEV-resistant C57Bl/6 mice starting at day 8  
424 after infection (14). However, in that report IL-6 was not measured until day-8 after  
425 infection. We show herein that the antiviral effect of IL-6 takes place within the first 6 h

426 after infection. Therefore the beneficial effect of IL-6 is early after TMEV infection, but  
427 not later. It is also possible that TMEV-resistant C57Bl/6 mice may have a different  
428 mechanism to control TMEV infection in macrophages compared with TMEV-resistant  
429 B10.S mice. The results herein indicate that during the first day after infection in vitro,  
430 IL-6 protein production by macrophages from TMEV-resistant B10.S mice is greater  
431 than from macrophages of TMEV-susceptible SJL/J and this difference may play an  
432 important role in determining the severity of the infection. Moreover, significant levels of  
433 IL-6 are not produced until 8 h after TMEV infection and IL-6 is only effective at  
434 controlling TMEV replication during the first 8 h after infection. Therefore, IL-6 produced  
435 by the TMEV infected macrophages helps to protect uninfected macrophages from  
436 subsequent infection. Altogether, the present data suggest the CNS demyelination that  
437 SJL/J mice develop several weeks to months after infection with TMEV may be due in  
438 part to an inability of SJL/J macrophages to control viral replication very early after the  
439 initial infection. Similarly, B10.S mice that do not develop demyelination following  
440 infection with TMEV may stem from better early control of TMEV replication in  
441 macrophages. Our results indicate that ability to control TMEV replication in  
442 macrophages begins during the first 24h after infection is to a large extent dependent on  
443 macrophage production of IL-6, which then protects uninfected cells from infection.  
444 Thus, IL-6 production by virally infected cells may have dual roles in viral infection by  
445 contributing to both anti-viral immunity and subsequent pathology.

446 Several reports have proposed beneficial effects of exogenous IL-6 treatment in  
447 TMEV infection of SJL/J mice (35, 39). In fact, recombinant exogenous IL-6 was shown  
448 to suppress chronic demyelination and reduce virus replication in SJL/J mice infected

449 with TMEV. Other reports have shown that IL-6 is involved in neuronal health facilitating  
450 neuronal differentiation, neurite outgrowth, survival, regeneration (47) and  
451 oligodendrocyte differentiation (52). In stark contrast to these reports are others which  
452 have suggested that chronic expression of IL-6 in TMEV-susceptible SJL/J mice is  
453 responsible for development of demyelinating disease (17). Hou et al. (14) showed that  
454 IL-6 levels in the brain and spinal cord of SJL/J mice are elevated 8-90 days post  
455 infection compared with TMEV resistant C57Bl/6 mice. It is possible that the lower early  
456 acute IL-6 response of SJL/J mice to TMEV infection is inadequate to control viral  
457 replication, thereby rendering these mice susceptible to chronic viral infection and a  
458 chronically elevated IL-6 response. Alternatively, enhanced IL-6 in the CNS of TMEV  
459 infected SJL/J mice may not be a reflection of a heightened IL-6 response of infected  
460 cells to TMEV but may be due to enhanced infiltration of macrophages into the CNS,  
461 which then are infected with TMEV and produce IL-6.

462         The anti-viral and pathological effects of IL-6 notwithstanding, the signaling  
463 pathways that lead to IL-6 expression in response to TMEV remain unclear. Previous  
464 reports showed that ERK and p38 MAPK pathways are involved in both innate anti-viral  
465 immunity (7) and IL-6 expression (49). The results herein confirm that activation of ERK  
466 MAPKs contributes to IL-6 expression in response to TMEV. However, maximum ERK  
467 activation occurs within 30 min after TMEV infection of macrophages and we have not  
468 noticed differences in the intensity of ERK activation between SJL and B10.S  
469 macrophages following TMEV infection (37, 38). Activated ERK MAPKs have been  
470 shown to phosphorylate downstream transcription factors such as the cyclic AMP  
471 response element binding protein (CREB) at serine 133 (4, 51, 45), CREB response

472 elements (CRE) are located at the IL-6 promoter (8), activated CREB binds to IL-6  
473 promoter CRE (10), and CREB activity is required for IL-6 (36). Therefore, TMEV  
474 induction of IL-6 is likely to depend on the ERK activation of the transcription factor,  
475 CREB.

476         The results herein also suggest that TMEV induction of IL-6 in macrophages is in  
477 part dependent upon macrophage production of IL-12, which is also the primary inducer  
478 of Th1 development during adaptive immune responses (15). In addition, our results  
479 show that IL-12 enhancement of IL-6 expression is dependent on the ERK-MAPK  
480 pathway. Other reports have shown in IL-12 p35 knockout mice that IL-12 plays a  
481 significant role in IL-6 expression during viral infection (18), IL-6 expression depends in  
482 part on activation of ERK (19), and that IL-12 activates ERK-MAPK (20). These results  
483 are consistent with our previous data that showed significantly more IL-12 production  
484 from B10.S macrophages compared with SJL/J macrophages following TMEV infection  
485 (37). Therefore, IL-12 expression by macrophages that is responsible for Th1  
486 development during adaptive immune responses also contributes to IL-6 expression  
487 during innate immune responses, both roles of IL-12 are likely to be beneficial to early  
488 anti-viral immunity.

489         Overall the data here suggest a direct, interferon-like role for IL-6 in the  
490 suppression of TMEV replication, which is exemplified by its ability to activate STAT1  
491 and induce expression of STAT1 dependent genes. Indeed early reports regarding IL-  
492 6, which was originally called Interferon-beta 2, showed it to have anti-viral activity (44).  
493 While it is not clear how IL-6 decreases TMEV replication, it is known to be a potent

494 inducer of iNOS and nitric oxide production (31), which has anti-viral properties (41).  
495 Our data confirm that IL-6 induces significant nitric oxide production from macrophages  
496 and that nitric oxide does indeed control TMEV replication. However, preventing IL-6-  
497 induced nitric oxide production did not reverse the anti-viral effect of IL-6. This is most  
498 likely because IL-6 also activated STAT1 and STAT3 that lead to the induction of  
499 STAT1-downstream anti-viral genes, IRF1, IRF7, and IRF9. Therefore the data suggest  
500 that early after infection with TMEV, IL-6 controls acute viral infection by inducing  
501 multiple innate antiviral programs.

502         One of the discrepancies noted herein is that between IL-6 mRNA expression  
503 and IL-6 protein secretion at time points beyond 8 h post infection. These data suggest  
504 that TMEV induced IL-6 production is under translational or inhibitory RNA control,  
505 which is especially notable in the TMEV-susceptible SJL/J macrophages. At 24 and 28  
506 h post infection SJL/J macrophages expressed more IL-6 mRNA but produced less IL-6  
507 compared with B10.S macrophages. In one study K homology (KH)-type splicing  
508 regulatory protein (KHSRP) was shown to target the AU-rich elements in the IL-6 mRNA  
509 3' untranslated region which restricts its translation but does not lead to degradation of  
510 mRNA (6). In another study microRNA-365 was shown to inhibit the translation of IL-6  
511 mRNA without affecting IL-6 mRNA (50). Our results suggest that differences in  
512 translational or inhibitory control of IL-6 expression following infection of macrophages  
513 could contribute to susceptibility versus resistance of macrophages to TMEV.

514         In summary, our findings show that strain differences in onset and amount of IL-6  
515 upregulation upon infection with TMEV correlate with previously observed differences in

516 disease outcomes between B10.S and SJL/J mice. Whereas B10.S mice with  
517 enhanced early IL-6 are capable of clearing TMEV, in SJL/J mice TMEV replication is  
518 not reduced by IL-6 unless exogenous IL-6 is added, suggesting that these mice do not  
519 produce IL-6 quickly enough or in sufficient quantity to control viral replication. The  
520 present investigation suggests that enhanced early expression of IL-6 by macrophages  
521 during infection with a macrophage-trophic virus, TMEV, is responsible for better control  
522 of viral replication in TMEV-resistant B10.S macrophages compared with TMEV-  
523 susceptible SJL/J macrophages. These results indicate that the persistent TMEV  
524 infection of macrophages in the CNS of SJL/J mice could be the result of insufficient IL-  
525 6 production that controls early TMEV infection through multiple anti-viral mechanisms  
526 that include nitric oxide production and STAT1 activation. Furthermore, the present  
527 data clearly shows that IL-12 and ERK MAPK play a major role in TMEV-induced IL-6.  
528 While ERK MAPK is required for IL-6 expression in response to TMEV its activation is  
529 equivalent in both strains. In contrast our previous studies point to differences in TLR  
530 signaling and IL-12 expression that may contribute to strain differences in TMEV-  
531 induced IL-6. These results are supported by our data showing that TMEV-induced IL-6  
532 is also antiviral in RAW264.7 cells, IL-6 expression is enhanced by IL-12, and IL-6 leads  
533 to activation of STAT1, expression of its downstream genes IRF1, IRF7, IRF 9, and IFN-  
534 beta, as well as induction of nitric oxide. These results are significant because  
535 persistent infection of SJL/J macrophages by TMEV contributes to the inflammatory  
536 autoimmune demyelination in the CNS of TMEV infected SJL/J mice, which mimics  
537 human Multiple Sclerosis, that is not seen in TMEV infected B10.S mice. It remains to  
538 be determined if IL-6 is effective at controlling TMEV replication during established



539 persistent TMEV infection of macrophage populations. Nevertheless, this study  
 540 suggests therapeutic strategies that promote early IL-6 antiviral pathways could prevent  
 541 chronic viral infections.

542

543 **ACKNOWLEDGEMENTS**

544 This work was supported by funding from the University of Nebraska Medical Center  
 545 College of Dentistry and University of Nebraska Lincoln, School of Biological Sciences,  
 546 and supported by Award Number 8P30GM10350903 and 5P20GM103489 from the  
 547 National Center for Research Resources (NCRR), a component of the National  
 548 Institutes of Health (NIH). The content is solely the responsibility of the authors and  
 549 does not necessarily represent the official views of the National Center for Research  
 550 Resources or the National Institutes of Health.

551

552 **REFERENCES**

553

554 1. **Al-Salleh, F., and T. M. Petro.** 2007. TLR3 and TLR7 are involved in  
 555 expression of IL-23 subunits while TLR3 but not TLR7 is involved in expression  
 556 of IFN-beta by Theiler's virus-infected RAW264.7 cells. *Microbes Infect* **9**:1384-  
 557 92.

558 2. **Andoh, A., M. Shimada, S. Bamba, T. Okuno, Y. Araki, Y. Fujiyama, and T.**  
 559 **Bamba.** 2002. Extracellular signal-regulated kinases 1 and 2 participate in  
 560 interleukin-17 plus tumor necrosis factor-alpha-induced stabilization of

- 561 interleukin-6 mRNA in human pancreatic myofibroblasts. *Biochim Biophys Acta*  
 562 **1591**:69-74.
- 563 3. **Banerjee, S., K. Narayanan, T. Mizutani, and S. Makino.** 2002. Murine  
 564 coronavirus replication-induced p38 mitogen-activated protein kinase activation  
 565 promotes interleukin-6 production and virus replication in cultured cells. *J Virol*  
 566 **76**:5937-48.
- 567 4. **Cammarota, M., L. R. Bevilaqua, P. R. Dunkley, and J. A. Rostas.** 2001.  
 568 Angiotensin II promotes the phosphorylation of cyclic AMP-responsive element  
 569 binding protein (CREB) at Ser133 through an ERK1/2-dependent mechanism. *J*  
 570 *Neurochem* **79**:1122-8.
- 571 5. **Clatch, R. J., S. D. Miller, R. Metzner, M. C. Dal Canto, and H. L. Lipton.**  
 572 1990. Monocytes/macrophages isolated from the mouse central nervous system  
 573 contain infectious Theiler's murine encephalomyelitis virus (TMEV). *Virology*  
 574 **176**:244-54.
- 575 6. **Dhamija, S., N. Kuehne, R. Winzen, A. Doerrie, O. Dittrich-Breiholz, B. K.**  
 576 **Thakur, M. Kracht, and H. Holtmann.** 2011. Interleukin-1 activates synthesis of  
 577 interleukin-6 by interfering with a KH-type splicing regulatory protein (KSRP)-  
 578 dependent translational silencing mechanism. *J Biol Chem* **286**:33279-88.
- 579 7. **Dong, C., R. J. Davis, and R. A. Flavell.** 2002. MAP kinases in the immune  
 580 response. *Annu Rev Immunol* **20**:55-72.
- 581 8. **Droogmans, L., I. Cludts, Y. Cleuter, R. Kettmann, and A. Burny.** 1992.  
 582 Nucleotide sequence of the bovine interleukin-6 gene promoter. *DNA Seq* **3**:115-  
 583 7.

- 584 9. **Gadient, R. A., and U. Otten.** 1994. Identification of interleukin-6 (IL-6)-  
 585 expressing neurons in the cerebellum and hippocampus of normal adult rats.  
 586 *Neurosci Lett* **182**:243-246.
- 587 10. **Grassl, C., B. Luckow, D. Schlondorff, and U. Dendorfer.** 1999.  
 588 Transcriptional regulation of the interleukin-6 gene in mesangial cells. *J Am Soc*  
 589 *Nephrol* **10**:1466-77.
- 590 11. **Hama, T., M. Miyamoto, H. Tsukui, C. Nishio, and H. Hatanaka.** 1989.  
 591 Interleukin-6 as a neurotrophic factor for promoting the survival of cultured basal  
 592 forebrain cholinergic neurons from postnatal rats. *Neurosci Lett* **104**:340-4.
- 593 12. **Harker, J. A., G. M. Lewis, L. Mack, and E. I. Zuniga.** 2011. Late Interleukin-6  
 594 Escalates T Follicular Helper Cell Responses and Controls a Chronic Viral  
 595 Infection. *Science* **334**:825-829.
- 596 13. **Hemann, U., C. Gerhartz, B. Heesel, J. r. Sasse, G. n. Kurapkat, J.**  
 597 **Grotzinger, A. Wollmer, Z. Zhong, J. E. D. Jr, L. Graeve, P. C. Heinrich, and**  
 598 **F. Horn.** 1996. Differential Activation of Acute Phase Response Factor/Stat3 and  
 599 Stat1 via the Cytoplasmic Domain of the Interleukin 6 Signal Transducer gp130.  
 600 *J Biol Chem* **271**:12999-13007.
- 601 14. **Hou, W., H. S. Kang, and B. S. Kim.** 2009. Th17 cells enhance viral persistence  
 602 and inhibit T cell cytotoxicity in a model of chronic virus infection. *J Exp Med*  
 603 **206**:313-28.
- 604 15. **Hsieh, C. S., S. E. Macatonia, C. S. Tripp, S. F. Wolf, A. O'Garra, and K. M.**  
 605 **Murphy.** 1993. Development of TH1 CD4+ T cells through IL-12 produced by  
 606 *Listeria*-induced macrophages. *Science* **260**:547-9.

- 607 16. **Ishihara, K., and T. Hirano.** 2002. IL-6 in autoimmune disease and chronic  
 608 inflammatory proliferative disease. *Cytokine Growth Factor Rev* **13**:357-68.
- 609 17. **Jin, Y.-H., M. Mohindru, M. H. Kang, A. C. Fuller, B. Kang, D. Gallo, and B. S.**  
 610 **Kim.** 2007. Differential Virus Replication, Cytokine Production, and Antigen-  
 611 Presenting Function by Microglia from Susceptible and Resistant Mice Infected  
 612 with Theiler's Virus. *J Virol* **81**:11690-11702.
- 613 18. **Kapil, P., R. Atkinson, C. Ramakrishna, D. J. Cua, C. C. Bergmann, and S. A.**  
 614 **Stohlman.** 2009. Interleukin-12 (IL-12), but not IL-23, deficiency ameliorates viral  
 615 encephalitis without affecting viral control. *J Virol* **83**:5978-86.
- 616 19. **Kim, S. H., J. Kim, and R. P. Sharma.** 2004. Inhibition of p38 and ERK MAP  
 617 kinases blocks endotoxin-induced nitric oxide production and differentially  
 618 modulates cytokine expression. *Pharmacol Res* **49**:433-439.
- 619 20. **Kondadasula, S. V., J. M. Roda, R. Parihar, J. Yu, A. Lehman, M. A. Caligiuri,**  
 620 **S. Tridandapani, R. W. Burry, and W. E. Carson.** 2008. Colocalization of the  
 621 IL-12 receptor and FcγRIIIa to natural killer cell lipid rafts leads to activation of  
 622 ERK and enhanced production of interferon-γ. *Blood* **111**:4173-4183.
- 623 21. **Leonard, M., M. P. Ryan, A. J. Watson, H. Schramek, and E. Healy.** 1999.  
 624 Role of MAP kinase pathways in mediating IL-6 production in human primary  
 625 mesangial and proximal tubular cells. *Kidney Int* **56**:1366-77.
- 626 22. **Li, X., S. Leung, S. Qureshi, J. E. Darnell, Jr., and G. R. Stark.** 1996.  
 627 Formation of STAT1-STAT2 heterodimers and their role in the activation of IRF-1  
 628 gene transcription by interferon-alpha. *J Biol Chem* **271**:5790-4.

- 629 23. **Marie, I., J. E. Durbin, and D. E. Levy.** 1998. Differential viral induction of  
 630 distinct interferon-alpha genes by positive feedback through interferon regulatory  
 631 factor-7. *Embo J* **17**:6660-9.
- 632 24. **Mihara, M., M. Hashizume, H. Yoshida, M. Suzuki, and M. Shiina.** 2011. IL-  
 633 6/IL-6 receptor system and its role in physiological and pathological conditions.  
 634 *Clin Sci (Lond)* **122**:143-59.
- 635 25. **Monteyne, P., F. Bihl, F. Levillayer, M. Brahic, and J. F. Bureau.** 1999. The  
 636 Th1/Th2 balance does not account for the difference of susceptibility of mouse  
 637 strains to Theiler's virus persistent infection. *J Immunol* **162**:7330-4.
- 638 26. **Moore, T. C., F. M. Al-Salleeh, D. M. Brown, and T. M. Petro.** 2011. IRF3  
 639 polymorphisms induce different innate anti-Theiler's virus immune responses in  
 640 RAW264.7 macrophages. *Virology* **418**:40-8.
- 641 27. **Moore, W. M., R. K. Webber, G. M. Jerome, F. S. Tjoeng, T. P. Misko, and M.**  
 642 **G. Currie.** 1994. L-N6-(1-Iminoethyl)lysine: A Selective Inhibitor of Inducible  
 643 Nitric Oxide Synthase. *J Med Chem* **37**:3886-3888.
- 644 28. **Mott, K. R., D. Underhill, S. L. Wechsler, T. Town, and H. Ghiasi.** 2009. A role  
 645 for the JAK-STAT1 pathway in blocking replication of HSV-1 in dendritic cells and  
 646 macrophages. *Virology* **6**:56.
- 647 29. **Nishiya, T., T. Uehara, H. Edamatsu, Y. Kaziro, H. Itoh, and Y. Nomura.** 1997.  
 648 Activation of Stat1 and subsequent transcription of inducible nitric oxide synthase  
 649 gene in C6 glioma cells is independent of interferon- $\gamma$ -induced MAPK activation  
 650 that is mediated by p21ras. *FEBS Letters* **408**:33-38.

- 651 30. **O’Gorman, W. E., P. Sampath, E. F. Simonds, R. Sikorski, M. O’Malley, P. O.**  
652 **Krutzik, H. Chen, V. Panchanathan, G. Chaudhri, G. Karupiah, D. B. Lewis,**  
653 **S. H. Thorne, and G. P. Nolan.** 2010. Alternate mechanisms of initial pattern  
654 recognition drive differential immune responses to related poxviruses. *Cell Host*  
655 *Microbe* **8**:174-85.
- 656 31. **Oh, Y. S., Y.-J. Lee, E. Y. Park, and H.-S. Jun.** 2011. Interleukin-6 treatment  
657 induces beta-cell apoptosis via STAT-3-mediated nitric oxide production.  
658 *Diabetes/Metabolism Research and Reviews* **27**:813-819.
- 659 32. **Okuda, Y., S. Sakoda, C. Bernard, H. Fujimura, Y. Saeki, T. Kishimoto, and**  
660 **T. Yanagihara.** 1998. IL-6-deficient mice are resistant to the induction of  
661 experimental autoimmune encephalomyelitis provoked by myelin oligodendrocyte  
662 glycoprotein. *Int. Immunol.* **10**:703-708.
- 663 33. **Oleszak, E. L., J. R. Chang, H. Friedman, C. D. Katsetos, and C. D.**  
664 **Platsoucas.** 2004. Theiler's virus infection: a model for multiple sclerosis. *Clin*  
665 *Microbiol Rev* **17**:174-207.
- 666 34. **Palma, J. P., D. Kwon, N. A. Clipstone, and B. S. Kim.** 2003. Infection with  
667 Theiler's Murine Encephalomyelitis Virus Directly Induces Proinflammatory  
668 Cytokines in Primary Astrocytes via NF- $\kappa$ B Activation: Potential Role for  
669 the Initiation of Demyelinating Disease. *J. Virol.* **77**:6322-6331.
- 670 35. **Pavelko, K. D., C. L. Howe, K. M. Drescher, J. D. Gamez, A. J. Johnson, T.**  
671 **Wei, R. M. Ransohoff, and M. Rodriguez.** 2003. Interleukin-6 Protects Anterior  
672 Horn Neurons from Lethal Virus-Induced Injury. *J Neurosci* **23**:481-492.

- 673 36. **Persson, E., O. S. Voznesensky, Y. F. Huang, and U. H. Lerner.** 2005.  
 674 Increased expression of interleukin-6 by vasoactive intestinal peptide is  
 675 associated with regulation of CREB, AP-1 and C/EBP, but not NF-kappaB, in  
 676 mouse calvarial osteoblasts. *Bone* **37**:513-29.
- 677 37. **Petro, T. M.** 2005. Disparate expression of IL-12 by SJL/J and B10.S  
 678 macrophages during Theiler's virus infection is associated with activity of TLR7  
 679 and mitogen-activated protein kinases. *Microbes Infect* **7**:224-32.
- 680 38. **Petro, T. M.** 2005. ERK-MAP-kinases differentially regulate expression of IL-23  
 681 p19 compared with p40 and IFN-beta in Theiler's virus-infected RAW264.7 cells.  
 682 *Immunol Lett* **97**:47-53.
- 683 39. **Rodriguez, M., K. D. Pavelko, C. W. McKinney, and J. L. Leibowitz.** 1994.  
 684 Recombinant human IL-6 suppresses demyelination in a viral model of multiple  
 685 sclerosis. *J Immunol* **153**:3811-3821.
- 686 40. **Samoilova, E. B., J. L. Horton, B. Hilliard, T.-S. T. Liu, and Y. Chen.** 1998. IL-  
 687 6-Deficient Mice Are Resistant to Experimental Autoimmune Encephalomyelitis:  
 688 Roles of IL-6 in the Activation and Differentiation of Autoreactive T Cells. *J*  
 689 *Immunol* **161**:6480-6486.
- 690 41. **Sanders, S. P., E. S. Siekierski, J. D. Porter, S. M. Richards, and D. Proud.**  
 691 1998. Nitric Oxide Inhibits Rhinovirus-Induced Cytokine Production and Viral  
 692 Replication in a Human Respiratory Epithelial Cell Line. *J Virol* **72**:934-942.
- 693 42. **Sawada, T., L. A. Falk, P. Rao, W. J. Murphy, and D. H. Pluznik.** 1997. IL-6  
 694 induction of protein-DNA complexes via a novel regulatory region of the inducible

- 695 nitric oxide synthase gene promoter: role of octamer binding proteins. *J Immunol*  
 696 **158**:5267-76.
- 697 43. **Schobitz, B., E. R. de Kloet, W. Sutanto, and F. Holsboer.** 1993. Cellular  
 698 localization of interleukin 6 mRNA and interleukin 6 receptor mRNA in rat brain.  
 699 *Eur J Neurosci* **5**:1426-35.
- 700 44. **Sehgal, P. B., D. C. Helfgott, U. Santhanam, S. B. Tatter, R. H. Clarick, J.**  
 701 **Ghrayeb, and L. T. May.** 1988. Regulation of the acute phase and immune  
 702 responses in viral disease. Enhanced expression of the beta 2-  
 703 interferon/hepatocyte-stimulating factor/interleukin 6 gene in virus-infected  
 704 human fibroblasts. *J Exp Med* **167**:1951-1956.
- 705 45. **Shaywitz, A. J., and M. E. Greenberg.** 1999. CREB: a stimulus-induced  
 706 transcription factor activated by a diverse array of extracellular signals. *Annu Rev*  
 707 *Biochem* **68**:821-61.
- 708 46. **Shresta, S., K. L. Sharar, D. M. Prigozhin, H. M. Snider, P. R. Beatty, and E.**  
 709 **Harris.** 2005. Critical Roles for Both STAT1-Dependent and STAT1-Independent  
 710 Pathways in the Control of Primary Dengue Virus Infection in Mice. *J Immunol*  
 711 **175**:3946-3954.
- 712 47. **Spooren, A., K. Kolmus, G. Laureys, R. Clinckers, J. De Keyser, G.**  
 713 **Haegeman, and S. Gerlo.** 2011. Interleukin-6, a mental cytokine. *Brain Res Rev*  
 714 **67**:157-183.
- 715 48. **Starnes, H. F., Jr., M. K. Pearce, A. Tewari, J. H. Yim, J. C. Zou, and J. S.**  
 716 **Abrams.** 1990. Anti-IL-6 monoclonal antibodies protect against lethal



717 Escherichia coli infection and lethal tumor necrosis factor-alpha challenge in  
 718 mice. J Immunol **145**:4185-91.

719 49. **Tuyt, L. M., W. H. Dokter, K. Birkenkamp, S. B. Koopmans, C. Lummen, W.**  
 720 **Kruijer, and E. Vellenga.** 1999. Extracellular-regulated kinase 1/2, Jun N-  
 721 terminal kinase, and c-Jun are involved in NF-kappa B-dependent IL-6  
 722 expression in human monocytes. J Immunol **162**:4893-902.

723 50. **Xu, Z., S.-B. Xiao, P. Xu, Q. Xie, L. Cao, D. Wang, R. Luo, Y. Zhong, H.-C.**  
 724 **Chen, and L.-R. Fang.** 2011. miR-365, a Novel Negative Regulator of  
 725 Interleukin-6 Gene Expression, Is Cooperatively Regulated by Sp1 and NF-kB. J  
 726 Biol Chem **286**:21401-21412.

727 51. **Zanassi, P., M. Paolillo, A. Feliciello, E. V. Avvedimento, V. Gallo, and S.**  
 728 **Schinelli.** 2001. cAMP-dependent protein kinase induces cAMP-response  
 729 element-binding protein phosphorylation via an intracellular calcium  
 730 release/ERK-dependent pathway in striatal neurons. J Biol Chem **276**:11487-95.

731 52. **Zhang, P.-L., M. Izrael, E. Ainbinder, L. Ben-Simchon, J. Chebath, and M.**  
 732 **Revel.** 2006. Increased myelinating capacity of embryonic stem cell derived  
 733 oligodendrocyte precursors after treatment by interleukin-6/soluble interleukin-6  
 734 receptor fusion protein. Mol Cell Neurosci **31**:387-398.

735  
 736  
 737  
 738

739 **FIG 1** TMEV induces more IL-6 expression in macrophages. IL-6 mRNA or protein  
740 secretion was induced from SJL/J or B10.S macrophages 3 and 8 h after infection with  
741 TMEV. SJL/J or B10.S macrophages ( $1 \times 10^6$ ) were infected with  $2-10 \times 10^5$  PFU of  
742 TMEV in the absence (A, B), or presence of 10 ng/ml recombinant (rec) IL-6 (C, E),  
743 0.01–10 ng/ml rec IL-6 (D), 3  $\mu$ g/ml anti-IL-6-Fab (E), or SJL/J macrophages were  
744 treated with 10 ng/ml rec IL-6 which was removed after 30 min (D), or SJLJ  
745 macrophages were treated with 10 ng/ml rec IL-6 or rec IFN $\beta$  starting at 1, 3, 6, or 7 h  
746 PI (F). RNA was reverse transcribed and relative levels of IL-6 mRNA (A) was evaluated  
747 by real-time PCR, IL-6 protein by ELISA (B), and TMEV by real-time PCR (C, D, E,F).  
748 Data are means of 3-5 samples per time point evaluated by the Student *t* test; \*  
749 indicates  $p \leq 0.05$ , \*\*  $p \leq 0.01$ .

750

751 **FIG 2** ERK MAPK activation is required for optimum TMEV-induced IL-6 expression. 1  
752  $\times 10^6$  macrophages were untreated (nil or control) or pretreated with 20  $\mu$ M SB203580  
753 (SB) or 40  $\mu$ M U0126 (U) for 30 min before infection with  $2 \times 10^5$  PFU of TMEV. After 24  
754 h, relative levels of IL-6 mRNA was evaluated by real-time PCR (A), IL-6 protein  
755 secretion evaluated by ELISA (B), and TMEV RNA by real-time PCR (C). Data are  
756 means of 5 samples from two experiments evaluated by the Student *t* test. \* with  
757 brackets indicates  $p \leq 0.05$ .

758

759

760 **FIG 3** Optimum IL-6 expression in B10.S compared with SJL/J macrophages following  
761 TLR4 and TLR3 stimulation depends on ERK MAPK activation.  $1 \times 10^6$  macrophages  
762 were untreated (control) or pretreated with U0126 (U) for 30 min before stimulation with  
763  $1 \mu\text{g/ml}$  LPS (A, B) or  $50 \mu\text{g/ml}$  polyI:C (C, D). After 24 h (A, C), relative levels of IL-6  
764 mRNA was evaluated by real-time PCR and (B, D) IL-6 protein secretion evaluated by  
765 ELISA. Data are means of 5 samples evaluated by the Student *t* test. \* indicates  
766 comparisons considered significantly different;  $p \leq 0.05$ .

767

768 **FIG 4** IL-12 and IFN $\beta$  enhance IL-6 expression and decrease TMEV replication during  
769 TMEV infection of B10.S and SJL/J macrophages.  $1 \times 10^6$  macrophages from SJL/J and  
770 B10.S mice were untreated (control) or treated with IL-12 p70 (1 ng/ml), IFN $\beta$  (10  
771 ng/ml), or IL-12 p40 (10 ng/ml) 45 min before and during infection with  $2 \times 10^5$  PFU of  
772 TMEV. After 24 and 48 h of infection relative levels of IL-6 mRNA was evaluated by  
773 real-time PCR (A), IL-6 protein was evaluated by ELISA (B), and TMEV RNA was  
774 evaluated by real-time PCR (C). Data are means of 3 samples each evaluated by the  
775 Student *t* test. \* indicates comparisons considered significantly different;  $p \leq 0.05$ .

776

777

778

779

780 **FIG 5** IL-12 and ERK-MAPK contribute to IL-6 expression during TMEV infection of  
781 RAW264.7 macrophages.  $1 \times 10^6$  RAW264.7 macrophage cells were untreated (control)  
782 or treated with IL-12 p70 (1 ng/ml) with or without 40  $\mu$ M U0126 ERK MAPK inhibitor 45  
783 min before and during infection with  $1 \times 10^6$  PFU of TMEV. After 24 h of infection  
784 relative levels of IL-6 mRNA (A) was evaluated by real-time PCR and (B) IL-6 protein  
785 was evaluated by ELISA. Data are means of 3 samples each evaluated by the Student *t*  
786 test. This experiment was repeated three times. \* indicates  $p \leq 0.05$ , \*\*  $p \leq 0.01$ , n.d.=  
787 not detectable.

788

789 **FIG 6** IL-6 constrains TMEV replication during infection of RAW264.7 macrophages.  $1$   
790  $\times 10^6$  RAW264.7 macrophage cells were untreated (control) or treated with 10ng/ml IL-6  
791 (A, C) or 0.01- 10 ng/ml IL-6 (B) during infection with  $1 \times 10^6$  PFU of TMEV. After 3, 6  
792 (D), or 24 h (A, B, C) of infection relative levels of TMEV (A, B, D) or IL-6 (C) mRNA  
793 were evaluated by real-time PCR. Means are of 6 samples from two experiments each  
794 evaluated by the Student *t* test. \* indicates  $p \leq 0.05$ , \*\*  $p \leq 0.01$ .

795

796 **FIG 7** IL-6 activates STAT1 and STAT3 in macrophages. (A) Western blots of  
797 phospho-STAT1, phospho-STAT3, total STAT3, and beta-tubulin at 30 min and 6 h in  
798 untreated or RAW264.7 cells treated with IL-6 or IFN $\beta$  with or without TMEV infection.  
799 (B) TMEV RNA measured by qRT-PCR in RAW264.7 cells transfected with siSTAT1-  
800 RNA (3 pmol) or siSTAT3-RNA (3 pmol) and then infected with TMEV at 1 MOI after 36

801 h. Real-time PCR of IRF1 (B), IRF7 (C), IRF9 (D), and IFN $\beta$  (E) in RAW cells treated  
802 with 10 ng/ml IL-6 prior to and during TMEV infection after 3 and 6 h infection. Data are  
803 means of 5 samples each evaluated by the Student *t* test. \* indicates means  
804 considered significantly different;  $p \leq 0.05$ ; n.d.= not detectable.

805

806

807

808

809

810

811

812

813

814

815

816


817

818


819 Figure 1. Moore et al.

820


821 **A**




**B**




**C**




**D**



**E**




**F**




822 Figure 2. Moore et al.

823



824 Figure 3. Moore et al.


825






826 Figure 4. Moore et al.

827



828 Figure 5. Moore et al.


829



830

Figure 6. Moore et al.

831



832 Figure 7. Moore et al.

