

March 2008

Measurement of the Shape of the Boson- Transverse Momentum Distribution in $pp \rightarrow Z/\gamma^*$ $= \rightarrow e^+e^- + X$ Events Produced at $\sqrt{s} = 1.96$ TeV

V.M. Abazov

Joint Institute for Nuclear Research, Dubna, Russia

Kenneth A. Bloom

University of Nebraska - Lincoln, kbloom2@unl.edu

Gregory Snow

University of Nebraska - Lincoln, gsnow1@unl.edu

D0 Collaboration

Follow this and additional works at: <http://digitalcommons.unl.edu/physicsbloom>



Part of the [Physics Commons](#)

Abazov, V. M.; Bloom, Kenneth A.; Snow, Gregory; and Collaboration, D0, "Measurement of the Shape of the Boson-Transverse Momentum Distribution in $pp \rightarrow Z/\gamma^* \rightarrow e^+e^- + X$ Events Produced at $\sqrt{s} = 1.96$ TeV" (2008). *Kenneth Bloom Publications*. 230. <http://digitalcommons.unl.edu/physicsbloom/230>

This Article is brought to you for free and open access by the Research Papers in Physics and Astronomy at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Kenneth Bloom Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Measurement of the Shape of the Boson-Transverse Momentum Distribution in $p\bar{p} \rightarrow Z/\gamma^* \rightarrow e^+e^- + X$ Events Produced at $\sqrt{s} = 1.96$ TeV

V. M. Abazov,³⁶ B. Abbott,⁷⁶ M. Abolins,⁶⁶ B. S. Acharya,²⁹ M. Adams,⁵² T. Adams,⁵⁰ E. Aguilo,⁶ S. H. Ahn,³¹ M. Ahsan,⁶⁰ G. D. Alexeev,³⁶ G. Alkhazov,⁴⁰ A. Alton,^{65,*} G. Alverson,⁶⁴ G. A. Alves,² M. Anastasoae,³⁵ L. S. Ancu,³⁵ T. Andeen,⁵⁴ S. Anderson,⁴⁶ B. Andrieu,¹⁷ M. S. Anzels,⁵⁴ Y. Arnaud,¹⁴ M. Arov,⁶¹ M. Arthaud,¹⁸ A. Askew,⁵⁰ B. Åsman,⁴¹ A. C. S. Assis Jesus,³ O. Atramentov,⁵⁰ C. Autermann,²¹ C. Avila,⁸ C. Ay,²⁴ F. Badaud,¹³ A. Baden,⁶² L. Bagby,⁵³ B. Baldin,⁵¹ D. V. Bandurin,⁶⁰ S. Banerjee,²⁹ P. Banerjee,²⁹ E. Barberis,⁶⁴ A.-F. Barfuss,¹⁵ P. Bargassa,⁸¹ P. Baringer,⁵⁹ J. Barreto,² J. F. Bartlett,⁵¹ U. Bassler,¹⁸ D. Bauer,⁴⁴ S. Beale,⁶ A. Bean,⁵⁹ M. Begalli,³ M. Begel,⁷² C. Belanger-Champagne,⁴¹ L. Bellantoni,⁵¹ A. Bellavance,⁵¹ J. A. Benitez,⁶⁶ S. B. Beri,²⁷ G. Bernardi,¹⁷ R. Bernhard,²³ I. Bertram,⁴³ M. Besançon,¹⁸ R. Beuselinck,⁴⁴ V. A. Bezzubov,³⁹ P. C. Bhat,⁵¹ V. Bhatnagar,²⁷ C. Biscarat,²⁰ G. Blazey,⁵³ F. Blekman,⁴⁴ S. Blessing,⁵⁰ D. Bloch,¹⁹ K. Bloom,⁶⁸ A. Boehnlein,⁵¹ D. Boline,⁶³ T. A. Bolton,⁶⁰ G. Borissov,⁴³ T. Bose,⁷⁸ A. Brandt,⁷⁹ R. Brock,⁶⁶ G. Brooijmans,⁷¹ A. Bross,⁵¹ D. Brown,⁸² N. J. Buchanan,⁵⁰ D. Buchholz,⁵⁴ M. Buehler,⁸² V. Buescher,²² V. Bunichev,³⁸ S. Burdin,^{43,†} S. Burke,⁴⁶ T. H. Burnett,⁸³ C. P. Buszello,⁴⁴ J. M. Butler,⁶³ P. Calfayan,²⁵ S. Calvet,¹⁶ J. Cammin,⁷² W. Carvalho,³ B. C. K. Casey,⁵¹ N. M. Cason,⁵⁶ H. Castilla-Valdez,³³ S. Chakrabarti,¹⁸ D. Chakraborty,⁵³ K. M. Chan,⁵⁶ K. Chan,⁶ A. Chandra,⁴⁹ F. Charles,^{19,**} E. Cheu,⁴⁶ F. Chevallier,¹⁴ D. K. Cho,⁶³ S. Choi,³² B. Choudhary,²⁸ L. Christofek,⁷⁸ T. Christoudias,⁴⁴ S. Cihangir,⁵¹ D. Claes,⁶⁸ Y. Coadou,⁶ M. Cooke,⁸¹ W. E. Cooper,⁵¹ M. Corcoran,⁸¹ F. Couderc,¹⁸ M.-C. Cousinou,¹⁵ S. Crépe-Renaudin,¹⁴ D. Cutts,⁷⁸ M. Ćwiok,³⁰ H. da Motta,² A. Das,⁴⁶ G. Davies,⁴⁴ K. De,⁷⁹ S. J. de Jong,³⁵ E. De La Cruz-Burelo,⁶⁵ C. De Oliveira Martins,³ J. D. Degenhardt,⁶⁵ F. Déliot,¹⁸ M. Demarteau,⁵¹ R. Demina,⁷² D. Denisov,⁵¹ S. P. Denisov,³⁹ S. Desai,⁵¹ H. T. Diehl,⁵¹ M. Diesburg,⁵¹ A. Dominguez,⁶⁸ H. Dong,⁷³ L. V. Dudko,³⁸ L. Dufлот,¹⁶ S. R. Dugad,²⁹ D. Duggan,⁵⁰ A. Duperrin,¹⁵ J. Dyer,⁶⁶ A. Dyshkant,⁵³ M. Eads,⁶⁸ D. Edmunds,⁶⁶ J. Ellison,⁴⁹ V. D. Elvira,⁵¹ Y. Enari,⁷⁸ S. Eno,⁶² P. Ermolov,³⁸ H. Evans,⁵⁵ A. Evdokimov,⁷⁴ V. N. Evdokimov,³⁹ A. V. Ferapontov,⁶⁰ T. Ferbel,⁷² F. Fiedler,²⁴ F. Filthaut,³⁵ W. Fisher,⁵¹ H. E. Fisk,⁵¹ M. Ford,⁴⁵ M. Fortner,⁵³ H. Fox,²³ S. Fu,⁵¹ S. Fuess,⁵¹ T. Gadfort,⁸³ C. F. Galea,³⁵ E. Gallas,⁵¹ E. Galyaev,⁵⁶ C. Garcia,⁷² A. Garcia-Bellido,⁸³ V. Gavrilov,³⁷ P. Gay,¹³ W. Geist,¹⁹ D. Gelé,¹⁹ C. E. Gerber,⁵² Y. Gershtein,⁵⁰ D. Gillberg,⁶ G. Ginter,⁷² N. Gollub,⁴¹ B. Gómez,⁸ A. Goussiou,⁵⁶ P. D. Grannis,⁷³ H. Greenlee,⁵¹ Z. D. Greenwood,⁶¹ E. M. Gregores,⁴ G. Grenier,²⁰ Ph. Gris,¹³ J.-F. Grivaz,¹⁶ A. Grohsjean,²⁵ S. Grünendahl,⁵¹ M. W. Grünewald,³⁰ J. Guo,⁷³ F. Guo,⁷³ P. Gutierrez,⁷⁶ G. Gutierrez,⁵¹ A. Haas,⁷¹ N. J. Hadley,⁶² P. Haefner,²⁵ S. Hagopian,⁵⁰ J. Haley,⁶⁹ I. Hall,⁶⁶ R. E. Hall,⁴⁸ L. Han,⁷ K. Hanagaki,⁵¹ P. Hansson,⁴¹ K. Harder,⁴⁵ A. Harel,⁷² R. Harrington,⁶⁴ J. M. Hauptman,⁵⁸ R. Hauser,⁶⁶ J. Hays,⁴⁴ T. Hebbeker,²¹ D. Hedin,⁵³ J. G. Hegeman,³⁴ J. M. Heinmiller,⁵² A. P. Heinson,⁴⁹ U. Heintz,⁶³ C. Hensel,⁵⁹ K. Herner,⁷³ G. Hesketh,⁶⁴ M. D. Hildreth,⁵⁶ R. Hirosky,⁸² J. D. Hobbs,⁷³ B. Hoeneisen,¹² H. Hoeth,²⁶ M. Hohlfeld,²² S. J. Hong,³¹ S. Hossain,⁷⁶ P. Houben,³⁴ Y. Hu,⁷³ Z. Hubacek,¹⁰ V. Hynek,⁹ I. Iashvili,⁷⁰ R. Illingworth,⁵¹ A. S. Ito,⁵¹ S. Jabeen,⁶³ M. Jaffré,¹⁶ S. Jain,⁷⁶ K. Jakobs,²³ C. Jarvis,⁶² R. Jesik,⁴⁴ K. Johns,⁴⁶ C. Johnson,⁷¹ M. Johnson,⁵¹ A. Jonckheere,⁵¹ P. Jonsson,⁴⁴ A. Juste,⁵¹ D. Käfer,²¹ E. Kajfasz,¹⁵ A. M. Kalinin,³⁶ J. R. Kalk,⁶⁶ J. M. Kalk,⁶¹ S. Kappler,²¹ D. Karmanov,³⁸ P. Kasper,⁵¹ I. Katsanos,⁷¹ D. Kau,⁵⁰ R. Kaur,²⁷ V. Kaushik,⁷⁹ R. Kehoe,⁸⁰ S. Kermiche,¹⁵ N. Khalatyan,⁵¹ A. Khanov,⁷⁷ A. Kharchilava,⁷⁰ Y. M. Kharzhev,³⁶ D. Khatidze,⁷¹ H. Kim,³² T. J. Kim,³¹ M. H. Kirby,⁵⁴ M. Kirsch,²¹ B. Klima,⁵¹ J. M. Kohli,²⁷ J.-P. Konrath,²³ M. Kopal,⁷⁶ V. M. Korablev,³⁹ A. V. Kozelov,³⁹ D. Krop,⁵⁵ T. Kuhl,²⁴ A. Kumar,⁷⁰ S. Kunori,⁶² A. Kupco,¹¹ T. Kurča,²⁰ J. Kvita,⁹ F. Lacroix,¹³ D. Lam,⁵⁶ S. Lammers,⁷¹ G. Landsberg,⁷⁸ P. Lebrun,²⁰ W. M. Lee,⁵¹ A. Leflat,³⁸ F. Lehner,⁴² J. Lellouch,¹⁷ J. Leveque,⁴⁶ P. Lewis,⁴⁴ J. Li,⁷⁹ Q. Z. Li,⁵¹ L. Li,⁴⁹ S. M. Lietti,⁵ J. G. R. Lima,⁵³ D. Lincoln,⁵¹ J. Linnemann,⁶⁶ V. V. Lipaev,³⁹ R. Lipton,⁵¹ Y. Liu,⁷ Z. Liu,⁶ L. Lobo,⁴⁴ A. Lobodenko,⁴⁰ M. Lokajicek,¹¹ P. Love,⁴³ H. J. Lubatti,⁸³ A. L. Lyon,⁵¹ A. K. A. Maciel,² D. Mackin,⁸¹ R. J. Madaras,⁴⁷ P. Mättig,²⁶ C. Magass,²¹ A. Magerkurth,⁶⁵ P. K. Mal,⁵⁶ H. B. Malbouisson,³ S. Malik,⁶⁸ V. L. Malyshev,³⁶ H. S. Mao,⁵¹ Y. Maravin,⁶⁰ B. Martin,¹⁴ R. McCarthy,⁷³ A. Melnitchouk,⁶⁷ A. Mendes,¹⁵ L. Mendoza,⁸ P. G. Mercadante,⁵ M. Merkin,³⁸ K. W. Merritt,⁵¹ J. Meyer,^{22,§} A. Meyer,²¹ T. Millet,²⁰ J. Mitrevski,⁷¹ J. Molina,³ R. K. Mommsen,⁴⁵ N. K. Mondal,²⁹ R. W. Moore,⁶ T. Moulik,⁵⁹ G. S. Muanza,²⁰ M. Mulders,⁵¹ M. Mulhearn,⁷¹ O. Mundal,²² L. Mundim,³ E. Nagy,¹⁵ M. Naimuddin,⁵¹ M. Narain,⁷⁸ N. A. Naumann,³⁵ H. A. Neal,⁶⁵ J. P. Negret,⁸ P. Neustroev,⁴⁰ H. Nilsen,²³ H. Nogima,³ A. Nomerotski,⁵¹ S. F. Novaes,⁵ T. Nunnemann,²⁵ V. O'Dell,⁵¹ D. C. O'Neil,⁶ G. Obrant,⁴⁰ C. Ochando,¹⁶ D. Onoprienko,⁶⁰ N. Oshima,⁵¹ J. Osta,⁵⁶ R. Otec,¹⁰ G. J. Otero y Garzón,⁵¹ M. Owen,⁴⁵ P. Padley,⁸¹ M. Pangilinan,⁷⁸ N. Parashar,⁵⁷ S.-J. Park,⁷² S. K. Park,³¹ J. Parsons,⁷¹ R. Partridge,⁷⁸ N. Parua,⁵⁵ A. Patwa,⁷⁴ G. Pawloski,⁸¹ B. Penning,²³ M. Perfilov,³⁸ K. Peters,⁴⁵ Y. Peters,²⁶ P. Pétröf,¹⁶

M. Petteni,⁴⁴ R. Piegaia,¹ J. Piper,⁶⁶ M.-A. Pleier,²² P.L. M. Podesta-Lerma,^{33,‡} V.M. Podstavkov,⁵¹ Y. Pogorelov,⁵⁶ M.-E. Pol,² P. Polozov,³⁷ B. G. Pope,⁶⁶ A. V. Popov,³⁹ C. Potter,⁶ W. L. Prado da Silva,³ H. B. Prosper,⁵⁰ S. Protopopescu,⁷⁴ J. Qian,⁶⁵ A. Quadt,^{22,§} B. Quinn,⁶⁷ A. Rakitine,⁴³ M. S. Rangel,² K. Ranjan,²⁸ P. N. Ratoff,⁴³ P. Renkel,⁸⁰ S. Reucroft,⁶⁴ P. Rich,⁴⁵ M. Rijssenbeek,⁷³ I. Ripp-Baudot,¹⁹ F. Rizatdinova,⁷⁷ S. Robinson,⁴⁴ R. F. Rodrigues,³ M. Rominsky,⁷⁶ C. Royon,¹⁸ P. Rubinov,⁵¹ R. Ruchti,⁵⁶ G. Safronov,³⁷ G. Sajot,¹⁴ A. Sánchez-Hernández,³³ M. P. Sanders,¹⁷ A. Santoro,³ G. Savage,⁵¹ L. Sawyer,⁶¹ T. Scanlon,⁴⁴ D. Schaile,²⁵ R. D. Schamberger,⁷³ Y. Scheglov,⁴⁰ H. Schellman,⁵⁴ P. Schieferdecker,²⁵ T. Schliephake,²⁶ C. Schwanenberger,⁴⁵ A. Schwartzman,⁶⁹ R. Schwienhorst,⁶⁶ J. Sekaric,⁵⁰ H. Severini,⁷⁶ E. Shabalina,⁵² M. Shamim,⁶⁰ V. Shary,¹⁸ A. A. Shchukin,³⁹ R. K. Shivpuri,²⁸ V. Siccaldi,¹⁹ V. Simak,¹⁰ V. Sirotenko,⁵¹ P. Skubic,⁷⁶ P. Slattery,⁷² D. Smirnov,⁵⁶ J. Snow,⁷⁵ G. R. Snow,⁶⁸ S. Snyder,⁷⁴ S. Söldner-Rembold,⁴⁵ L. Sonnenschein,¹⁷ A. Sopczak,⁴³ M. Sosebee,⁷⁹ K. Soustruznik,⁹ M. Souza,² B. Spurlock,⁷⁹ J. Stark,¹⁴ J. Steele,⁶¹ V. Stolin,³⁷ D. A. Stoyanova,³⁹ J. Strandberg,⁶⁵ S. Strandberg,⁴¹ M. A. Strang,⁷⁰ M. Strauss,⁷⁶ E. Strauss,⁷³ R. Ströhmer,²⁵ D. Strom,⁵⁴ L. Stutte,⁵¹ S. Sumowidagdo,⁵⁰ P. Svoisky,⁵⁶ A. Sznajder,³ M. Talby,¹⁵ P. Tamburello,⁴⁶ A. Tanasijczuk,¹ W. Taylor,⁶ J. Temple,⁴⁶ B. Tiller,²⁵ F. Tissandier,¹³ M. Titov,¹⁸ V. V. Tokmenin,³⁶ T. Toole,⁶² I. Torchiani,²³ T. Trefzger,²⁴ D. Tsybychev,⁷³ B. Tuchming,¹⁸ C. Tully,⁶⁹ P. M. Tuts,⁷¹ R. Unalan,⁶⁶ S. Uvarov,⁴⁰ L. Uvarov,⁴⁰ S. Uzunyan,⁵³ B. Vachon,⁶ P. J. van den Berg,³⁴ R. Van Kooten,⁵⁵ W. M. van Leeuwen,³⁴ N. Varelas,⁵² E. W. Varnes,⁴⁶ I. A. Vasilyev,³⁹ M. Vaupel,²⁶ P. Verdier,²⁰ L. S. Vertogradov,³⁶ M. Verzocchi,⁵¹ F. Villeneuve-Seguiier,⁴⁴ P. Vint,⁴⁴ P. Vokac,¹⁰ E. Von Toerne,⁶⁰ M. Voutilainen,^{68,||} R. Wagner,⁶⁹ H. D. Wahl,⁵⁰ L. Wang,⁶² M. H. L. S Wang,⁵¹ J. Warchol,⁵⁶ G. Watts,⁸³ M. Wayne,⁵⁶ M. Weber,⁵¹ G. Weber,²⁴ A. Wenger,^{23,¶} N. Wermes,²² M. Wetstein,⁶² A. White,⁷⁹ D. Wicke,²⁶ G. W. Wilson,⁵⁹ S. J. Wimpenny,⁴⁹ M. Wobisch,⁶¹ D. R. Wood,⁶⁴ T. R. Wyatt,⁴⁵ Y. Xie,⁷⁸ S. Yacoub,⁵⁴ R. Yamada,⁵¹ M. Yan,⁶² T. Yasuda,⁵¹ Y. A. Yatsunenko,³⁶ K. Yip,⁷⁴ H. D. Yoo,⁷⁸ S. W. Youn,⁵⁴ J. Yu,⁷⁹ A. Zatserklyaniy,⁵³ C. Zeitnitz,²⁶ T. Zhao,⁸³ B. Zhou,⁶⁵ J. Zhu,⁷³ M. Zielinski,⁷² D. Zieminska,⁵⁵ A. Zieminski,^{55,*} L. Zivkovic,⁷¹ V. Zutshi,⁵³ and E. G. Zverev³⁸

(The D0 Collaboration)

¹Universidad de Buenos Aires, Buenos Aires, Argentina

²LAFEX, Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil

³Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

⁴Universidade Federal do ABC, Santo André, Brazil

⁵Instituto de Física Teórica, Universidade Estadual Paulista, São Paulo, Brazil

⁶University of Alberta, Edmonton, Alberta, Canada,

Simon Fraser University, Burnaby, British Columbia, Canada,

York University, Toronto, Ontario, Canada,

and McGill University, Montreal, Quebec, Canada

⁷University of Science and Technology of China, Hefei, People's Republic of China

⁸Universidad de los Andes, Bogotá, Colombia

⁹Center for Particle Physics, Charles University, Prague, Czech Republic

¹⁰Czech Technical University, Prague, Czech Republic

¹¹Center for Particle Physics, Institute of Physics, Academy of Sciences of the Czech Republic, Prague, Czech Republic

¹²Universidad San Francisco de Quito, Quito, Ecuador

¹³Laboratoire de Physique Corpusculaire, IN2P3-CNRS, Université Blaise Pascal, Clermont-Ferrand, France

¹⁴Laboratoire de Physique Subatomique et de Cosmologie, IN2P3-CNRS, Université de Grenoble I, Grenoble, France

¹⁵CPPM, IN2P3-CNRS, Université de la Méditerranée, Marseille, France

¹⁶Laboratoire de l'Accélérateur Linéaire, IN2P3-CNRS et Université Paris-Sud, Orsay, France

¹⁷LPNHE, IN2P3-CNRS, Universités Paris VI and VII, Paris, France

¹⁸DAPNIA/Service de Physique des Particules, CEA, Saclay, France

¹⁹IPHC, Université Louis Pasteur et Université de Haute Alsace, CNRS, IN2P3, Strasbourg, France

²⁰IPNL, Université Lyon 1, CNRS/IN2P3, Villeurbanne, France and Université de Lyon, Lyon, France

²¹III. Physikalisches Institut A, RWTH Aachen, Aachen, Germany

²²Physikalisches Institut, Universität Bonn, Bonn, Germany

²³Physikalisches Institut, Universität Freiburg, Freiburg, Germany

²⁴Institut für Physik, Universität Mainz, Mainz, Germany

²⁵Ludwig-Maximilians-Universität München, München, Germany

²⁶Fachbereich Physik, University of Wuppertal, Wuppertal, Germany

²⁷Panjab University, Chandigarh, India

²⁸Delhi University, Delhi, India

²⁹Tata Institute of Fundamental Research, Mumbai, India

- ³⁰University College Dublin, Dublin, Ireland
³¹Korea Detector Laboratory, Korea University, Seoul, Korea
³²SungKyunKwan University, Suwon, Korea
³³CINVESTAV, Mexico City, Mexico
³⁴FOM-Institute NIKHEF and University of Amsterdam/NIKHEF, Amsterdam, The Netherlands
³⁵Radboud University Nijmegen/NIKHEF, Nijmegen, The Netherlands
³⁶Joint Institute for Nuclear Research, Dubna, Russia
³⁷Institute for Theoretical and Experimental Physics, Moscow, Russia
³⁸Moscow State University, Moscow, Russia
³⁹Institute for High Energy Physics, Protvino, Russia
⁴⁰Petersburg Nuclear Physics Institute, St. Petersburg, Russia
⁴¹Lund University, Lund, Sweden, Royal Institute of Technology and Stockholm University, Stockholm, Sweden, and Uppsala University, Uppsala, Sweden
⁴²Physik Institut der Universität Zürich, Zürich, Switzerland
⁴³Lancaster University, Lancaster, United Kingdom
⁴⁴Imperial College, London, United Kingdom
⁴⁵University of Manchester, Manchester, United Kingdom
⁴⁶University of Arizona, Tucson, Arizona 85721, USA
⁴⁷Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA
⁴⁸California State University, Fresno, California 93740, USA
⁴⁹University of California, Riverside, California 92521, USA
⁵⁰Florida State University, Tallahassee, Florida 32306, USA
⁵¹Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA
⁵²University of Illinois at Chicago, Chicago, Illinois 60607, USA
⁵³Northern Illinois University, DeKalb, Illinois 60115, USA
⁵⁴Northwestern University, Evanston, Illinois 60208, USA
⁵⁵Indiana University, Bloomington, Indiana 47405, USA
⁵⁶University of Notre Dame, Notre Dame, Indiana 46556, USA
⁵⁷Purdue University Calumet, Hammond, Indiana 46323, USA
⁵⁸Iowa State University, Ames, Iowa 50011, USA
⁵⁹University of Kansas, Lawrence, Kansas 66045, USA
⁶⁰Kansas State University, Manhattan, Kansas 66506, USA
⁶¹Louisiana Tech University, Ruston, Louisiana 71272, USA
⁶²University of Maryland, College Park, Maryland 20742, USA
⁶³Boston University, Boston, Massachusetts 02215, USA
⁶⁴Northeastern University, Boston, Massachusetts 02115, USA
⁶⁵University of Michigan, Ann Arbor, Michigan 48109, USA
⁶⁶Michigan State University, East Lansing, Michigan 48824, USA
⁶⁷University of Mississippi, University, Mississippi 38677, USA
⁶⁸University of Nebraska, Lincoln, Nebraska 68588, USA
⁶⁹Princeton University, Princeton, New Jersey 08544, USA
⁷⁰State University of New York, Buffalo, New York 14260, USA
⁷¹Columbia University, New York, New York 10027, USA
⁷²University of Rochester, Rochester, New York 14627, USA
⁷³State University of New York, Stony Brook, New York 11794, USA
⁷⁴Brookhaven National Laboratory, Upton, New York 11973, USA
⁷⁵Langston University, Langston, Oklahoma 73050, USA
⁷⁶University of Oklahoma, Norman, Oklahoma 73019, USA
⁷⁷Oklahoma State University, Stillwater, Oklahoma 74078, USA
⁷⁸Brown University, Providence, Rhode Island 02912, USA
⁷⁹University of Texas, Arlington, Texas 76019, USA
⁸⁰Southern Methodist University, Dallas, Texas 75275, USA
⁸¹Rice University, Houston, Texas 77005, USA
⁸²University of Virginia, Charlottesville, Virginia 22901, USA
⁸³University of Washington, Seattle, Washington 98195, USA

(Received 5 December 2007; published 11 March 2008)

We present a measurement of the shape of the Z/γ^* boson transverse momentum (q_T) distribution in $p\bar{p} \rightarrow Z/\gamma^* \rightarrow e^+e^- + X$ events at a center-of-mass energy of 1.96 TeV using 0.98 fb⁻¹ of data collected with the D0 detector at the Fermilab Tevatron collider. The data are found to be consistent with the resummation prediction at low q_T , but above the perturbative QCD calculation in the region of $q_T > 30$ GeV/ c .

Using events with $q_T < 30$ GeV/ c , we extract the value of g_2 , one of the nonperturbative parameters for the resummation calculation. Data at large boson rapidity y are compared with the prediction of resummation and with alternative models that employ a resummed form factor with modifications in the small Bjorken x region of the proton wave function.

DOI: [10.1103/PhysRevLett.100.102002](https://doi.org/10.1103/PhysRevLett.100.102002)

PACS numbers: 13.85.-t, 12.38.Qk, 13.38.Dg

A complete understanding of weak vector boson production is essential for maximizing the sensitivity to new physics at hadron colliders. Studies of the Z/γ^* boson production play a particularly valuable role in that its kinematics can be precisely determined through measurement of its leptonic decays. Throughout this Letter, we use the notation “Z boson” to mean “ Z/γ^* boson,” unless specified otherwise.

Z boson production also serves as an ideal testing ground for predictions of quantum chromodynamics (QCD), since the boson’s transverse momentum, q_T , can be measured over a wide range of values and can be correlated with its rapidity. At large q_T (approximately greater than 30 GeV/ c), the radiation of a single parton with large transverse momentum dominates the cross section, and fixed-order perturbative QCD (pQCD) calculations [1], currently available at next-to-next-to leading order (NNLO) [2], should yield reliable predictions. At lower q_T , multiple soft-gluon emission can not be neglected, and the fixed-order perturbation calculation no longer gives accurate results. A soft-gluon resummation technique developed by Collins, Soper, and Sterman (CSS) [3] gives reliable predictions in the low- q_T region. A prescription has been proposed [4] for matching the low- and high- q_T regions in order to provide a continuous prediction for all values of q_T . The CSS resummation formalism allows the inclusion of contributions from large logarithms of the form $\ln^n(q_T^2/Q^2)$ to all orders of perturbation theory in an effective resummed form factor, where Q^2 represents the invariant mass corresponding to the four-momentum transfer. The CSS resummation can be done either in impact parameter (b) space or in transverse momentum (q_T) space. In the case of b -space resummation, this form factor can be parameterized with the following nonperturbative function first introduced by Brock, Landry, Nadolsky, and Yuan (BLNY) [5]:

$$S_{\text{NP}}(b, Q^2) = \left[g_1 + g_2 \ln\left(\frac{Q}{2Q_0}\right) + g_1 g_3 \ln(100x_i x_j) \right] b^2, \quad (1)$$

where x_i and x_j are the fractions of the incident hadron momenta carried by the colliding partons, Q_0 is a scale typical of the onset of nonperturbative effects, and g_1 , g_2 , and g_3 are phenomenological nonperturbative parameters that must be obtained from fits to the data. The Z boson q_T distribution at the Fermilab Tevatron is by far most sensitive to the value of g_2 and quite insensitive to the value of g_3 . Thus, a measurement of the Z boson q_T spectrum can

be used to test this formalism and to determine the value of g_2 .

Recent studies of data from deep inelastic scattering (DIS) experiments [6,7] indicate that the resummed form factor in the above equation may need to be modified for processes involving a small- x parton in the initial state. Reference [8] indicates how such a modification would influence the q_T distributions of vector and Higgs bosons produced in hadronic collisions. A wider q_T distribution is predicted for Z bosons with large rapidity (called “small- x broadening”). Z bosons produced at the Tevatron in the rapidity range $2 < |y| < 3$ probe processes involving a parton with $0.002 < x < 0.006$, and can be used to test the modified form factor at small x .

Z boson q_T distributions have been published previously by the CDF [9] and D0 [10] collaborations using about 100 pb^{-1} of data at $\sqrt{s} = 1.8$ TeV. In this Letter, we report a new measurement with larger statistics and improved precision. This measurement is also the first to present a q_T distribution for large-rapidity Z bosons.

The data sample used in this measurement was collected using a set of inclusive single-electron triggers with the D0 detector [11] at the Fermilab Tevatron collider, and the integrated luminosity is $980 \pm 60 \text{ pb}^{-1}$ [12].

Our selection criteria for Z bosons require two isolated electromagnetic clusters that have a shower shape consistent with that of an electron. Electron candidates are required to have transverse momentum greater than 25 GeV/ c . The electron pairs must have a reconstructed invariant mass $70 < M(ee) < 110 \text{ GeV}/c^2$. If an event has both its candidate electrons in the central calorimeter (CC events), each electron must be spatially matched to a reconstructed track. Because the tracking efficiency decreases with rapidity in the endcap region, events with one or two endcap calorimeter electron candidates (CE and EE events, respectively) are required to have at least one electron with a matching track. After these requirements, 23 959 CC, 30 344 CE, and 9598 EE events are selected; 5412 of these have a Z boson with $|y| > 2$.

Electron identification efficiencies are measured using a combination of data and a GEANT-based [13] simulation of the D0 detector. The electron identification efficiencies are measured from Z data. The dependence of the overall selection efficiency on the Z boson q_T is parameterized from the GEANT simulation. A measurement of this shape from the data agrees well with the simulation within statistical uncertainties.

The dominant backgrounds are from photon plus jet events and di-jet events, with photons and jets misidenti-

fied as electrons. The kinematic properties of these events are obtained from events that satisfy most of the Z selection criteria, but fail the electron shower shape requirement. The normalization of the background is obtained by fitting to a sum of a signal shape obtained from a parameterized simulation of the detector response and the invariant mass distribution from the background sample to the invariant mass distribution of the data sample. The background fractions are $(1.30 \pm 0.14)\%$, $(8.55 \pm 0.26)\%$, and $(4.71 \pm 0.30)\%$ for CC, CE, and EE events, respectively. Other backgrounds are negligible.

The data are corrected for acceptances within a range of generated Z masses of 40 to 200 GeV/c^2 , and for selection efficiencies using a parameterized simulation. We use RESBOS [14] as the event generator which incorporates the resummation calculation in b -space using the BLNY parameterization for low q_T and a NLO pQCD calculation for high q_T . We use PHOTOS [15] to simulate the effects of final state photon radiation. The overall acceptance times efficiency falls slowly from a value of 0.27 at low q_T to a minimum of 0.19 at $q_T = 40 \text{ GeV}/c$ and slowly increases for larger q_T .

The measured spectrum is further corrected for detector resolution effects using the RUN (Regularized Unfolding) program [16] to obtain the true differential cross section. Its performance was verified by comparing the true and unfolded spectrum generated using pseudoexperiments. The measured Z q_T resolution is about 2 GeV/c ; the bin width we choose is 2.5 GeV/c for $q_T < 30 \text{ GeV}/c$. The typical correlation between adjacent bins is around 30%. Due to limited statistics, the chosen bin width is 10 GeV/c for $30 < q_T < 100 \text{ GeV}/c$ and 40 GeV/c for $100 < q_T < 260 \text{ GeV}/c$.

Systematic uncertainties on the unfolded q_T spectrum arise from uncertainties on the electron energy calibration, the electron energy resolution, the dependence of the overall selection efficiency on q_T , and the effect of parton distribution functions (PDFs) on the acceptance. The uncertainties on the unfolded spectrum are estimated from the resulting change when the smearing parameters are varied within their uncertainties. CTEQ 6.1M is used as the default PDF. Uncertainties due to the PDFs are estimated using the procedure described in Ref. [17]. The uncertainty due to the choice of unfolding parameters in the RUN program is also estimated and included in the final systematic uncertainty.

The final results in the $q_T < 30 \text{ GeV}/c$ range are shown in Fig. 1 for the inclusive sample and for the sample with $|y| > 2$. Each data point is plotted at the average value of the expected distribution over the bin [18]. For the theoretical calculation, we use RESBOS with published values of the nonperturbative parameters [5]. Good agreement between data and the prediction is observed for all rapidity ranges, which indicates that the BLNY parameterization works well for the low q_T region.




FIG. 1 (color online). The normalized differential cross section as a function of q_T for (a) the inclusive sample and (b) the sample with Z boson $|y| > 2$ with $q_T < 30 \text{ GeV}/c$. The points are the data, the solid curve is the RESBOS prediction, and the dashed line in (b) is the prediction from the form factor modified after studies of small- x DIS data.

Z boson events produced at large rapidities ($|y| > 2$) are also used to test the small- x prediction. We compare data with the theoretical predictions with and without the form factor as modified from studies of small- x DIS data [8]. All curves are normalized to 1 for $q_T < 30 \text{ GeV}/c$. The default values for the parameters g_1 , g_2 , and g_3 [5] obtained from large- x data are used. The $\chi^2/\text{d.o.f.}$ between the data and the RESBOS calculation using the default parameters is 0.8/1 for $q_T < 5 \text{ GeV}/c$ and 11.1/11 for $q_T < 30 \text{ GeV}/c$, while that for the modified calculation is 5.7/1 for $q_T < 5 \text{ GeV}/c$ and 31.9/11 for $q_T < 30 \text{ GeV}/c$. It remains to be seen if retuning of the nonperturbative parameters could improve the agreement for the modified calculations.

Figure 2 shows the measured differential cross section in the range $q_T < 260 \text{ GeV}/c$ compared to (1) the RESBOS calculation with its default parameters [5], (2) RESBOS with a NLO to NNLO K factor by Arnold and Reno [19] incorporated into RESBOS by its authors, (3) a pQCD




FIG. 2 (color online). The normalized differential cross section as a function of q_T compared to four theoretical calculations for (a) the entire range measured and (b) the fractional differences between data and the theoretical predictions. The four theoretical calculations are RESBOS with its default parameters, RESBOS with a NLO to NNLO K factor by Arnold and Reno, the NNLO calculation by Melnikov and Petriello, and the NNLO calculation but rescaled to data at $q_T = 30$ GeV/ c .

calculation at NNLO [2] using the MRST 2001 NNLO PDF set [20] divided by the NNLO calculation of the total cross section [21], and (4) the NNLO calculation but rescaled to the data at $q_T = 30$ GeV/ c . The agreement between data and RESBOS, with or without the K factor, is good for $q_T < 30$ GeV/ c . At higher values of q_T , the data are not in agreement with the RESBOS calculation. The data agree better with the NNLO calculation and RESBOS prediction with the Arnold-Reno K factor, but agrees best when the NNLO results are rescaled by a factor of 1.25 so that they match the data at $q_T = 30$ GeV/ c . This indicates that the shape from these calculations agrees with the data, and that the source of the discrepancy is in the normalization. Table I summarizes the measured values for each q_T bin together with statistical and systematic uncertainties.

TABLE I. The normalized differential cross section for Z events produced in bins of q_T . The first uncertainty is statistical, and the second is systematic.

$\langle q_T \rangle$ (GeV/ c)	$1/\sigma \times d\sigma/dq_T$ (GeV/ c) $^{-1}$
1.1	$(5.32 \pm 0.13 \pm 0.24) \times 10^{-2}$
4.0	$(8.08 \pm 0.12 \pm 0.19) \times 10^{-2}$
6.2	$(6.33 \pm 0.11 \pm 0.14) \times 10^{-2}$
8.7	$(4.43 \pm 0.09 \pm 0.11) \times 10^{-2}$
11.3	$(3.15 \pm 0.08 \pm 0.08) \times 10^{-2}$
13.7	$(2.46 \pm 0.07 \pm 0.06) \times 10^{-2}$
16.2	$(1.86 \pm 0.06 \pm 0.05) \times 10^{-2}$
18.7	$(1.42 \pm 0.05 \pm 0.05) \times 10^{-2}$
21.3	$(1.09 \pm 0.04 \pm 0.03) \times 10^{-2}$
23.7	$(9.40 \pm 0.40 \pm 0.20) \times 10^{-3}$
26.4	$(6.90 \pm 0.30 \pm 0.20) \times 10^{-3}$
28.5	$(5.50 \pm 0.30 \pm 0.10) \times 10^{-3}$
34.6	$(3.90 \pm 0.10 \pm 0.10) \times 10^{-3}$
44.6	$(2.10 \pm 0.07 \pm 0.06) \times 10^{-3}$
54.6	$(1.10 \pm 0.05 \pm 0.03) \times 10^{-3}$
64.6	$(7.30 \pm 0.40 \pm 0.20) \times 10^{-4}$
73.4	$(4.20 \pm 0.30 \pm 0.20) \times 10^{-4}$
85.4	$(2.50 \pm 0.20 \pm 0.10) \times 10^{-4}$
95.1	$(1.60 \pm 0.17 \pm 0.08) \times 10^{-4}$
117.5	$(6.00 \pm 0.50 \pm 0.30) \times 10^{-5}$
157.5	$(1.10 \pm 0.20 \pm 0.07) \times 10^{-5}$
195.5	$(3.00 \pm 1.00 \pm 0.30) \times 10^{-6}$
245.5	$(7.10 \pm 6.10 \pm 0.60) \times 10^{-7}$

The CSS model parameter most sensitive to the shape at low q_T ($q_T < 30$ GeV/ c) is g_2 . In a fit, we fix other phenomenological parameters to the values obtained in Ref. [5] and only vary g_2 . A minimum $\chi^2/\text{d.o.f.}$ of 9/11 between the model and the inclusive data for $q_T < 30$ GeV/ c is found when $g_2 = 0.77 \pm 0.06$ (GeV/ c) 2 .

In conclusion, we have measured the normalized differential spectrum, $\frac{1}{\sigma} \frac{d\sigma}{dq_T}$, for Z boson events produced in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV with boson mass $40 < M < 200$ GeV/ c^2 and $q_T < 260$ GeV/ c . This represents the highest center-of-mass energy measurement of this quantity over the largest phase space available to date. The overall uncertainty of this measurement has been reduced compared with the previous measurements. We find that for $q_T < 30$ GeV/ c , the CSS resummation model used in RESBOS describes the data very well at all rapidities. Our data with $|y| > 2$ disfavor a variant of this model that incorporates an additional small- x form factor when g_1 , g_2 , and g_3 from large- x data is used. Using the BLNY parameterization for events with $q_T < 30$ GeV/ c , we obtain $g_2 = 0.77 \pm 0.06$ (GeV/ c) 2 , which is comparable with the current world average value [5]. We observe a disagreement between our data and NNLO calculations in the region $q_T > 30$ GeV/ c , where our distribution is higher than predicted by a factor of 1.25. However, the

NNLO calculation agrees in shape with our data when normalized at $q_T = 30 \text{ GeV}/c$.

We thank C. Balazs, Q. H. Cao, P. Nadolsky, F. Petriello, and C. P. Yuan for many useful discussions. We thank the staffs at Fermilab and collaborating institutions, and acknowledge support from the DOE and NSF (USA); CEA and CNRS/IN2P3 (France); FASI, Rosatom and RFBR (Russia); CAPES, CNPq, FAPERJ, FAPESP, and FUNDUNESP (Brazil); DAE and DST (India); Colciencias (Colombia); CONACyT (Mexico); KRF and KOSEF (Korea); CONICET and UBACyT (Argentina); FOM (The Netherlands); Science and Technology Facilities Council (United Kingdom); MSMT and GACR (Czech Republic); CRC Program, CFI, NSERC, and WestGrid Project (Canada); BMBF and DFG (Germany); SFI (Ireland); The Swedish Research Council (Sweden); CAS and CNSF (China); Alexander von Humboldt Foundation; and the Marie Curie Program.

*Visitor from Augustana College, Sioux Falls, SD, USA.

†Visitor from The University of Liverpool, Liverpool, United Kingdom.

‡Visitor from ICN-UNAM, Mexico City, Mexico.

§Visitor from II. Physikalisches Institut, Georg-August-University Göttingen, Germany.

||Visitor from Helsinki Institute of Physics, Helsinki, Finland.

¶Visitor from Universität Zürich, Zürich, Switzerland.

**Deceased.

- [1] P. B. Arnold and M. H. Reno, Nucl. Phys. B **319**, 37 (1989); R. J. Gonsalves, J. Pawlowski, and C-F. Wai, Phys. Rev. D **40**, 2245 (1989).
- [2] K. Melnikov and F. Petriello, Phys. Rev. D **74**, 114017 (2006).
- [3] J. Collins, D. Soper, and G. Sterman, Nucl. Phys. B **250**, 199 (1985).
- [4] P. B. Arnold and R. Kauffman, Nucl. Phys. B **349**, 381 (1991).
- [5] F. Landry *et al.*, Phys. Rev. D **67**, 073016 (2003). The values are $g_1 = 0.21 \pm 0.01 \text{ (GeV}/c)^2$, $g_2 = 0.68_{-0.02}^{+0.01} \text{ (GeV}/c)^2$, $g_3 = -0.6_{-0.04}^{+0.05}$, $Q_0 = 1.6 \text{ GeV}/c$.
- [6] P. Nadolsky, D. R. Stump, and C. P. Yuan, Phys. Rev. D **61**, 014003 (1999); **64**, 059903(E) (2001).
- [7] P. Nadolsky, D. R. Stump, and C. P. Yuan, Phys. Rev. D **64**, 114011 (2001).
- [8] S. Berge *et al.*, Phys. Rev. D **72**, 033015 (2005).
- [9] T. Affolder *et al.* (CDF Collaboration), Phys. Rev. Lett. **84**, 845 (2000).
- [10] B. Abbott *et al.* (D0 Collaboration), Phys. Rev. D **61**, 032004 (2000); B. Abbott *et al.* (D0 Collaboration), Phys. Rev. Lett. **84**, 2792 (2000).
- [11] V. Abazov *et al.* (D0 Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **565**, 463 (2006).
- [12] T. Andeen *et al.*, Fermilab Report No. FERMILAB-TM-2365, 2007.
- [13] R. Brun and F. Carminati, CERN Program Library Long Writup Report No. W5013, 1993 (unpublished).
- [14] C. Balazs and C. P. Yuan, Phys. Rev. D **56**, 5558 (1997).
- [15] E. Barberio and Z. Was, Comput. Phys. Commun. **79**, 291 (1994).
- [16] V. Blobel, The RUN manual, Regularized Unfolding for High-Energy Physics Experiments, OPAL Technical Note Report No. TN361, 1996.
- [17] J. Pumplin *et al.*, J. High Energy Phys. 07 (2002) pp. 012; D. Stump *et al.*, J. High Energy Phys. 10 (2003) 046.
- [18] G. D. Lafferty and T. R. Wyatt, Nucl. Instrum. Methods Phys. Res., Sect. A **355**, 541 (1995).
- [19] P. B. Arnold and M. H. Reno, Nucl. Phys. B **319**, 37 (1989); **330**, 284(E) (1990).
- [20] A. D. Martin, R. G. Roberts, W. J. Stirling, and R. S. Thorne, Phys. Lett. B **531**, 216 (2002).
- [21] R. Hamberg, W. van Neerven, and T. Matsuura, Nucl. Phys. B **359**, 343 (1991); **644**, 403(E) (2002).