

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

USDA National Wildlife Research Center - Staff  
Publications

U.S. Department of Agriculture: Animal and Plant  
Health Inspection Service

---

January 2003

# CHARACTERISTICS OF SPRING AND FALL BLACKBIRD ROOSTS IN THE NORTHERN GREAT PLAINS

George M. Linz

*USDA, APHIS, Wildlife Services, National Wildlife Research Center, Great Plains Field Station, 2110 Miriam Circle, Bismarck, ND 58501, USA, [george.m.linz@aphis.usda.gov](mailto:george.m.linz@aphis.usda.gov)*

Richard S. Sawin

*Department of Biological Sciences, North Dakota State University, Stevens Hall, Fargo, ND 58105, USA*

Mark W. Lutman

*Department of Biological Sciences, North Dakota State University, Stevens Hall, Fargo, ND 58105, USA*

H. Jeffrey Homan

*USDA, APHIS, Wildlife Services, National Wildlife Research Center, Great Plains Field Station, 2110 Miriam Circle, Bismarck, ND 58501, USA*

Linda B. Penry

*USDA, APHIS, Wildlife Services, National Wildlife Research Center, Great Plains Field Station, 2110 Miriam Circle, Bismarck, ND 58501, USA*

~~See next page for additional authors~~

Follow this and additional works at: [http://digitalcommons.unl.edu/icwdm\\_usdanwrc](http://digitalcommons.unl.edu/icwdm_usdanwrc)



Part of the [Environmental Sciences Commons](#)

---

Linz, George M.; Sawin, Richard S.; Lutman, Mark W.; Homan, H. Jeffrey; Penry, Linda B.; and Bleier, William J., "CHARACTERISTICS OF SPRING AND FALL BLACKBIRD ROOSTS IN THE NORTHERN GREAT PLAINS" (2003). *USDA National Wildlife Research Center - Staff Publications*. 242.  
[http://digitalcommons.unl.edu/icwdm\\_usdanwrc/242](http://digitalcommons.unl.edu/icwdm_usdanwrc/242)

This Article is brought to you for free and open access by the U.S. Department of Agriculture: Animal and Plant Health Inspection Service at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in USDA National Wildlife Research Center - Staff Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

---

**Authors**

George M. Linz, Richard S. Sawin, Mark W. Lutman, H. Jeffrey Homan, Linda B. Penry, and William J. Bleier

## CHARACTERISTICS OF SPRING AND FALL BLACKBIRD ROOSTS IN THE NORTHERN GREAT PLAINS

GEORGE M. LINZ, USDA, APHIS, Wildlife Services, National Wildlife Research Center, Great Plains Field Station, 2110 Miriam Circle, Bismarck, ND 58501, USA

RICHARD S. SAWIN, Department of Biological Sciences, North Dakota State University, Stevens Hall, Fargo, ND 58105, USA

MARK W. LUTMAN, Department of Biological Sciences, North Dakota State University, Stevens Hall, Fargo, ND 58105, USA

H. JEFFREY HOMAN, USDA, APHIS, Wildlife Services, National Wildlife Research Center, Great Plains Field Station, 2110 Miriam Circle, Bismarck, ND 58501, USA

LINDA B. PENRY, USDA, APHIS, Wildlife Services, National Wildlife Research Center, Great Plains Field Station, 2110 Miriam Circle, Bismarck, ND 58501, USA

WILLIAM J. BLEIER, Department of Biological Sciences, North Dakota State University, Stevens Hall, Fargo, ND 58105, USA

**Abstract:** To successfully reduce blackbird (Icteridae) damage to sunflower crops in the northern Great Plains, wildlife damage managers must be able to identify sites where large roosts could form in spring and fall. In 1998 and 1999, we measured several habitat and land-use characteristics of spring and fall blackbird roosts. Our objective was to quantify the relationship between roost formation and wetland habitat and land-use. We used data from 7 spring roosts and 12 fall blackbird roosts to set search guidelines that might increase the efficacy of locating potential roost sites in the Prairie Pothole Region of the northern Great Plains. Analysis of land use with aerial photography and on-site measurements of wetlands showed that roosts were usually established in larger-sized wetlands with sufficient standing water to produce large robust stands of cattail (*Typha* spp.). Large roosts often formed near harvested cornfields in spring and near ripening sunflower in fall.

**Key words:** blackbirds, cattails, northern Great Plains, Prairie Pothole Region, sunflower, *Typha* spp.

Proceedings of the 10<sup>th</sup> Wildlife Damage Management Conference. (K.A. Fagerstone, G.W. Witmer, Eds). 2003

---

### INTRODUCTION

Blackbirds, including red-winged blackbirds (*Agelaius phoeniceus*), common grackles (*Quiscalus quiscula*), and yellow-headed blackbirds (*Xanthocephalus xanthocephalus*), are known in the northern Great Plains for their predation on sunflower. Each spring, an estimated 52 million blackbirds migrate north into the northern Great Plains of the United States

and southern Canada to nest (Peer et al. 2003). In fall, these birds and their progeny, totaling 75 million, drift southward feeding on sunflower and other grain crops. During both migratory periods, these birds prefer cattail-choked (*Typha* spp.) wetlands as night roosts because they provide shelter from severe weather and predators. Local habitat characteristics of wetland roosts and surrounding land-use might be useful for

making *a priori* predictions of locations of active blackbird roosts. To be useful, predictive habitat characteristics should be practical to assess and consistent over relatively long periods of time. In this paper, we assess the practicality of using intrinsic and extrinsic site characteristics to predict use of wetlands as roost sites. The objective of our study was to find spring and fall blackbird roosts in the northern Great Plains and identify habitat and land-use traits common to these roosts. This knowledge should increase our ability to predict the locations of potential roosts based on the presence or absence of certain habitat and land-use variables. Wildlife managers and agricultural producers could implement damage management techniques more efficiently if accurate predictions of potential roost sites were available. For example, sunflower producers who plant near cattail-dominated wetlands might implement a management program by treating the wetland with glyphosate-based herbicide to eliminate the potential fall roost site and thus reduce the chances of significant bird damage (Linz and Hanzel 1997). On the other hand, wildlife managers could implement population management at spring roost sites to reduce numbers in the pre-breeding population (Linz et al. 2002a, 2002b).

## **METHODS**


### **Study Areas**

Spring 1998 and 1999.--We searched for spring migratory roosts in a 9-county area in east-central South Dakota (Figure 1). This area was chosen because it is considered to be the most northern region


where migrating flocks of blackbirds are large enough to warrant a baiting program (Sawin 1999). Eastern South Dakota lies in the Coteau Des Prairies physiographic region in the Prairie Pothole Region (PPR) of the northern Great Plains. The landscape is characterized by low, rolling hills and a variety of glacial till deposits; undrained lakes and small wetland basins abound (Johnson and Higgins 1997). Formerly unbroken tall-grass prairie, the land now consists mainly of pasture and row crops. Shelterbelts and windbreaks divide expansive field crops and surround farmsteads and residential areas. Corn and soybean are the principal crops in eastern South Dakota, comprising 92% of the planted crops. The annual long-term average temperature and precipitation in eastern South Dakota are 6.7° C and 58.4 cm, respectively (SDASS 2001).

Fall 1998 and 1999.--We searched for migratory blackbird roosts in Stutsman County, North Dakota (Figure 1). Stutsman County is located in the PPR and consists of many wetlands, croplands, and in the western part of the county, rangeland. Currently, Stutsman County is one of the top producers of sunflowers in the state, producing 117,000,000 kg per year (NDASS 2000). On average, Stutsman County receives 43.2 cm of rainfall per year and has an average temperature of 5.7° C (NDASS 2000). The growing season for Stutsman County is about 125 days and typically starts around 15 May, the average date of last frost, and ends around 20 September, average date of first frost (NDASS 2000). This growing season, coupled with ample precipitation, makes the region a highly productive agricultural area.

**Figure 1a. Location of 7 spring-migratory blackbird roosts in east-central South Dakota occupied in both 1998 and 1999.**



**Figure 1b. Location of 12 fall-migratory blackbird roosts in Stutsman County, North Dakota occupied in both 1998 and 1999.**



## Roost Detection

We found candidate roosts by following evening flight-lines or by aerially surveying wetlands during the day and checking for blackbird activity in the mornings and evenings. The aerial surveys were conducted with a fixed-wing airplane. We verified roost size by counting birds departing and entering the roost in the mornings and evenings, respectively, using standardized block-count methods (Meanley 1965). A minimum of 20 ha of emergent vegetation was needed to admit wetlands into the spring roost study, and >10 ha was needed for entry in the fall study. Minimum peak counts of 50,000 and 10,000 were required for spring and fall roosts, respectively. The spring counts were conducted between mid-March and late-April 1998-1999; the fall counts between mid-August and mid-September 1998-1999.

## Habitat Analyses

In 1998 and 1999, color aerial photographs of the spring and fall roosts were taken at elevations between 1,280 m and 3,140 m above ground level. The aerial photographs were converted to digital images and analyzed using a PC-based Geographic Information System (ARC/INFO; version 7.1.2 Environmental Systems Research Institute, Inc., Redlands, CA). The images were georeferenced, rectified, and projected to Albers conic equal-area projections. This projection provides an accurate cartographic representation of both distance and area.

Upland areas surrounding the wetland basin were removed before classifying the area coverage (ha) of open water and emergent vegetation in each wetland. An on-site visual assessment of the classified image was done to determine if each classified image was roughly accurate. One hundred pixels from each habitat type were

randomly selected to perform the accuracy assessment. We characterized landscape features surrounding the spring roosts by using Landsat data classified by crop type by the National Agricultural Statistics Service Spatial Analysis Research Section. Land use within a 16-km radius of each roost wetland was determined by GIS. We used 4 habitat categories: (1) corn; (2) wetland; (3) all cultivated (corn, small grain, soybean); and (4) other (shelterbelts/woodlots, roads, buildings, town, and fallow fields).

For analysis of habitats surrounding the fall roosts, we used 1.6 km x 3.2 km aerial photographs obtained from the Stutsman County Farm Services Agency in Jamestown, North Dakota. We estimated habitat coverage within a 4.8 km radius surrounding the wetland roost sites using a non-mapping technique (Marcum and Loftsgaarden 1980). The photographs were projected on to a 60-cm X 60-cm grid, and 21 random points per 1.6 km X 1.6 km quadrat were surveyed to assess the surrounding habitat. We used 4 habitat categories: (1) sunflower; (2) wetland; (3) all cultivated (small grain, sunflower, and soybean); (4) other (shelterbelts/woodlots, roads, buildings, town, and fallow fields). Coverage of open water and emergent vegetation within the wetland basins of fall roost sites were analyzed according to methods described for the spring roost study.

## RESULTS

Spring.--Wetland area (158  $\nabla$  61 ha, 95% C.I.) and percentages of open water (30  $\nabla$  14%, 95% C.I.) and emergent vegetation (70  $\nabla$  14%, 95% C.I.) in the 7 spring roosts varied little between study years (Table 1).

**Table 1. Habitat variables within and around spring roosts in 1998 and 1999 in east-central South Dakota.**

| Category  | Year    |                |         |                |                |                |
|---|---------|----------------|---------|----------------|----------------|----------------|
|   | 1998    |                | 1999    |                | Years Combined |                |
|   | Mean    | 95% CI         | Mean    | 95% CI         | Mean           | 95% CI         |
| Wetland size (hectares)   | 154     | 91.1-216.6     | 162     | 98.1-226.8     | 158            | 97.0-219.4     |
| Percentage of wetland covered by open water                     | 28      | 13.0-42.6      | 32      | 17.2-46.2      | 30             | 15.6-44.0      |
| Percentage of wetland covered by emergent vegetation            | 72      | 57.4-87.0      | 68      | 53.8-82.8      | 70             | 56.0-84.4      |
| Percentage of land planted to corn within 16 km radius of roost | 26      | 19.3-32.2      | 23      | 17.4-29.1      | 24             | 18.5-30.5      |
| Percentage of land cultivated within 16 km radius of roost      | 59      | 53.7-65.2      | 57      | 49.0-64.8      | 58             | 51.4-64.9      |
| Percentage of land covered by wetlands                          | 14      | 9.4-18.6       | 12      | 5.1-18.1       | 13             | 7.3-18.3       |
| Maximum number of blackbirds using roost                        | 144,279 | 10,187-278,370 | 110,114 | 15,238-204,991 | 127,196        | 14,482-239,910 |

Similarly, use of land surrounding the wetlands was similar between years, with percentages of land planted to corn and all cultivated crops combined, averaging 24% (∇ 6.5%, 95% C.I.) and 58% (∇ 6.9%, 95% C.I.), respectively. Wetlands comprised 13% (∇ 5.3%, 95% C.I.) of the surrounding landscape. Blackbird numbers in spring roosts averaged 144,000 and 110,000 in 1998 and 1999, respectively.

Fall.--Wetland area (94 ∇ 71 ha, 95% C.I.) and percentages of open water (51 ∇ 11%, 95% C.I.) and emergent vegetation (49 ∇ 11%, 95% C.I.) found in 12 fall roosts also varied little between study years (Table 2). Use of land surrounding the wetlands was only measured in 1998. Wetlands covered 20% of the surrounding land,

whereas sunflower and all cultivated crops combined covered 17% (∇ 3.2%, 95% C.I.) and 51% (∇ 6.9%, 95% C.I.) of the land area, respectively. Blackbird numbers in spring averaged 27,000 and 14,000 in 1998 and 1999, respectively.

Seasonal Comparison.--Aerial coverage of wetlands harboring spring blackbird roosts was 68% ( $\bar{x}$  = 158 ha) larger than fall ( $\bar{x}$  = 94 ha) roosts (Table 3). Emergent vegetation covered 70% of the spring roost wetlands and only 49% of the fall roost wetlands. The remaining portions of the wetlands were comprised of open water. Blackbird roosts were 6 times larger ( $\bar{x}$  = 127,000 ha) in the spring than in the fall ( $\bar{x}$  = 21,000 ha).

**Table 2. Habitat variables within and around fall roosts in 1998 and 1999 in central North Dakota.**

| Category  | Year   |               |        |              |                |              |
|---|--------|---------------|--------|--------------|----------------|--------------|
|   | 1998   |               | 1999   |              | Years Combined |              |
|   | Mean   | 95% CI        | Mean   | 95% CI       | Mean           | 95% CI       |
| Wetland size (hectares)   | 90     | 22.2-158.4    | 97     | 21.2-173.5   | 94             | 23.1-164.5   |
| Percentage of wetland covered by open water                           | 51     | 40.6-62.2     | 51     | 39.8-62.2    | 51             | 40.4-62.0    |
| Percentage of wetland covered by emergent vegetation                  | 49     | 37.8-59.4     | 49     | 37.8-60.2    | 49             | 38.0-59.6    |
| Percentage of land planted to sunflower within 4.8 km radius of roost | 17     | 14.2-20.3     | -      | -            | -              | -            |
| Percentage of land cultivated within 4.8 km radius of roost           | 51     | 43.7-57.5     | -      | -            | -              | -            |
| Percentage of land covered by wetlands                                | 20     | 15.7-23.9     | -      | -            | -              | -            |
| Maximum number of blackbirds using roost                              | 27,375 | 15,185-39,565 | 14,017 | 4,307-23,727 | 20,696         | 9,095-32,296 |

## DISCUSSION

The combination of cattail-dominated wetlands and grain crops provides a high quality stopover point for resident and migrating blackbirds. In March and April, blackbirds migrating through east-central South Dakota typically roost in large wetlands with dense emergent vegetation and feed primarily in harvested cornfields (Linz et al. 2002b). In August and September, blackbirds migrating through central North Dakota typically roost in smaller wetlands and primarily feed in ripening sunflower fields (Lutman 2000).

Patterns of wetland type and abundance vary across the PPR of the northern Great Plains (Batt et al. 1989). As a result, some locales are more likely than

others to attract blackbird roosts. East-central South Dakota and central North Dakota are two areas in the PPR that consistently contain large blackbird roosts (Barras 1996, Sawin 1999, Lutman 2000). At the highest level of choice, migrating blackbirds are probably constrained by the general features of the landscape as they proceed along the migration route. Inside of this scale, however, a more refined choice is made based upon the intrinsic features within a particular location (Hutto 1985). Thus, roost-site selection can be viewed as a hierarchical series of decisions made at a variety of spatial and temporal scales, similar to other habitat-use decisions made by birds (Johnson 1980, Hutto 1985).



**Table 3. Flock numbers and habitat variables within and around fall roosts in central North Dakota and spring roosts in east-central South Dakota in 1998 and 1999.**

| Category <sup>1</sup>   | Season            |              |                     |                |
|---|-------------------|--------------|---------------------|----------------|
|   | Fall <sup>1</sup> |              | Spring <sup>2</sup> |                |
|   | Mean              | 95% CI       | Mean                | 95% CI         |
| Wetland size (hectares)   | 94                | 23.1-164.5   | 158                 | 97.0-219.4     |
| Percentage of wetland covered by open water                               | 51                | 40.4-62.0    | 30                  | 15.6-44.0      |
| Percentage of wetland covered by emergent vegetation                      | 49                | 38.0-59.6    | 70                  | 56.0-84.4      |
| Percentage of land planted to sunflower <sup>3</sup> or corn <sup>4</sup> | 17                | 14.2-20.3    | 24                  | 18.5-30.5      |
| Percentage of land cultivated   | 51                | 43.7-57.4    | 58                  | 51.0-65.0      |
| Percentage of land covered by wetlands                                    | 20                | 15.7-23.9    | 13                  | 7.3-18.3       |
| Maximum number of blackbirds using roost                                  | 20,696            | 9,095-32,296 | 127,196             | 14,482-239,910 |

<sup>1</sup>Roosts located in central North Dakota

<sup>2</sup>Roosts located in east-central South Dakota

<sup>3</sup>Percentage of land planted within a 4.8 km radius of the roost

<sup>4</sup>Percentage of land planted within a 16.0 km radius of the roost

We found that most spring and fall blackbird roosts occurred in semipermanent wetlands with large expanses of dense emergent vegetation and abundant food available in the surrounding landscape. Although these similarities were present to a certain degree in spring and fall roosts, we did observe that fall roosts were notably smaller in number of birds per roost, wetland size, and ratio of open water: emergent vegetation. It was not unexpected to us that the number of birds per fall roost was smaller compared to spring, because the fall migration is a much more disorganized process than is the spring migration.

Differences in wetland size and open water: emergent vegetation ratios were perhaps a reflection of the inherent landscape pattern differences between east-

central South Dakota and central North Dakota. Wetlands for the latter were 20% of the surrounding landscape, whereas wetlands comprised only 13% of the South Dakota study area. Lastly, the proximity of an easily available food resource might influence roost-site choice, particularly in late summer and early fall when crops are ripening (Meanley 1965, Eiserer 1984, Lutman 2000). Having a food resource next to a roost site makes foraging more energy efficient for blackbirds and maximizes their feeding time and calorie intake. Otis and Kilburn (1988) showed that cattail-dominated wetlands near sunflower fields were attractive to blackbirds. Therefore, it is not surprising that many sunflower fields planted close to roosts often receive more damage than fields further away.

The exact point at which roosting habitat becomes unattractive to blackbirds does not seem easy to predict. Changes in roost size from year to year might be related to changes in the quality of the wetland itself, or the habitat that surrounds it. Land use around potential roost wetlands might affect spring roost status, but probably not to the degree that it affects fall roost selection, particularly the nearness of a preferred food source (Sawin 1999, Lutman 2000). During fall, many more small-sized wetlands are available, unlike spring, when roost choice is limited mainly to the larger-sized wetlands. Likewise, the number of blackbirds at a roost is affected by wetland size, amount of cattails, and water depth. Of all factors, water-depth (and concomitantly, cattail coverage) perhaps is the most important characteristic governing selection of a fall roost site (Meanley 1965, Lutman 2000). We have never found stable roosts in wetlands lacking standing water in the northern Great Plains. Periodic fluctuations in water levels throughout this region result in successional changes in wetlands that alter the distribution and abundance of quality roosting habitat (Van Der Valk and Davis 1979, Weller 1975). The location and size of active roosts will shift to reflect these climatically-induced habitat changes. Some wetlands serve as roosts for a number of consecutive years, while others last for only one season (Barras 1996, Sawin 1999, Lutman 2000).

### **Management Implications**

In the spring, look for roosts in large cattail-dominated wetlands that have harbored blackbirds in previous years. In the fall, search for roosts in wetlands located near sunflower fields. The best method of finding these roosts is to first aerially survey the landscape for likely locations and then follow-up with ground inspection. Aerial searches require approximately one-sixth of

the time required for ground surveys and provide a much better overview of the landscape in the study area. First-time searchers will find it useful to view previously identified roosts from the air and on the ground in order to become familiar with the appearance of potential roost wetlands. Conduct early season roost estimates as they are good indicators that a substantial roost might be forming.

### **ACKNOWLEDGMENTS**

We thank C. Gehri, M. Klich, R. Kostecke, and R. Wimberly for assistance with field work. R. Carlson and G. Nuechterlein provided comments on earlier drafts of this manuscript. These studies were part of the second and third author's M.S. thesis research. Funding was provided jointly by the National Wildlife Research Center (Protocol QA-437, as amended), within the Wildlife Services program of the United States Department of Agriculture, Animal and Plant Health Inspection Service, and by the Department of Biological Sciences at North Dakota State University.

### **LITERATURE CITED**

- BARRAS, A.E. 1996. Evaluation of the efficacy of DRC-1339 for reducing spring migratory blackbird populations. M.S. Thesis, North Dakota State University, Fargo, ND, USA.
- BATT, D.J.B., M.G. ANDERSON, C.D. ANDERSON, AND F.D. CASWELL. 1989. The use of prairie potholes by North American ducks. Pages 204-227 in A. van der Valk, editor. Northern Prairie Wetlands. Iowa University Press, Ames, IA, USA.
- EISERER, L.A. 1984. Communal roosting in birds. *Bird Behaviour* 5:61-80.
- HUTTO, R.L. 1985. Habitat selection by nonbreeding, migratory land birds. Pages 455-476 in Martin L. Cody, editor. *Habitat selection in birds*. Academic Press, Incorporated, Orlando, FL, USA.
- JOHNSON, R.R., AND K.F. HIGGINS. 1997. Wetlands resources of eastern South Dakota. South Dakota State University, Brookings, SD, USA.
- JOHNSON, D.H. 1980. The comparison of usage and

- availability measurements for evaluating resource preference. *Ecology* 61:65-71.
- LINZ, G.M., A.E. BARRAS, R.S. SAWIN, H.J. HOMAN, D.L. BERGMAN, AND W.J. BLEIER. 2002*b*. Spring migration phenology and habitat use of red-winged blackbirds in eastern South Dakota. *Vertebrate Pest Conference* 20:97-102.
- \_\_\_\_\_, G.A. KNUTSEN, H.J. HOMAN, AND W.J. BLEIER. 2002*a*. Baiting blackbirds in stubble grain fields during spring migration in South Dakota. *Crop Protection* 22:261-264.
- \_\_\_\_\_, AND J.J. HANZEL. 1997. Birds and sunflower. Pages 381-394 in Albert A. Schneiter editor. *Sunflower technology and production*. Soil Science Society of America, Incorporated, Publishers, Madison, WI, USA.
- LUTMAN, M.W. 2000. Location and habitat characteristics of late-summer blackbird roosts in the sunflower-growing region of North Dakota. M.S. Thesis, North Dakota State University, Fargo, ND, USA.
- MARCUM, C. L., AND D. O. LOFTSGAARDEN. 1980. A non-mapping technique for studying habitat preferences. *Journal of Wildlife Management* 44:963-968.
- MEANLEY, B. 1965. The roosting behavior of red-winged blackbirds in the southern United States. *Wilson Bulletin* 77:217-228.
- NORTH DAKOTA AGRICULTURAL SERVICE (NDASS). 2000. *North Dakota Agricultural Statistics*. Vol 69.
- OTIS, D.L., AND C.M. KILBURN. 1988. Influence of environmental factors on blackbird damage to sunflower. U.S. Fish and Wildlife Service, Technical Report 16.
- PEER, B.D., H.J. HOMAN, G.M. LINZ, AND W.J. BLEIER. 2003. Impact of blackbird damage to sunflower: bioenergetic and economic models. *Ecological Applications* 13:248-56.
- SAWIN, R.S. 1999. Habitat characteristics of spring blackbird roosts in eastern South Dakota. M.S. Thesis, North Dakota State University, Fargo, ND, USA.
- SOUTH DAKOTA AGRICULTURAL SERVICE (SDASS). 2001. *South Dakota Agricultural Statistics*. Vol 61.
- VAN DER VALK, A.G., AND C.B. DAVIS. 1979. A reconstruction of the recent vegetational history of a prairie marsh, Eagle Lake, Iowa, from its seed bank. *Aquatic Botany* 6:29-51.
- WELLER, M.W. 1975. Studies of cattail in relation to management for wildlife. *Iowa State Journal Resource* 49:383-412.