Brief Report

Substitution of conserved cysteine residues in wheat streak mosaic virus HC-Pro abolishes virus transmission by the wheat curl mite

B. A. Young¹, G. L. Hein², R. French¹, D. C. Stenger¹

Received 7 May 2007; Accepted 14 June 2007; Published online 7 August 2007 © Springer-Verlag 2007

Summary

Substitutions in the amino-proximal region of wheat streak mosaic virus (WSMV) HC-Pro were evaluated for effects on transmission by the wheat curl mite (*Aceria tosichella* Keifer). Alanine substitution at cysteine residues 16, 46 and 49 abolished vector transmission. Although alanine substitution at Cys₂₀ had no effect, substitution with arginine reduced vector transmission efficiency. Random substitutions at other positions (Lys₇ to Asn, Asn₁₉ to Ile, and Arg₄₅ to Lys) did not affect vector transmission. These results suggest that a zinc-finger-like motif (His₁₃-X2-Cys₁₆-X29-Cys₄₆-X2-Cys₄₉) in WSMV HC-Pro is essential for vector transmission.

*

Wheat streak mosaic virus (WSMV) is the type species of the genus *Tritimovirus* in the family *Potyviridae* [15]. Whereas viruses of the genus

Correspondence: Drake C. Stenger, USDA-ARS, San Joaquin Valley Agricultural Sciences Center, 9611 South Riverbend Avenue, Parlier, CA 93648, USA e-mail: dstenger@fresno.ars.usda.gov

Potyvirus are transmitted non-persistently by aphids, WSMV is transmitted semi-persistently by the wheat curl mite (Aceria tosichella Keifer). Both tritimoviruses [14, 17] and potyviruses [7, 19] encode HC-Pro homologues required for vector transmission. Although HC-Pro sequence conservation among the two genera is limited [6], functional domains of WSMV HC-Pro [16, 18] involved in vector transmission (amino-proximal) and polyprotein maturation (carboxy-proximal) are positioned similarly to that of potyvirus HC-Pro.

Alignment of WSMV HC-Pro with potyvirus HC-Pro is mostly ambiguous except for the proteinase domain in the carboxy-proximal third of the protein [6]. Nonetheless, the amino-proximal region of both potyvirus and tritimovirus HC-Pro are cysteine-rich. HC-Pro of both genera self-interact [5, 8, 11, 20]; mutations introduced into the cysteine-rich region of potyvirus HC-Pro have been shown to abolish self interaction [20] and aphid transmission [1, 2, 9]. Whether amino acid substitutions in the corresponding region of WSMV HC-Pro affect virus transmission by the wheat curl mite is unknown. Here, we address this issue by evaluating infectivity and vector transmission phe-

¹ United States Department of Agriculture – Agricultural Research Service and Department of Plant Pathology, University of Nebraska, Lincoln, NE, USA

² Department of Entomology, University of Nebraska Panhandle Research Center, Scottsbluff, NE, USA

B. A. Young et al.

notypes of WSMV bearing amino acid substitutions in the amino-proximal region of HC-Pro.

An infectious cDNA clone of the WSMV Sidney 81 isolate was constructed previously [4] and subsequently modified [12] to contain unique SalI and ApaI sites (pS81-SA) flanking the HC-Pro coding region. This SalI-ApaI fragment was subcloned into pALTER (Promega, Madison, WI) and served as a template for site-directed mutagenesis. Codons for each of four cysteine residues (Cys₁₆, Cys₂₀, Cys_{46} , and Cys_{49}) in the 5'-proximal coding region of WSMV HC-Pro were mutated individually to alanine codons. Site-directed mutagenesis also was used to generate double mutants in which pairs of cysteine codons (Cys₁₆ + Cys₂₀ and Cys₄₆ + Cys₄₉) were altered to alanine codons in the same construct. Primers (sequences in WSMV minus strand) used for site-directed mutagenesis (underline denotes nucleotide substitutions) were 5'-TTG CGCAGTTGTTGGCCACATCGTGATATTT CGG-3' (Cys₁₆ to Ala), 5'-CAGGTAACTTGC<u>TGC</u> GTTGTTGCACA-3' (Cys₂₀ to Ala), 5'-GTTC TACATTGGTCTGCCCTCAGATTATG-3' (Cys₄₆ to Ala), 5'-CCACTCTGTTCTTGCTTGGTCGCA CCTC-3' (Cys₄₉ to Ala), 5'-TCCAGGTAACTTGC TGCGTTGTTTGCCACATCGTGATATTTC-3' (Cys₁₆ to Ala and Cys₂₀ to Ala), and 5'-TTCC ACTCTGTTCT<u>TGC</u>TTGGTC<u>TGC</u>CCTCAGATT ATGTA-3' (Cys₄₆ to Ala and Cys₄₉ to Ala). The four single and two double mutations were verified by nucleotide sequencing. Each mutant SalI-ApaI fragment was excised from the pALTER derivatives and subsequently used to replace the SalI-ApaI fragment of pS81-SA. RNA was transcribed from each mutant construct and inoculated to wheat seedlings (cv. Arapahoe) essentially as described [4]. Transcripts from all six mutant constructs were infectious to wheat and generated wild-type systemic symptoms (Table 1). The infection status of transcript-inoculated plants was verified by reverse transcription-polymerase chain reaction (RT-PCR) of the HC-Pro region as described [12]. The stability of mutations introduced into WSMV HC-Pro was assessed by sequencing of plasmids bearing cloned inserts of RT-PCR products amplified from systemically infected test plants. No evidence for reversion was seen; all cysteine-to-alanine substitu-

Table 1. Infectivity of wheat streak mosaic virus with cysteine-to-alanine substitutions in HC-Pro

WSMV	Experiment	1 ^b	Experiment 2 ^b		
genotype ^a	Symptoms	RT- PCR	Symptoms	RT- PCR	
Cys ₁₆ to Ala	10/10	10/10	9/10	4/4	
Cys ₂₀ to Ala	10/10	10/10	10/10	4/4	
Cys ₁₆ to Ala+	7/10	7/10	10/10	4/4	
Cys ₂₀ to Ala	•	,	•	,	
Cys ₄₅ to Ala	7/10	9/10	8/10	4/4	
Cys ₄₉ to Ala	9/10	10/10	9/10	4/4	
Cys ₄₅ to Ala+	8/10	9/10	9/10	4/4	
Cys ₄₉ to Ala	•	•	•		
No virus control	0/10	0/4	0/10	0/4	

^a Subscript denotes HC-Pro amino acid coordinate. Stability of mutations verified by sequencing of RT-PCR products cloned from systemically infected wheat plants.

tions were retained by WSMV in systemically infected plants (data not shown).

Previously, a series of 35 randomly generated point mutations in WSMV HC-Pro were evaluated for systemic infectivity, the ability to generate primary infection foci, and autoproteolytic activity [18]. Among the random mutations analyzed, four nonsynonymous point substitutions (Lys₇ to Asn, Asn₁₉ to Ile, Cys₂₀ to Arg, and Arg₄₅ to Lys) that occurred within the amino-proximal region of WSMV HC-Pro had no effect on pathogenicity or proteinase activity. These four randomly generated mutations, along with the cysteine-to-alanine site-directed mutations described above, were evaluated for phenotypic effects on virus transmission by the wheat curl mite.

Wheat curl mite transmission assays were conducted essentially as described [17]. Wheat seedlings to be used as source plants for vector transmission assays were inoculated with transcripts derived from each of the ten mutant constructs. Wheat seedlings inoculated with transcripts of pS81-SA or non-inoculated wheat seedlings served as positive or negative (mock) control source plants, respectively. Total RNA was extracted [10] from leaf samples 12–14 days post inoculation and used as template to determine infection status of source

^b Number of plants infected/number of inoculated plants evaluated. Infection status of plants determined by visual inspection for symptoms and by RT-PCR.

Table 2. Vector transmission of wheat streak mosaic virus (WSMV) mutants bearing amino acid substitutions in HC-Pro

WSMV Genotype	Transmission to test plants ^a Acquisition source plant								
	A	В	С	D	Е	F			
	Cys ₁₆ to Ala	0/4	0/4	0/4	0/4	0/4	ND^b	0/20 (0%)	
Cys ₂₀ to Ala	3/4	3/4	4/4	3/4	3/4	4/4	20/24 (83%)*		
Cys ₁₆ to Ala &	0/4	0/4	0/4	0/4	0/4	ND^b	0/20 (0%)		
Cys ₂₀ to Ala	,	,	,	,	,		, , ,		
Cys ₄₆ to Ala	0/4	0/4	0/4	0/4	0/4	ND^b	0/20 (0%)		
Cys ₄₉ to Ala	0/4	0/4	0/4	0/4	0/4	0/4	0/24 (0%)		
Cys ₄₆ to Ala &	0/4	0/4	0/4	0/4	0/4	0/4	0/24 (0%)		
Cys ₄₉ to Ala	,	,	,	,	,	,	, , , ,		
Lys ₇ to Asn	4/4	4/4	3/4	3/4	3/4	2/4	19/24 (79%)*		
Asn ₁₉ to Ile	4/4	4/4	4/4	2/4	2/4	4/4	20/24 (83%)*		
Cys ₂₀ to Arg	0/4	3/4	1/4	0/4	0/4	0/4	4/24 (17%)		
Arg ₄₅ to Lys	4/4	4/4	4/4	1/4	3/4	4/4	20/24 (83%)*		
Wild type (S81-SA)	3/4	4/4	3/4	4/4	4/4	3/4	21/24 (87%)*		
No virus (mock)	0/4	0/4	0/4	0/4	0/4	0/4	0/24 (0%)		

^a Number of test plants infected/number test plants inoculated.

plants by RT-PCR as described [12]. The presence of each mutation was verified by sequencing of one RT-PCR product per mutant per experimental replicate (data not shown). Source plants (three per mutant per experimental replicate, except as noted in Table 2) were each colonized with ten wheat curl mites transferred from an aviruliferous colony. After a three-week acquisition access period, wheat curl mite progeny were transferred in groups of ten from each source plant to healthy wheat seedlings serving as test plants (four per source plant). Following a four-week inoculation access period, test plants were harvested and leaves stored at -80 °C. To prevent movement of wheat curl mites among plants, each source plant and each test plant was individually caged. The infection status of each test plant was determined by RT-PCR as described above.

Results of the wheat curl mite transmission assays are presented in Table 2. Of the four single cysteine-to-alanine substitutions, the only mutant that retained vector transmission competence was Cys₂₀ to Ala. The remaining three single substitution mutants and the two WSMV genomes bear-

ing double cysteine-to-alanine mutations were not transmitted by the wheat curl mite. Three of the four randomly generated substitution mutants (Lys₇ to Asn, Asn₁₉ to Ile, and Arg₄₅ to Lys) were transmitted by the wheat curl mite at levels (79-83%) similar to that of the positive control S81-SA (87%). The mutant bearing the Cys₂₀-to-Arg substitution retained vector transmission competence, albeit at a reduced level (17%) compared to S81-SA or the alanine substitution mutant at Cys₂₀ (83%). All test plants colonized with wheat curl mites transferred from uninoculated (mock) source plants tested negative for WSMV infection by RT-PCR. Direct sequencing of RT-PCR products (two per mutant construct) amplified from WSMVinfected test plants revealed no evidence for reversion (data not shown) following transmission by the wheat curl mite.

We propose that the amino-proximal region of WSMV HC-Pro contains a non-canonical zinc finger motif [3] (His₁₃-X2-Cys₁₆-X29-Cys₄₆-X2-C₄₉), in which the three conserved cysteine residues (Cys_{16,46,49}), along with a conserved histidine resi-

^b ND Not determined.

^c Asterisk denotes vector transmission efficiency not significantly different (P>0.05) from wild type (S81-SA) based on ANOVA and Tukey's multiple comparison tests.

2110 B. A. Young et al.




Fig. 1. Multiple alignment of the amino-proximal region of HC-Pro for five wheat streak mosaic virus (WSMV) strains (Sidney 81, Type, TK1, Czech, and El Batán 3), oat necrotic mottle virus (ONMV), and members of five species of the genus *Potyvirus* (zucchini yellow mosaic virus [ZYMV], turnip mosaic virus [TuMV], tobacco vein mottling virus [TVMV], tobacco etch virus [TEV], and potato virus Y [PVY]). Coordinates of amino acid substitutions introduced into Sidney 81 HC-Pro are indicated at the top. Cysteine residues are shown in red, and includes four Sidney 81 residues mutated to alanine (and also arginine for Cys₂₀). Green denotes other Sidney 81 HC-Pro amino acid residues subject to substitution mutations. Asterisks (*) at the bottom denote a conserved histidine residue and three conserved cysteine residues constituting a zinc-finger-like motif in WSMV HC-Pro (His₁₃-X2-Cys₁₆-X29-Cys₄₆-X2-Cys₄₉)

due (His₁₃) in a favorable position relative to a cysteine residue (His-X2-C), constitute the key residues that interact with a zinc ion (Fig. 1). This same zinc-finger-like motif occurs in HC-Pro of oat necrotic mottle virus (ONMV), the tritimovirus species most closely related to WSMV [13], and is similar to that conserved among divergent potyviruses (Fig. 1). That mutations in the zinc-fingerlike motif of HC-Pro for both potyviruses and WSMV abolished vector transmission suggests that, despite very different modes of transmission by distinct vector taxa, at least some aspect(s) of the mechanism by which HC-Pro mediates vector transmission is similar among genera of the family Potyviridae. Although a fourth cysteine residue (Cys₂₀) could be mutated to alanine without affecting transmission by the wheat curl mite, substitution of this residue with arginine reduced vector transmission efficiency. If the hypothesis (wheat curl mite transmission of WSMV requires zinc bound to the zinc-finger-like motif of HC-Pro) is correct, then these observations suggest that Cys₂₀ does not directly interact with the metal ion, but instead is likely part of a loop structure which tolerates some substitutions. Interestingly, ONMV

HC-Pro has leucine at this position rather than cysteine (Fig. 1), further suggesting that the zinc-finger-like motif in tritimovirus HC-Pro does not require Cys₂₀. To test the hypothesis stated above, non-transmissible WSMV HC-Pro mutants should be evaluated for zinc ion binding.

Acknowledgments

We thank Kathryn Stenberg for technical assistance. Mention of proprietary or brand names is necessary to report factually on available data; however, the USDA neither guarantees nor warrants the standard of the product, and the use of the name by USDA implies no approval to the exclusion of others that also may be suitable. This article is in the public domain and not copyrightable. It may be freely reprinted with customary crediting of source.

References

- Atreya CD, Atreya PL, Thornbury DW, Pirone TP (1992) Site-directed mutations in the potyvirus HC-PRO gene affect helper component activity, virus accumulation, and symptom expression in infected tobacco plants. Virology 191: 106–111
- 2. Atreya CD, Pirone TP (1993) Mutational analysis of the helper component-proteinase gene of a potyvirus:

- Effects of amino acid substitutions, deletions, and gene replacement on virulence and aphid transmissibility. Proc Natl Acad Sci USA 90: 11919–11923
- Berg JM (1990) Zinc fingers and other metal-binding domains. J Biol Chem 265: 6513–6516
- 4. Choi I-R, French R, Hein GL, Stenger DC (1999) Fully biologically active *in vitro* transcripts of the eriophyid mite-transmitted wheat streak mosaic tritimovirus. Phytopathology 89: 1182–1185
- Choi I-R, Stenger DC, French R (2000) Multiple interactions among proteins encoded by the mite-transmitted wheat streak mosaic tritimovirus. Virology 267: 185–198
- French R, Stenger DC (2005) Complete nucleotide sequences of Agropyron mosaic virus and Hordeum mosaic virus support reciprocal monophyly of the genera Potyvirus and Rymovirus in the family Potyviridae. Arch Virol 150: 299–312
- Govier DA, Kassanis B (1974) A virus-induced component of plant sap needed when aphids acquire potato virus Y from purified preparation. Virology 61: 420–426
- Guo D, Merits A, Saarma M (1999) Self-association and mapping of interaction domains of helper component-proteinase of potato potyvirus. J Gen Virol 80: 1127–1131
- Llave C, Martinez B, Diaz-Ruiz JR, Lopez-Abella D (2002) Amino acid substitutions within the cys-rich domain of the tobacco etch potyvirus HC-Pro result in loss of transmissibility by aphids. Arch Virol 147: 2365–2375
- McNeil JE, French R, Hein GL, Baenziger PS, Eskridge KM (1996) Characterization of genetic variability among natural populations of wheat streak mosaic virus. Phytopathology 86: 1222–1227
- Ruiz-Ferrer V, Boskovic J, Alfonso C, Rivas G, Llorca O, Lopez-Abella D, Lopez-Moya JJ (2005) Structural analysis of tobacco etch potyvirus HC-Pro oligomers involved in aphid transmission. J Virol 79: 3758–3765

- 12. Stenger DC, French R (2004) Functional replacement of *Wheat streak mosaic virus* HC-Pro with the corresponding cistron from a diverse array of viruses in the family *Potyviridae*. Virology 323: 257–267
- Stenger DC, French R (2004) Complete nucleotide sequence of *Oat necrotic mottle virus*: a distinct *Tritimovirus* species (family *Potyviridae*) most closely related to *Wheat streak mosaic virus*. Arch Virol 149: 633–640
- Stenger DC, French R, Gildow FE (2005) Complete deletion of Wheat streak mosaic virus HC-Pro: a null mutant is viable for systemic infection. J Virol 79: 12077–12080
- 15. Stenger DC, Hall JS, Choi I-R, French R (1998) Phylogenetic relationships within the family *Potyviridae*: Wheat streak mosaic virus and brome streak mosaic virus are not members of the genus *Rymovirus*. Phytopathology 88: 782–787
- Stenger DC, Hein GL, French R (2006) Nested deletion analysis of Wheat streak mosaic virus HC-Pro: mapping of domains affecting polyprotein processing and eriophyid mite transmission. Virology 350: 465–474
- 17. Stenger DC, Hein GL, Gildow FE, Horken KM, French R (2005) Plant virus HC-Pro is a determinant of eriophyid mite transmission. J Virol 79: 9054–9061
- Stenger DC, Young BA, French R (2006) Random mutagenesis of Wheat streak mosaic virus HC-Pro: noninfectious interfering mutations in a gene dispensable for systemic infection of plants. J Gen Virol 87: 2741–2747
- Thornbury DW, Hellmann GM, Rhoads RE, Pirone TP (1985) Purification and characterization of potyvirus helper component. Virology 144: 260–267
- 20. Urcuqui-Inchima S, Maia IG, Drugeon G, Haenni A-L, Bernadi F (1999) Effect of mutations within the cys-rich region of potyvirus helper component-proteinase on self interaction. J Gen Virol 80: 2809–2812