

1980

Den and Den-Site Characteristics of Coyotes (*Canis latrans*) in Southeastern Nebraska

Donald P. Althoff

Pennsylvania State University - Main Campus

Follow this and additional works at: <http://digitalcommons.unl.edu/tnas>



Part of the [Life Sciences Commons](#)

Althoff, Donald P., "Den and Den-Site Characteristics of Coyotes (*Canis latrans*) in Southeastern Nebraska" (1980). *Transactions of the Nebraska Academy of Sciences and Affiliated Societies*. 277.

<http://digitalcommons.unl.edu/tnas/277>

This Article is brought to you for free and open access by the Nebraska Academy of Sciences at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Transactions of the Nebraska Academy of Sciences and Affiliated Societies by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

DEN AND DEN-SITE CHARACTERISTICS OF COYOTES (*CANIS LATRANS*) IN SOUTHEASTERN NEBRASKA

Donald P. Althoff

Land and Water Research Building
Pennsylvania State University
University Park, Pennsylvania 16802

Sixteen pup-rearing den-sites of coyotes (*Canis latrans*) were located in southeastern Nebraska from 1976-1978. Although there was no selection for sites close to water or away from roads, dens were always located 400 m or farther from occupied farmsteads. The mean distance between neighboring dens was 3.87 km. Twelve sites were located within 1 m of brome grass (*Bromus inermis*); 15 dens were located where two cover-types met. Twelve dens were dug in loam soil; four in clay. Mean number of entrances for 10 excavated dens was 1.4, with the majority opening to the east and north.

† † †

INTRODUCTION

Denning habits of coyotes are important in understanding their ecology. Studies of coyote dens in several regions have indicated considerable variation in the type of sites selected as well as in other characteristics (Young and Jackson, 1951; Gier, 1968; Hallett, 1977; Rock, 1978). However, denning habits of coyotes in an area of intensive cultivation such as southeastern Nebraska have not been reported.

Here, I report on den ecology of coyotes in southeastern Nebraska. Data were collected from 1976-1978 while studying the spatial relationships and livestock depredations of coyotes in the same area (Andelt, 1976; Althoff, 1978). In this paper, "den" refers to the opening, burrow, and chamber, and "den-site" refers to the den and the immediate surrounding area.

STUDY AREA

Dens were located in Lancaster and Seward counties (Fig. 1). The topography of the area consists of low, rolling hills with sloping-upland and bottom-land areas. Grain sorghum, corn, wheat, oats, alfalfa, and native grasses, predominantly brome, constitute most of the vegetation in this intensively farmed portion of Nebraska. Waterways consist of brome and other native grasses, while most woody vegetation occurs along creek bottoms and fence rows.

RESULTS AND DISCUSSION

Den-Site Identification

Sixteen active pup-rearing den-sites were located in the study area (Fig. 1). Andelt (1976) and local residents provided location of six dens in 1976; I located eight dens in 1977 and two in 1978 during ground searches conducted from May through July. Pup-rearing dens were distinguished from non-pup-rearing dens and badger (*Taxidea taxus*) and rabbit (*Sylvilagus floridanus*) burrows by the presence of pups, food scraps, worn trails, bedsites, and/or diameter of den entrances.

Den-Site Characteristics

Frequency distributions of den-sites from nearest water sources, roads (including major highways and county roads), and occupied farmsteads are shown in Figure 2. Apparently, close proximity to water was not a critical site factor since more than one-half of the dens were 200 m or farther from a pond, creek, or intermittent stream. Hallett (1977) and Rock (1978) reported that most coyotes den within 100 m of water in Missouri and Saskatchewan, respectively. Although they did not present data, Young and Jackson (1951) and Gier (1968) concluded that coyotes commonly den near a water source.

Distances of den-sites from roads and occupied farmsteads provide an index to coyote tolerance of human activity. No den was located closer than 400 m to an occupied homestead; most were farther than 600 m (Fig. 2). In contrast, eight dens were less than 400 m from a road. A grid of county-maintained roads divides most of the land in the Lancaster County study area into 2.56 km² (sq mi) sections (Fig. 1), with one or more occupied farmsteads usually located in each section. Those dens within 400 m of a road were always situated in the farthest possible area from any homestead in the same or adjoining section.




FIGURE 1. Location of 15 pup-rearing den-sites, Lancaster County study area. One den-site located in Seward County is not shown. Coyote family home-range boundaries were determined from radio telemetry data (Andelt, 1976; Althoff 1978).




FIGURE 2. Distances between coyote dens and nearest water source, road, and occupied homestead.

The mean distance to the nearest neighboring den over the two-year period in the Lancaster County study area was 3.87 km. This compares with a mean of 2.1 km and 5.2 km between neighboring dens in parts of Missouri (Hallett, 1977) and Saskatchewan (Rock, 1978), respectively. Statistical analysis of den dispersion (Table I) was based on the method of Clark and Evans (1954). The ratio, R , between observed and expected mean-distances from each den to its nearest neighbor indicates the departure of the dispersion pattern from random. The R ratio will equal 0 if there is no spacing, 1 if there is random spacing, and 2.1491 if maximum spacing between individuals exists. Statistical significance is based on C , the standard variate of the normal curve. Thus, den dispersion as indicated by R appeared to be non-random toward maximum spacing, but the C values showed no statistical significance ($P > 0.05$, Table I). However, home ranges of coyotes based on radio telemetry (Andelt, 1976; Althoff, 1978) indicated that pup-rearing dens are in the core and not on the edge of the home range (Fig. 1).

Habitats of den-sites ranged from open patches of native grass to edges of dense thickets. Eleven den-sites were located in fence rows, native-grass patches, drainage ditches, or draws where vegetation seldom exceeded a height of 0.5 m; five sites were in thickets. None of the dens was located in a woodlot. Brome grass was within 1 m of 12 dens; four dens were concealed by the grass (see Althoff, 1979, for description of camouflaged den-entrances). Cover types of sorghum, wheat, or corn were within 10 m of all but one den. This is not surprising considering the intensity of agriculture in the area. Although not as much of the study area was comprised of croplands, Rock (1978) located most coyote dens in native pasture in Saskatchewan. Hallett (1977) found most Missouri dens situated along the edge between two cover-types.

Suitable denning-habitat within the core of home ranges probably dictated exact placement of dens and accounted for some variation of distances between neighboring dens. The data strongly suggest that avoidance of human activity as well

TABLE I. Dispersion of occupied coyote dens based on nearest neighbor method (Clark and Evans, 1954) for determining spatial relationships.

Year	Area (km ²)	Number of Dens*	R†	C†	Level of Significance
1976	100	4	1.45	1.73	> 0.05
1977	180	6	1.40	1.86	> 0.05
Combined	280	10	1.46	1.67	> 0.05

*Secondary den-sites and one den-site outside Lancaster County study area were not used in analysis.

†See text for description of R and C values.

as neighboring coyote families are important factors in site location.

Den Characteristics

Twelve dens were dug in loam soil; four in clay. Coyotes may have selected the former because of the ease of digging in loam soils. Judging from the amount of soil in front of the den, I suspect most of the dens were originally dug the previous year but not necessarily by coyotes. Badgers are common in the area and were likely responsible for excavating most of the original burrows. The slope of the terrain where the dens were located ranged from 0° to 90° (flat ground to vertical banks) with eight in the range of 10° to 60°. In other regions, coyotes commonly select well-drained locations as sites for pup-rearing dens (Young and Jackson, 1951; Gier, 1968; Hallett, 1977; Rock, 1978).

The mean number of actively used openings per den was 1.4 (range: 1 to 3). (This does not include 1976 dens because I did not visit those sites during the pup-rearing period.) Andelt (personal communication) found eight active holes at a den-site in 1976. Although the same female coyote prepared this den-site in 1977 and 1978, it was not used for rearing pups in either year.

The orientation of 14 openings at 10 dens in 1977 and 1978 is shown in Figure 3. North- and east-facing openings were most common; no den opened to the southeast or northwest. In contrast, Gier (1968), Hallett (1977), and Rock (1978) reported that most of the dens they located were oriented toward warmer, south-facing slopes. Perhaps the slope of the terrain or prevailing wind are more important factors than southern exposure in site selection. Dens opening to the north, northeast, or east would usually be protected from

south-southwesterly prevailing winds such as the case in southeastern Nebraska.

The mean diameter of the main entrances was 32 cm (range: 28 cm to 40 cm) for the six dens which were completely excavated. The distance between the main entrance and the main chamber was 2.5 cm to 11.0 cm. Soil depth to the main-chamber ceiling ranged from 61 cm to 122 cm. Chamber heights, floor-to-ceiling, ranged from 18 cm to 51 cm.




FIGURE 3. Orientation of 14 den-openings at 10 pup-rearing dens.

CONCLUSIONS

These observations suggest that coyotes were highly selective in the placement of dens. Although no single characteristic was noted at all den sites, a "typical" pup-rearing den in southeastern Nebraska might be described as follows: Located at least 400 m from the nearest occupied farmstead in an area where two or more cover types meet, one of which is brome grass, the den would be dug in loam soil with a single, active entrance, at least 28 cm in diameter, opening to the north or east.

ACKNOWLEDGMENTS

I thank W. F. Andelt, S. A. Kilpatrick, B. R. Mahan, and G. Retzlaff for help in locating dens, and H. T. Gier and B. R. Mahan for critically reviewing earlier drafts of the manuscript.

REFERENCES

- Althoff, D. P. 1978. Social and spatial relationships of coyote families and neighboring coyotes. Master of Science Thesis, University of Nebraska-Lincoln: 80p.
- _____. 1979. Camouflaged coyote den entrances. *Prairie Naturalist*, 11:23-25.
- Andelt, W. F. 1976. Ecology of suspected damaging coyotes and their interaction with domestic poultry and livestock. Master of Science Thesis, University of Nebraska-Lincoln: 85p.
- Clark, P. J., and F. C. Evans. 1954. Distance to nearest neighbor as a measure of spatial relationships in populations. *Ecology*, 35:445-453.
- Gier, H. T. 1968. Coyotes in Kansas. *Kansas Agricultural Experiment Station Bulletin*, 393:118p.
- Hallett, D. L. 1977. Post-natal mortality, movements, and den sites of Missouri coyotes. Master of Science Thesis, University of Missouri, Columbia: 144p.
- Rock, T. W. 1978. An evaluation of seasonal coyote control techniques and sheep losses in Saskatchewan. Master of Science Thesis, University of Nevada, Reno: 54p.
- Young, S. P., and H. H. T. Jackson. 1951. *The clever coyote*. Harrisburg, Pennsylvania, The Stackpole Company: 411p.