

2009

Evidence of WW and WZ Production with lepton + jets Final States in pp Collisions at $\sqrt{s} = 1.96$ TeV

V. M. Abazov

Joint Institute for Nuclear Research, Dubna, Russia

Kenneth A. Bloom

University of Nebraska - Lincoln, kbloom2@unl.edu

Gregory R. Snow

University of Nebraska-Lincoln, gsnow1@unl.edu

D0 Collaboration

Follow this and additional works at: <http://digitalcommons.unl.edu/physicsbloom>



Part of the [Physics Commons](#)

Abazov, V. M.; Bloom, Kenneth A.; Snow, Gregory R.; and Collaboration, D0, "Evidence of WW and WZ Production with lepton + jets Final States in pp Collisions at $\sqrt{s} = 1.96$ TeV" (2009). *Kenneth Bloom Publications*. 297.
<http://digitalcommons.unl.edu/physicsbloom/297>

This Article is brought to you for free and open access by the Research Papers in Physics and Astronomy at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Kenneth Bloom Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Evidence of WW and WZ Production with lepton + jets Final States in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV

V. M. Abazov,³⁶ B. Abbott,⁷⁵ M. Abolins,⁶⁵ B. S. Acharya,²⁹ M. Adams,⁵¹ T. Adams,⁴⁹ E. Aguilo,⁶ M. Ahsan,⁵⁹ G. D. Alexeev,³⁶ G. Alkhazov,⁴⁰ A. Alton,^{64,*} G. Alverson,⁶³ G. A. Alves,² M. Anastasoiaie,³⁵ L. S. Ancu,³⁵ T. Andeen,⁵³ B. Andrieu,¹⁷ M. S. Anzelc,⁵³ M. Aoki,⁵⁰ Y. Arnaud,¹⁴ M. Arov,⁶⁰ M. Arthaud,¹⁸ A. Askew,^{49,†} B. Åsman,⁴¹ A. C. S. Assis Jesus,³ O. Atramentov,⁴⁹ C. Avila,⁸ F. Badaud,¹³ L. Bagby,⁵⁰ B. Baldin,⁵⁰ D. V. Bandurin,⁵⁹ P. Banerjee,²⁹ S. Banerjee,²⁹ E. Barberis,⁶³ A.-F. Barfuss,¹⁵ P. Bargassa,⁸⁰ P. Baringer,⁵⁸ J. Barreto,² J. F. Bartlett,⁵⁰ U. Bassler,¹⁸ D. Bauer,⁴³ S. Beale,⁶ A. Bean,⁵⁸ M. Begalli,³ M. Begel,⁷³ C. Belanger-Champagne,⁴¹ L. Bellantoni,⁵⁰ A. Bellavance,⁵⁰ J. A. Benitez,⁶⁵ S. B. Beri,²⁷ G. Bernardi,¹⁷ R. Bernhard,²³ I. Bertram,⁴² M. Besançon,¹⁸ R. Beuselinck,⁴³ V. A. Bezzubov,³⁹ P. C. Bhat,⁵⁰ V. Bhatnagar,²⁷ G. Blazey,⁵² F. Blekman,⁴³ S. Blessing,⁴⁹ K. Bloom,⁶⁷ A. Boehnlein,⁵⁰ D. Boline,⁶² T. A. Bolton,⁵⁹ E. E. Boos,³⁸ G. Borissov,⁴² T. Bose,⁷⁷ A. Brandt,⁷⁸ R. Brock,⁶⁵ G. Brooijmans,⁷⁰ A. Bross,⁵⁰ D. Brown,⁸¹ X. B. Bu,⁷ N. J. Buchanan,⁴⁹ D. Buchholz,⁵³ M. Buehler,⁸¹ V. Buescher,²² V. Bunichev,³⁸ S. Burdin,^{42,‡} T. H. Burnett,⁸² C. P. Buszello,⁴³ P. Calfayan,²⁵ S. Calvet,¹⁶ J. Cammin,⁷¹ M. A. Carrasco-Lizarraga,³³ E. Carrera,⁴⁹ W. Carvalho,³ B. C. K. Casey,⁵⁰ H. Castilla-Valdez,³³ S. Chakrabarti,⁷² D. Chakraborty,⁵² K. M. Chan,⁵⁵ A. Chandra,⁴⁸ E. Cheu,⁴⁵ D. K. Cho,⁶² S. Choi,³² B. Choudhary,²⁸ L. Christofek,⁷⁷ T. Christoudias,⁴³ S. Cihangir,⁵⁰ D. Claes,⁶⁷ J. Clutter,⁵⁸ M. Cooke,⁵⁰ W. E. Cooper,⁵⁰ M. Corcoran,⁸⁰ F. Couderc,¹⁸ M.-C. Cousinou,¹⁵ S. Crépe-Renaudin,¹⁴ V. Cuplov,⁵⁹ D. Cutts,⁷⁷ M. Cwiok,³⁰ H. da Motta,² A. Das,⁴⁵ G. Davies,⁴³ K. De,⁷⁸ S. J. de Jong,³⁵ E. De La Cruz-Burelo,³³ C. De Oliveira Martins,³ K. DeVaughan,⁶⁷ F. Déliot,¹⁸ M. Demarteau,⁵⁰ R. Demina,⁷¹ D. Denisov,⁵⁰ S. P. Denisov,³⁹ S. Desai,⁵⁰ H. T. Diehl,⁵⁰ M. Diesburg,⁵⁰ A. Dominguez,⁶⁷ T. Dorland,⁸² A. Dubey,²⁸ L. V. Dudko,³⁸ L. Duflot,¹⁶ S. R. Dugad,²⁹ D. Duggan,⁴⁹ A. Duperrin,¹⁵ S. Dutt,²⁷ J. Dyer,⁶⁵ A. Dyshkant,⁵² M. Eads,⁶⁷ D. Edmunds,⁶⁵ J. Ellison,⁴⁸ V. D. Elvira,⁵⁰ Y. Enari,⁷⁷ S. Eno,⁶¹ P. Ermolov,^{38,§} H. Evans,⁵⁴ A. Evdokimov,⁷³ V. N. Evdokimov,³⁹ A. V. Ferapontov,⁵⁹ T. Ferbel,^{61,71} F. Fiedler,²⁴ F. Filthaut,³⁵ W. Fisher,⁵⁰ H. E. Fisk,⁵⁰ M. Fortner,⁵² H. Fox,⁴² S. Fu,⁵⁰ S. Fuess,⁵⁰ T. Gadfort,⁷⁰ C. F. Galea,³⁵ C. Garcia,⁷¹ A. Garcia-Bellido,⁷¹ V. Gavrilov,³⁷ P. Gay,¹³ W. Geist,¹⁹ W. Geng,^{15,65} C. E. Gerber,⁵¹ Y. Gershtein,^{49,†} D. Gillberg,⁶ G. Ginther,⁷¹ B. Gómez,⁸ A. Goussiou,⁸² P. D. Grannis,⁷² H. Greenlee,⁵⁰ Z. D. Greenwood,⁶⁰ E. M. Gregores,⁴ G. Grenier,²⁰ Ph. Gris,¹³ J.-F. Grivaz,¹⁶ A. Grohsjean,²⁵ S. Grünendahl,⁵⁰ M. W. Grünewald,³⁰ F. Guo,⁷² J. Guo,⁷² G. Gutierrez,⁵⁰ P. Gutierrez,⁷⁵ A. Haas,⁷⁰ N. J. Hadley,⁶¹ P. Haefner,²⁵ S. Hagopian,⁴⁹ J. Haley,⁶⁸ I. Hall,⁶⁵ R. E. Hall,⁴⁷ L. Han,⁷ K. Harder,⁴⁴ A. Harel,⁷¹ J. M. Hauptman,⁵⁷ J. Hays,⁴³ T. Hebbeker,²¹ D. Hedin,⁵² J. G. Hegeman,³⁴ A. P. Heinson,⁴⁸ U. Heintz,⁶² C. Hensel,^{22,§} K. Herner,⁷² G. Hesketh,⁶³ M. D. Hildreth,⁵⁵ R. Hirosky,⁸¹ T. Hoang,⁴⁹ J. D. Hobbs,⁷² B. Hoeneisen,¹² M. Hohlfeld,²² S. Hossain,⁷⁵ P. Houben,³⁴ Y. Hu,⁷² Z. Hubacek,¹⁰ V. Hynek,⁹ I. Iashvili,⁶⁹ R. Illingworth,⁵⁰ A. S. Ito,⁵⁰ S. Jabeen,⁶² M. Jaffré,¹⁶ S. Jain,⁷⁵ K. Jakobs,²³ C. Jarvis,⁶¹ R. Jesik,⁴³ K. Johns,⁴⁵ C. Johnson,⁷⁰ M. Johnson,⁵⁰ D. Johnston,⁶⁷ A. Jonckheere,⁵⁰ P. Jonsson,⁴³ A. Juste,⁵⁰ E. Kajfasz,¹⁵ D. Karmanov,³⁸ P. A. Kasper,⁵⁰ I. Katsanos,⁷⁰ V. Kaushik,⁷⁸ R. Kehoe,⁷⁹ S. Kermiche,¹⁵ N. Khalatyan,⁵⁰ A. Khanov,⁷⁶ A. Kharchilava,⁶⁹ Y. N. Kharzheev,³⁶ D. Khatidze,⁷⁰ T. J. Kim,³¹ M. H. Kirby,⁵³ M. Kirsch,²¹ B. Klima,⁵⁰ J. M. Kohli,²⁷ J.-P. Konrath,²³ A. V. Kozelov,³⁹ J. Kraus,⁶⁵ T. Kuhl,²⁴ A. Kumar,⁶⁹ A. Kupco,¹¹ T. Kurča,²⁰ V. A. Kuzmin,³⁸ J. Kvita,⁹ F. Lacroix,¹³ D. Lam,⁵⁵ S. Lammers,⁷⁰ G. Landsberg,⁷⁷ P. Lebrun,²⁰ W. M. Lee,⁵⁰ A. Leflat,³⁸ J. Lellouch,¹⁷ J. Li,^{78,§§} L. Li,⁴⁸ Q. Z. Li,⁵⁰ S. M. Lietti,⁵ J. K. Lim,³¹ J. G. R. Lima,⁵² D. Lincoln,⁵⁰ J. Linnemann,⁶⁵ V. V. Lipaev,³⁹ R. Lipton,⁵⁰ Y. Liu,⁷ Z. Liu,⁶ A. Lobodenko,⁴⁰ M. Lokajicek,¹¹ P. Love,⁴² H. J. Lubatti,⁸² R. Luna-Garcia,^{33,||} A. L. Lyon,⁵⁰ A. K. A. Maciel,² D. Mackin,⁸⁰ R. J. Madaras,⁴⁶ P. Mättig,²⁶ A. Magerkurth,⁶⁴ P. K. Mal,⁸² H. B. Malbouisson,³ S. Malik,⁶⁷ V. L. Malyshev,³⁶ Y. Maravin,⁵⁹ B. Martin,¹⁴ R. McCarthy,⁷² M. M. Meijer,³⁵ A. Melnitchouk,⁶⁶ L. Mendoza,⁸ P. G. Mercadante,⁵ M. Merkin,³⁸ K. W. Merritt,⁵⁰ A. Meyer,²¹ J. Meyer,^{22,§} J. Mitrevski,⁷⁰ R. K. Mommsen,⁴⁴ N. K. Mondal,²⁹ R. W. Moore,⁶ T. Moulik,⁵⁸ G. S. Muanza,¹⁵ M. Mulhearn,⁷⁰ O. Mundal,²² L. Mundim,³ E. Nagy,¹⁵ M. Naimuddin,⁵⁰ M. Narain,⁷⁷ H. A. Neal,⁶⁴ J. P. Negret,⁸ P. Neustroev,⁴⁰ H. Nilsen,²³ H. Nogima,³ S. F. Novaes,⁵ T. Nunnemann,²⁵ D. C. O'Neil,⁶ G. Odrant,⁴⁰ C. Ochando,¹⁶ D. Onoprienko,⁵⁹ N. Oshima,⁵⁰ N. Osman,⁴³ J. Osta,⁵⁵ R. Otec,¹⁰ G. J. Otero y Garzón,⁵⁰ M. Owen,⁴⁴ P. Padley,⁸⁰ M. Pangilinan,⁷⁷ N. Parashar,⁵⁶ S.-J. Park,^{22,§} S. K. Park,³¹ J. Parsons,⁷⁰ R. Partridge,⁷⁷ N. Parua,⁵⁴ A. Patwa,⁷³ G. Pawloski,⁸⁰ B. Penning,²³ M. Perfilov,³⁸ K. Peters,⁴⁴ Y. Peters,²⁶ P. Pétrouff,¹⁶ M. Petteni,⁴³ R. Piegaia,¹ J. Piper,⁶⁵ M.-A. Pleier,²² P. L. M. Podesta-Lerma,^{33,¶} V. M. Podstavkov,⁵⁰ Y. Pogorelov,⁵⁵ M.-E. Pol,² P. Polozov,³⁷ B. G. Pope,⁶⁵ A. V. Popov,³⁹ C. Potter,⁶ W. L. Prado da Silva,³ H. B. Prosper,⁴⁹ S. Protopopescu,⁷³ J. Qian,⁶⁴ A. Quadt,^{22,§} B. Quinn,⁶⁶ A. Rakitine,⁴² M. S. Rangel,² K. Ranjan,²⁸ P. N. Ratoff,⁴² P. Renkel,⁷⁹ P. Rich,⁴⁴ M. Rijssenbeek,⁷² I. Ripp-Baudot,¹⁹ F. Rizatdinova,⁷⁶ S. Robinson,⁴³ R. F. Rodrigues,³ M. Rominsky,⁷⁵ C. Royon,¹⁸ P. Rubinov,⁵⁰

R. Ruchti,⁵⁵ G. Safronov,³⁷ G. Sajot,¹⁴ A. Sánchez-Hernández,³³ M. P. Sanders,¹⁷ B. Sanghi,⁵⁰ G. Savage,⁵⁰ L. Sawyer,⁶⁰ T. Scanlon,⁴³ D. Schaile,²⁵ R. D. Schamberger,⁷² Y. Scheglov,⁴⁰ H. Schellman,⁵³ T. Schliephake,²⁶ S. Schlobohm,⁸² C. Schwanenberger,⁴⁴ A. Schwartzman,⁶⁸ R. Schwienhorst,⁶⁵ J. Sekaric,⁴⁹ H. Severini,⁷⁵ E. Shabalina,⁵¹ M. Shamim,⁵⁹ V. Shary,¹⁸ A. A. Shchukin,³⁹ R. K. Shivpuri,²⁸ V. Siccaldi,¹⁹ V. Simak,¹⁰ V. Sirotenko,⁵⁰ P. Skubic,⁷⁵ P. Slattery,⁷¹ D. Smirnov,⁵⁵ G. R. Snow,⁶⁷ J. Snow,⁷⁴ S. Snyder,⁷³ S. Söldner-Rembold,⁴⁴ L. Sonnenschein,¹⁷ A. Sopczak,⁴² M. Sosebee,⁷⁸ K. Soustruznik,⁹ B. Spurlock,⁷⁸ J. Stark,¹⁴ V. Stolin,³⁷ D. A. Stoyanova,³⁹ J. Strandberg,⁶⁴ S. Strandberg,⁴¹ M. A. Strang,⁶⁹ E. Strauss,⁷² M. Strauss,⁷⁵ R. Ströhmer,²⁵ D. Strom,⁵³ L. Stutte,⁵⁰ S. Sumowidagdo,⁴⁹ P. Svoisky,³⁵ A. Sznajder,³ A. Tanasijczuk,¹ W. Taylor,⁶ B. Tiller,²⁵ F. Tissandier,¹³ M. Titov,¹⁸ V. V. Tokmenin,³⁶ I. Torchiani,²³ D. Tsybychev,⁷² B. Tuchming,¹⁸ C. Tully,⁶⁸ P. M. Tuts,⁷⁰ R. Unalan,⁶⁵ L. Uvarov,⁴⁰ S. Uvarov,⁴⁰ S. Uzunyan,⁵² B. Vachon,⁶ P. J. van den Berg,³⁴ R. Van Kooten,⁵⁴ W. M. van Leeuwen,³⁴ N. Varelas,⁵¹ E. W. Varnes,⁴⁵ I. A. Vasilyev,³⁹ P. Verdier,²⁰ L. S. Vertogradov,³⁶ M. Verzocchi,⁵⁰ D. Vilanova,¹⁸ F. Villeneuve-Seguié,⁴³ P. Vint,⁴³ P. Vokac,¹⁰ M. Voutilainen,^{67,**} R. Wagner,⁶⁸ H. D. Wahl,⁴⁹ M. H. L. S. Wang,⁵⁰ J. Warchol,⁵⁵ G. Watts,⁸² M. Wayne,⁵⁵ G. Weber,²⁴ M. Weber,^{50,††} L. Welty-Rieger,⁵⁴ A. Wenger,^{23,‡‡} N. Wermes,²² M. Wetstein,⁶¹ A. White,⁷⁸ D. Wicke,²⁶ M. R. J. Williams,⁴² G. W. Wilson,⁵⁸ S. J. Wimpenny,⁴⁸ M. Wobisch,⁶⁰ D. R. Wood,⁶³ T. R. Wyatt,⁴⁴ Y. Xie,⁷⁷ C. Xu,⁶⁴ S. Yacoub,⁵³ R. Yamada,⁵⁰ W.-C. Yang,⁴⁴ T. Yasuda,⁵⁰ Y. A. Yatsunenkov,³⁶ H. Yin,⁷ K. Yip,⁷³ H. D. Yoo,⁷⁷ S. W. Youn,⁵³ J. Yu,⁷⁸ C. Zeitnitz,²⁶ S. Zelitch,⁸¹ T. Zhao,⁸² B. Zhou,⁶⁴ J. Zhu,⁷² M. Zielinski,⁷¹ D. Zieminska,⁵⁴ A. Zieminski,^{54,§§} L. Zivkovic,⁷⁰ V. Zutshi,⁵² and E. G. Zverev³⁸

(D0 Collaboration)

¹Universidad de Buenos Aires, Buenos Aires, Argentina²LAFEX, Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil³Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil⁴Universidade Federal do ABC, Santo André, Brazil⁵Instituto de Física Teórica, Universidade Estadual Paulista, São Paulo, Brazil⁶University of Alberta, Edmonton, Alberta, Canada,

Simon Fraser University, Burnaby, British Columbia, Canada,

York University, Toronto, Ontario, Canada,

and McGill University, Montreal, Quebec, Canada

⁷University of Science and Technology of China, Hefei, People's Republic of China⁸Universidad de los Andes, Bogotá, Colombia⁹Center for Particle Physics, Charles University, Prague, Czech Republic¹⁰Czech Technical University, Prague, Czech Republic¹¹Center for Particle Physics, Institute of Physics, Academy of Sciences of the Czech Republic, Prague, Czech Republic¹²Universidad San Francisco de Quito, Quito, Ecuador¹³LPC, Université Blaise Pascal, CNRS/IN2P3, Clermont, France¹⁴LPSC, Université Joseph Fourier Grenoble 1, CNRS/IN2P3, Institut National Polytechnique de Grenoble, Grenoble, France¹⁵CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille, France¹⁶LAL, Université Paris-Sud, IN2P3/CNRS, Orsay, France¹⁷LPNHE, IN2P3/CNRS, Universités Paris VI and VII, Paris, France¹⁸CEA, Irfu, SPP, Saclay, France¹⁹IPHC, Université Louis Pasteur, CNRS/IN2P3, Strasbourg, France²⁰IPNL, Université Lyon 1, CNRS/IN2P3, Villeurbanne, France and Université de Lyon, Lyon, France²¹III. Physikalisches Institut A, RWTH Aachen University, Aachen, Germany²²Physikalisches Institut, Universität Bonn, Bonn, Germany²³Physikalisches Institut, Universität Freiburg, Freiburg, Germany²⁴Institut für Physik, Universität Mainz, Mainz, Germany²⁵Ludwig-Maximilians-Universität München, München, Germany²⁶Fachbereich Physik, University of Wuppertal, Wuppertal, Germany²⁷Panjab University, Chandigarh, India²⁸Delhi University, Delhi, India²⁹Tata Institute of Fundamental Research, Mumbai, India³⁰University College Dublin, Dublin, Ireland³¹Korea Detector Laboratory, Korea University, Seoul, Korea³²SungKyunKwan University, Suwon, Korea³³CINVESTAV, Mexico City, Mexico³⁴FOM-Institute NIKHEF and University of Amsterdam/NIKHEF, Amsterdam, The Netherlands

- ³⁵Radboud University Nijmegen/NIKHEF, Nijmegen, The Netherlands
³⁶Joint Institute for Nuclear Research, Dubna, Russia
³⁷Institute for Theoretical and Experimental Physics, Moscow, Russia
³⁸Moscow State University, Moscow, Russia
³⁹Institute for High Energy Physics, Protvino, Russia
⁴⁰Petersburg Nuclear Physics Institute, St. Petersburg, Russia
⁴¹Lund University, Lund, Sweden, Royal Institute of Technology and Stockholm University, Stockholm, Sweden, and Uppsala University, Uppsala, Sweden
⁴²Lancaster University, Lancaster, United Kingdom
⁴³Imperial College, London, United Kingdom
⁴⁴University of Manchester, Manchester, United Kingdom
⁴⁵University of Arizona, Tucson, Arizona 85721, USA
⁴⁶Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA
⁴⁷California State University, Fresno, California 93740, USA
⁴⁸University of California, Riverside, California 92521, USA
⁴⁹Florida State University, Tallahassee, Florida 32306, USA
⁵⁰Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA
⁵¹University of Illinois at Chicago, Chicago, Illinois 60607, USA
⁵²Northern Illinois University, DeKalb, Illinois 60115, USA
⁵³Northwestern University, Evanston, Illinois 60208, USA
⁵⁴Indiana University, Bloomington, Indiana 47405, USA
⁵⁵University of Notre Dame, Notre Dame, Indiana 46556, USA
⁵⁶Purdue University Calumet, Hammond, Indiana 46323, USA
⁵⁷Iowa State University, Ames, Iowa 50011, USA
⁵⁸University of Kansas, Lawrence, Kansas 66045, USA
⁵⁹Kansas State University, Manhattan, Kansas 66506, USA
⁶⁰Louisiana Tech University, Ruston, Louisiana 71272, USA
⁶¹University of Maryland, College Park, Maryland 20742, USA
⁶²Boston University, Boston, Massachusetts 02215, USA
⁶³Northeastern University, Boston, Massachusetts 02115, USA
⁶⁴University of Michigan, Ann Arbor, Michigan 48109, USA
⁶⁵Michigan State University, East Lansing, Michigan 48824, USA
⁶⁶University of Mississippi, University, Mississippi 38677, USA
⁶⁷University of Nebraska, Lincoln, Nebraska 68588, USA
⁶⁸Princeton University, Princeton, New Jersey 08544, USA
⁶⁹State University of New York, Buffalo, New York 14260, USA
⁷⁰Columbia University, New York, New York 10027, USA
⁷¹University of Rochester, Rochester, New York 14627, USA
⁷²State University of New York, Stony Brook, New York 11794, USA
⁷³Brookhaven National Laboratory, Upton, New York 11973, USA
⁷⁴Langston University, Langston, Oklahoma 73050, USA
⁷⁵University of Oklahoma, Norman, Oklahoma 73019, USA
⁷⁶Oklahoma State University, Stillwater, Oklahoma 74078, USA
⁷⁷Brown University, Providence, Rhode Island 02912, USA
⁷⁸University of Texas, Arlington, Texas 76019, USA
⁷⁹Southern Methodist University, Dallas, Texas 75275, USA
⁸⁰Rice University, Houston, Texas 77005, USA
⁸¹University of Virginia, Charlottesville, Virginia 22901, USA
⁸²University of Washington, Seattle, Washington 98195, USA
(Received 21 October 2008; published 21 April 2009)

We present first evidence for $WW + WZ$ production in lepton + jets final states at a hadron collider. The data correspond to 1.07 fb^{-1} of integrated luminosity collected with the D0 detector at the Fermilab Tevatron in $p\bar{p}$ collisions at $\sqrt{s} = 1.96 \text{ TeV}$. The observed cross section for $WW + WZ$ production is $20.2 \pm 4.5 \text{ pb}$, consistent with the standard model and more precise than previous measurements in fully leptonic final states. The probability that background fluctuations alone produce this excess is $< 5.4 \times 10^{-6}$, which corresponds to a significance of 4.4 standard deviations.

The production of vector-boson pairs in $p\bar{p}$ collisions (WW , WZ , or ZZ) provides important tests of the electro-weak sector of the standard model (SM). The next-to-leading-order (NLO) cross sections for WW and WZ production in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ GeV predicted by the SM are $\sigma(WW) = 12.4 \pm 0.8$ pb and $\sigma(WZ) = 3.7 \pm 0.3$ pb [1]. A discrepancy with this expectation or deviations in the predicted kinematic distributions could signal the presence of new physics, e.g., originating from anomalous trilinear gauge boson couplings [2]. The production of two weak bosons is also relevant to searches for the Higgs boson or for new particles in extensions of the SM. Production of WW and WZ in $p\bar{p}$ collisions at the Fermilab Tevatron Collider has thus far been observed only in fully leptonic decay modes [3,4]. Previous searches for WW and WZ in lepton + jets final states [5,6], which benefit from a higher branching ratio relative to fully leptonic channels, were hindered by large backgrounds from jets produced in association with a W boson (W + jets).

In this Letter we report first evidence from a hadron collider for the production of a W boson that decays leptonically, associated with a second vector boson V ($V = W$ or Z) that decays into $q\bar{q}$ ($WV \rightarrow \ell\nu q\bar{q}$; $\ell = e, \mu$). The limited dijet mass resolution ($\approx 18\%$ for dijets from W/Z decays) results in a significant overlap of the $W \rightarrow q\bar{q}$ and $Z \rightarrow q\bar{q}$ dijet mass peaks. We therefore consider WW and WZ simultaneously, assuming the ratio of their cross sections as predicted by the SM. The use of improved multivariate event classification and new statistical techniques [7], as well as an increased integrated luminosity, make the WV signal in lepton + jets final states more distinguishable from the W + jets background and more accessible to measurement than in the past [5,6]. This analysis also provides a valuable proving ground for such advanced techniques, now ubiquitous in Higgs searches at the Tevatron.

We analyze 1.07 fb^{-1} of data collected with the D0 detector [8] at a center-of-mass energy of 1.96 TeV at the Tevatron. Candidate $e\nu q\bar{q}$ events must pass a trigger based on a single electron or electron + jet(s) requirement that has an efficiency of $98_{-3}^{+2}\%$. A suite of triggers for $\mu\nu q\bar{q}$ candidate events achieves an efficiency of $>95\%$ at 95% confidence level.

To select $WV \rightarrow \ell\nu q\bar{q}$ candidates, we require a single reconstructed lepton (electron or muon) [9] with transverse momentum $p_T > 20$ GeV and pseudorapidity $|\eta| < 1.1(2.0)$ for electrons (muons), the imbalance in transverse energy to be $E_T > 20$ GeV, and at least two jets [10] with $p_T > 20$ GeV and $|\eta| < 2.5$. The jet of highest p_T must have $p_T > 30$ GeV. To reduce the background from processes that do not contain $W \rightarrow \ell\nu$, we require a “transverse” mass [11] of $M_T^{\ell\nu} > 35$ GeV. The lepton must be spatially matched to a track reconstructed in the central tracker that originates from the primary vertex. Electrons (muons) must be isolated from other particles in the calorimeter (and central tracker) [12].

Signal and background processes containing charged leptons are modeled via Monte Carlo (MC) simulation. The signal includes all possible W and Z decays, including their decays to leptons. The diboson signal (WW and WZ) is generated with PYTHIA [13] using CTEQ6L parton distribution functions (PDFs). The fixed-order matrix element (FOME) generator ALPGEN [14] with CTEQ6L1 PDFs is used to generate W + jets, Z + jets, and $t\bar{t}$ events to leading order at the parton level. The FOME generator COMPHEP [15] is used to produce single top-quark MC samples. ALPGEN and COMPHEP are interfaced to PYTHIA for subsequent parton showering and hadronization. All simulated events undergo a GEANT-based [16] detector simulation and are reconstructed using the same programs as used for D0 data. The MC samples are normalized using next-to-leading-order (NLO) or next-to-next-to-leading-order predictions for SM cross sections, except W + jets which is scaled to the data.

The probability for multijet events with misidentified leptons to pass all selection requirements is small; however, because of the copious production of multijet events, the background from this source cannot be ignored. For $\mu\nu q\bar{q}$, the multijet background is modeled with data that fail the muon isolation requirements, but pass all other selections. The normalization is determined from a fit to the $M_T^{\ell\nu}$ distribution. For $e\nu q\bar{q}$, the multijet background is estimated using a “loose-but-not-tight” data sample obtained by selecting events that pass loosened electron quality requirements, but fail the tight electron quality criteria [9]. This sample is normalized by the probability for a jet that passes the “loose” electron requirements to also pass the tight requirement. Both $\mu\nu q\bar{q}$ and $e\nu q\bar{q}$ multijet samples are corrected for contributions from all processes modeled through MC calculations.

Accurate modeling of the selected events is vital. The dominant background is W + jets, and the modeling of ALPGEN W + jets and sources of uncertainty are therefore studied in great detail. Comparison of ALPGEN with other generators and with data shows discrepancies [17] in jet η and dijet angular separation. Data are used to correct these quantities in the ALPGEN W + jets and Z + jets samples. The possible bias in this procedure from the presence of

TABLE I. Measured number of events for the signal and each background after the combined fit (with total uncertainties determined from the fit) and the number observed in data.

	$e\nu q\bar{q}$ channel	$\mu\nu q\bar{q}$ channel
Diboson signal	436 ± 36	527 ± 43
W + jets	10100 ± 500	11910 ± 590
Z + jets	387 ± 61	1180 ± 180
$t\bar{t}$ + single top	436 ± 57	426 ± 54
Multijet	1100 ± 200	328 ± 83
Total predicted	12460 ± 550	14370 ± 620
Data	12473	14392

TABLE II. The signal cross section extracted from a simultaneous fit of the WV cross section and the normalization factor for $W + \text{jets}$. Also given are expected and observed p values obtained by comparing the measurement with pseudoexperiments assuming no signal and the corresponding significance in number of standard deviations (s.d.) for a one-sided Gaussian integral.

Channel	Fitted signal σ (pb)	Expected p -value (significance)	Observed p -value (significance)
$evq\bar{q}$ RF Output	$18.0 \pm 3.7(\text{stat}) \pm 5.2(\text{syst}) \pm 1.1(\text{lum})$	6.8×10^{-3} (2.5 s.d.)	3.2×10^{-3} (2.7 s.d.)
$\mu\nu q\bar{q}$ RF Output	$22.8 \pm 3.3(\text{stat}) \pm 4.9(\text{syst}) \pm 1.4(\text{lum})$	1.8×10^{-3} (2.9 s.d.)	5.2×10^{-5} (3.9 s.d.)
Combined RF Output	$20.2 \pm 2.5(\text{stat}) \pm 3.6(\text{syst}) \pm 1.2(\text{lum})$	1.5×10^{-4} (3.6 s.d.)	5.4×10^{-6} (4.4 s.d.)
Combined Dijet Mass	$18.5 \pm 2.8(\text{stat}) \pm 4.9(\text{syst}) \pm 1.1(\text{lum})$	1.7×10^{-3} (2.9 s.d.)	4.4×10^{-4} (3.3 s.d.)

the diboson signal in data is small, but is nevertheless taken into account via a systematic uncertainty. Systematic effects on the differential distributions of the ALPGEN $W + \text{jets}$ and $Z + \text{jets}$ MC events from changes of the renormalization and factorization scales and of the parameters used in the MLM parton-jet matching algorithm [18] are also considered. Uncertainties on PDFs, as well as uncertainties from object reconstruction and identification, are evaluated for all MC samples. We consider the effect of systematic uncertainty both on the normalization and on the shape of differential distributions for the signal and backgrounds [19].

The signal and the backgrounds are further separated using a multivariate classifier to combine information from several kinematic variables. This analysis uses a Random Forest (RF) classifier [20,21]. Thirteen well-modeled kinematic variables [19] that demonstrate a difference in probability density between signal and at least one of the backgrounds, such as dijet mass and \cancel{E}_T , are used as inputs to the RF. The RF is trained using half of each MC sample. The other halves, along with the multijet background samples, are then evaluated by the RF and used in the measurement.

The signal cross section is determined from a fit of signal and background RF templates to the data by minimizing a Poisson χ^2 function with respect to variations in the systematic uncertainties [7]. The magnitude of systematic uncertainties is effectively constrained by the regions of the RF distribution with low signal over background. A Gaussian prior is used for each systematic uncertainty. Different uncertainties are assumed to be mutually independent, but those common to multiple samples or lepton channels are assumed to be 100% correlated.

The fit simultaneously varies the WV and $W + \text{jets}$ contributions, thereby also determining the normalization factor for the $W + \text{jets}$ MC sample. This obviates the need for using the predicted ALPGEN cross section, and provides a more rigorous approach that incorporates an unbiased uncertainty from $W + \text{jets}$ when extracting the WV cross section. The normalization factor from the fit for the $W + \text{jets}$ component is 1.53 ± 0.13 , similar to the expected ratio of NLO to LO cross sections [22]. The measured yields for the signal and each background are given in Table I. Table II contains the measured WV cross section

for each channel, separately and combined, showing consistent results between channels and the SM prediction of $\sigma(WV) = 16.1 \pm 0.9$ pb [1]. The combined fit yields a cross section of $20.2 \pm 2.5(\text{stat}) \pm 3.6(\text{syst}) \pm 1.2(\text{lum})$ pb. The RF output distributions following the combined fit are shown in Fig. 1, along with comparisons of consistency between the background-subtracted data




FIG. 1 (color online). (a) The RF output distribution from the combined $evq\bar{q}$ and $\mu\nu q\bar{q}$ channels for data and MC predictions following a fit of the MC calculations to data. (b) A comparison of the extracted signal (filled histogram) to background-subtracted data (points), along with the ± 1 standard deviation (s.d.) systematic uncertainty on the background. The residual distance between the data points and the extracted signal, divided by the total uncertainty, is given at the bottom.




FIG. 2 (color online). (a) The dijet mass distribution from the combined $evq\bar{q}$ and $\mu\nu q\bar{q}$ channels for data and MC predictions following the fit to the RF output. (b) A comparison of the extracted signal (filled histogram) to background-subtracted data (points), along with the ± 1 standard deviation (s.d.) systematic uncertainty on the background. The residual distance between the data points and the extracted signal, divided by the total uncertainty, is given at the bottom.

and the extracted signal. Figure 2 shows analogous plots for the dijet mass after the combined fit to the RF output. The dominant systematic uncertainties arise from the modeling of the W + jets background and the jet energy scale, contributing 2.4 pb and 1.9 pb to the total systematic uncertainty [19], respectively. The position of the dijet mass peaks in data and MC calculations are consistent within one-half standard deviation, which includes the relative data or MC uncertainty in energy scale. As a cross check, we also perform the measurement using only the dijet mass distribution. The result, also given in Table II, although less precise, is consistent with that obtained using the RF output.

The significance of the measurement is obtained via fits of the signal + background hypothesis to pseudodata samples drawn from the background-only hypothesis [23]. The observed (or expected) significance corresponds to the fraction of outcomes that yield a WV cross section at

least as large as that measured in data (as predicted by the SM). The probabilities that background fluctuations could produce the expected and observed signal in each channel (p values), separately and combined, are shown in Table II, along with their corresponding significance (equivalent one-sided Gaussian probabilities). The χ^2 fit with respect to variations in the systematic uncertainties [7] results in an improvement of the expected significance of the result from 2.4 (1.6) to 3.6 (2.9) standard deviations when using the RF output (dijet mass) discriminant.

In summary, we measure $\sigma(WV) = 20.2 \pm 4.5$ pb (with $V = W$ or Z) in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV. The probability that the backgrounds fluctuate to give an excess as large as observed in data is $< 5.4 \times 10^{-6}$, corresponding to a significance of 4.4 standard deviations. This represents the first evidence for WV production in lepton + jets events at a hadron collider. The result is more precise than previous independent measurements of WW and WZ yields in fully leptonic final states [3,4] and consistent with the SM prediction of $\sigma(WV) = 16.1 \pm 0.9$ pb [1]. This work clearly demonstrates the ability of the D0 experiment to isolate a small signal in a large background in a final state of direct relevance to searches for a low mass Higgs boson, and thereby validates the analytical methods used in searches for Higgs bosons at the Tevatron [24].

We thank the staffs at Fermilab and collaborating institutions, and acknowledge support from the DOE and NSF (USA); CEA and CNRS/IN2P3 (France); FASI, Rosatom and RFBR (Russia); CNPq, FAPERJ, FAPESP and FUNDUNESP (Brazil); DAE and DST (India); Colciencias (Colombia); CONACyT (Mexico); KRF and KOSEF (Korea); CONICET and UBACyT (Argentina); FOM (The Netherlands); STFC (United Kingdom); MSMT and GACR (Czech Republic); CRC Program, CFI, NSERC and WestGrid Project (Canada); BMBF and DFG (Germany); SFI (Ireland); The Swedish Research Council (Sweden); CAS and CNSF (China); and the Alexander von Humboldt Foundation (Germany).

*Visitor from Augustana College, Sioux Falls, SD, USA.
[†]Visitor from Rutgers University, Piscataway, NJ, USA.
[‡]Visitor from The University of Liverpool, Liverpool, UK.
[§]Visitor from II. Physikalisches Institut, Georg-August-University, Göttingen, Germany.
^{||}Visitor from Centro de Investigacion en Computacion - IPN, Mexico City, Mexico.
[¶]Visitor from ECFM, Universidad Autonoma de Sinaloa, Culiacán, Mexico.
^{**}Visitor from Helsinki Institute of Physics, Helsinki, Finland.
^{††}Visitor from Universität Bern, Bern, Switzerland.
^{‡‡}Visitor from Universität Zürich, Zürich, Switzerland.
^{§§}Deceased.

- [1] J. M. Campbell and R. K. Ellis, *Phys. Rev. D* **60**, 113006 (1999). Cross sections were calculated with the same parameter values given in the paper, except with $\sqrt{s} = 1.96$ TeV.
- [2] K. Hagiwara, S. Ishihara, R. Szalapski, and D. Zeppenfeld, *Phys. Rev. D* **48**, 2182 (1993).
- [3] V. M. Abazov *et al.* (D0 Collaboration), *Phys. Rev. Lett.* **94**, 151801 (2005); *Phys. Rev. D* **76**, 111104(R) (2007); *Phys. Rev. Lett.* **101**, 171803 (2008).
- [4] D. Acosta *et al.* (CDF Collaboration), *Phys. Rev. Lett.* **94**, 211801 (2005); A. Abulencia *et al.*, *Phys. Rev. Lett.* **98**, 161801 (2007); T. Aaltonen *et al.*, *Phys. Rev. Lett.* **100**, 201801 (2008).
- [5] B. Abbott *et al.* (D0 Collaboration), *Phys. Rev. D* **62**, 052005 (2000).
- [6] T. Aaltonen *et al.* (CDF Collaboration), *Phys. Rev. D* **76**, 111103(R) (2007).
- [7] W. Fisher, FERMILAB Report No. FERMILAB-TM-2386-E, 2006.
- [8] B. Abbott *et al.* (D0 Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **565**, 463 (2006).
- [9] V. M. Abazov *et al.* (D0 Collaboration), *Phys. Lett. B* **626**, 45 (2005).
- [10] G. C. Blazey *et al.*, arXiv:hep-ex/0005012. The seeded cone algorithm with radius 0.5 was used.
- [11] J. Smith, W. L. van Neerven, and J. A. M. Vermaseren, *Phys. Rev. Lett.* **50**, 1738 (1983).
- [12] V. M. Abazov *et al.* (D0 Collaboration), *Phys. Rev. Lett.* **100**, 139901 (2008).
- [13] T. Sjöstrand *et al.*, *Comput. Phys. Commun.* **135**, 238 (2001). Verison 6.3 was used.
- [14] M. L. Mangano *et al.*, *J. High Energy Phys.* 07 (2003) 001. Version 2.05 was used.
- [15] A. Pukhov *et al.*, arXiv:hep-ph/9908288.
- [16] R. Brun and F. Carminati, CERN Program Library Long Writeup Report No. W5013, 1993.
- [17] J. Alwall *et al.*, *Eur. Phys. J. C* **53**, 473 (2008).
- [18] S. Höche *et al.*, arXiv:hep-ph/0602031.
- [19] See EPAPS Document No. E-PRLTAO-102-001919 for supplementary material. For more information on EPAPS, see <http://www.aip.org/pubservs/epaps.html>.
- [20] L. Breiman, *Mach. Learn.* **45**, 5 (2001).
- [21] I. Narsky, arXiv:physics/0507143.
- [22] J. M. Campbell and R. K. Ellis, *Phys. Rev. D* **65**, 113007 (2002).
- [23] V. M. Abazov *et al.* (D0 Collaboration), *Phys. Rev. D* **78**, 012005 (2008).
- [24] TEVNPH Working Group, for the CDF Collaboration and D0 Collaboration, arXiv:0804.3423.