

2009

Observation of Single Top-Quark Production

V. M. Abazov

Joint Institute for Nuclear Research, Dubna, Russia

Kenneth A. Bloom

University of Nebraska - Lincoln, kbloom2@unl.edu

Gregory Snow

University of Nebraska - Lincoln, gsnow1@unl.edu

D0 Collaboration

Follow this and additional works at: <http://digitalcommons.unl.edu/physicsbloom>



Part of the [Physics Commons](#)

Abazov, V. M.; Bloom, Kenneth A.; Snow, Gregory; and Collaboration, D0, "Observation of Single Top-Quark Production" (2009).
Kenneth Bloom Publications. 291.

<http://digitalcommons.unl.edu/physicsbloom/291>

This Article is brought to you for free and open access by the Research Papers in Physics and Astronomy at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Kenneth Bloom Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.



Observation of Single Top-Quark Production

V. M. Abazov,³⁶ B. Abbott,⁷⁴ M. Abolins,⁶⁴ B. S. Acharya,²⁹ M. Adams,⁵⁰ T. Adams,⁴⁸ E. Aguilo,⁶ M. Ahsan,⁵⁸ G. D. Alexeev,³⁶ G. Alkhalaf,⁴⁰ A. Alton,^{64,*} G. Alverson,⁶² G. A. Alves,² L. S. Ancu,³⁵ T. Andeen,⁵² M. S. Anzels,⁵² M. Aoki,⁴⁹ Y. Arnoud,¹⁴ M. Arov,⁵⁹ M. Arthaud,¹⁸ A. Askew,^{48,†} B. Åsman,⁴¹ O. Atramentov,^{48,†} C. Avila,⁸ J. BackusMayes,⁸¹ F. Badaud,¹³ L. Bagby,⁴⁹ B. Baldin,⁴⁹ D. V. Bandurin,⁵⁸ P. Banerjee,²⁹ S. Banerjee,²⁹ E. Barberis,⁶² A.-F. Barfuss,¹⁵ P. Bargassa,⁷⁹ P. Baringer,⁵⁷ J. Barreto,² J. F. Bartlett,⁴⁹ U. Bassler,¹⁸ D. Bauer,⁴³ S. Beale,⁶ A. Bean,⁵⁷ M. Begalli,³ M. Begel,⁷² C. Belanger-Champagne,⁴¹ L. Bellantoni,⁴⁹ A. Bellavance,⁴⁹ J. A. Benitez,⁶⁴ S. B. Beri,²⁷ G. Bernardi,¹⁷ R. Bernhard,²³ I. Bertram,⁴² M. Besançon,¹⁸ R. Beuselinck,⁴³ V. A. Bezzubov,³⁹ P. C. Bhat,⁴⁹ V. Bhatnagar,²⁷ G. Blazey,⁵¹ S. Blessing,⁴⁸ K. Bloom,⁶⁶ A. Boehnlein,⁴⁹ D. Boline,⁶¹ T. A. Bolton,⁵⁸ E. E. Boos,³⁸ G. Borissov,⁴² T. Bose,⁷⁶ A. Brandt,⁷⁷ R. Brock,⁶⁴ G. Brooijmans,⁶⁹ A. Bross,⁴⁹ D. Brown,¹⁹ X. B. Bu,⁷ N. J. Buchanan,⁴⁸ D. Buchholz,⁵² M. Buehler,⁸⁰ V. Buescher,²² V. Bunichev,³⁸ S. Burdin,^{42,‡} T. H. Burnett,⁸¹ C. P. Buszello,⁴³ P. Calfayan,²⁵ B. Calpas,¹⁵ S. Calvet,¹⁶ J. Cammin,⁷⁰ M. A. Carrasco-Lizarraga,³³ E. Carrera,⁴⁸ W. Carvalho,³ B. C. K. Casey,⁴⁹ H. Castilla-Valdez,³³ S. Chakrabarti,⁷¹ D. Chakraborty,⁵¹ K. M. Chan,⁵⁴ A. Chandra,⁴⁷ E. Cheu,⁴⁵ D. K. Cho,⁶¹ S. Choi,³² B. Choudhary,²⁸ L. Christofek,⁷⁶ T. Christoudias,⁴³ S. Cihangir,⁴⁹ D. Claes,⁶⁶ J. Clutter,⁵⁷ Y. Coadou,^{6,§} M. Cooke,⁴⁹ W. E. Cooper,⁴⁹ M. Corcoran,⁷⁹ F. Couderc,¹⁸ M.-C. Cousinou,¹⁵ S. Crépe-Renaudin,¹⁴ V. Cuplov,⁵⁸ D. Cutts,⁷⁶ M. Ćwiok,³⁰ A. Das,⁴⁵ G. Davies,⁴³ K. De,⁷⁷ S. J. de Jong,³⁵ E. De La Cruz-Burelo,³³ K. DeVaughan,⁶⁶ F. Déliot,¹⁸ M. Demarteau,⁴⁹ R. Demina,⁷⁰ D. Denisov,⁴⁹ S. P. Denisov,³⁹ S. Desai,⁴⁹ H. T. Diehl,⁴⁹ M. Diesburg,⁴⁹ A. Dominguez,⁶⁶ T. Dorland,⁸¹ A. Dubey,²⁸ L. V. Dudko,³⁸ L. Dufлот,¹⁶ D. Duggan,⁴⁸ A. Duperrin,¹⁵ S. Dutt,²⁷ A. Dyshkant,⁵¹ M. Eads,⁶⁶ D. Edmunds,⁶⁴ J. Ellison,⁴⁷ V. D. Elvira,⁴⁹ Y. Enari,⁷⁶ S. Eno,⁶⁰ P. Ermolov,^{38,|||} M. Escalier,¹⁵ H. Evans,⁵³ A. Evdokimov,⁷² V. N. Evdokimov,³⁹ A. V. Ferapontov,⁵⁸ T. Ferbel,^{61,70} F. Fiedler,²⁴ F. Filthaut,³⁵ W. Fisher,⁴⁹ H. E. Fisk,⁴⁹ M. Fortner,⁵¹ H. Fox,⁴² S. Fu,⁴⁹ S. Fuess,⁴⁹ T. Gadfort,⁶⁹ C. F. Galea,³⁵ A. Garcia-Bellido,⁷⁰ V. Gavrilov,³⁷ P. Gay,¹³ W. Geist,¹⁹ W. Geng,^{15,64} C. E. Gerber,⁵⁰ Y. Gershtein,^{48,†} D. Gillberg,⁶ G. Ginther,⁷⁰ B. Gómez,⁸ A. Goussiou,⁸¹ P. D. Grannis,⁷¹ S. Greder,¹⁹ H. Greenlee,⁴⁹ Z. D. Greenwood,⁵⁹ E. M. Gregores,⁴ G. Grenier,²⁰ Ph. Gris,¹³ J.-F. Grivaz,¹⁶ A. Grohsjean,²⁵ S. Grünendahl,⁴⁹ M. W. Grünewald,³⁰ F. Guo,⁷¹ J. Guo,⁷¹ G. Gutierrez,⁴⁹ P. Gutierrez,⁷⁴ A. Haas,⁶⁹ N. J. Hadley,⁶⁰ P. Haefner,²⁵ S. Hagopian,⁴⁸ J. Haley,⁶⁷ I. Hall,⁶⁴ R. E. Hall,⁴⁶ L. Han,⁷ K. Harder,⁴⁴ A. Harel,⁷⁰ J. M. Hauptman,⁵⁶ J. Hays,⁴³ T. Hebbeker,²¹ D. Hedin,⁵¹ J. G. Hegeman,³⁴ A. P. Heinson,⁴⁷ U. Heintz,⁶¹ C. Hensel,^{22,||} K. Herner,⁶³ G. Hesketh,⁶² M. D. Hildreth,⁵⁴ R. Hirosky,⁸⁰ T. Hoang,⁴⁸ J. D. Hobbs,⁷¹ B. Hoeneisen,¹² M. Hohlfield,²² S. Hossain,⁷⁴ P. Houben,³⁴ Y. Hu,⁷¹ Z. Hubacek,¹⁰ N. Huske,¹⁷ V. Hynek,¹⁰ I. Iashvili,⁶⁸ R. Illingworth,⁴⁹ A. S. Ito,⁴⁹ S. Jabeen,⁶¹ M. Jaffré,¹⁶ S. Jain,⁷⁴ K. Jakobs,²³ D. Jamin,¹⁵ C. Jarvis,⁶⁰ R. Jesik,⁴³ K. Johns,⁴⁵ C. Johnson,⁶⁹ M. Johnson,⁴⁹ D. Johnston,⁶⁶ A. Jonckheere,⁴⁹ P. Jonsson,⁴³ A. Juste,⁴⁹ E. Kajfasz,¹⁵ D. Karmanov,³⁸ P. A. Kasper,⁴⁹ I. Katsanos,⁶⁶ V. Kaushik,⁷⁷ R. Kehoe,⁷⁸ S. Kermiche,¹⁵ N. Khalatyan,⁴⁹ A. Khanov,⁷⁵ A. Kharchilava,⁶⁸ Y. N. Kharzhev,³⁶ D. Khatidze,⁶⁹ T. J. Kim,³¹ M. H. Kirby,⁵² M. Kirsch,²¹ B. Klima,⁴⁹ J. M. Kohli,²⁷ J.-P. Konrath,²³ A. V. Kozelov,³⁹ J. Kraus,⁶⁴ T. Kuhl,²⁴ A. Kumar,⁶⁸ A. Kupco,¹¹ T. Kurča,²⁰ V. A. Kuzmin,³⁸ J. Kvita,⁹ F. Lacroix,¹³ D. Lam,⁵⁴ S. Lammers,⁵³ G. Landsberg,⁷⁶ P. Lebrun,²⁰ W. M. Lee,⁴⁹ A. Leflat,³⁸ J. Lellouch,^{17,|||} J. Li,⁷⁷ L. Li,⁴⁷ Q. Z. Li,⁴⁹ S. M. Lietti,⁵ J. K. Lim,³¹ D. Lincoln,⁴⁹ J. Linnemann,⁶⁴ V. V. Lipaev,³⁹ R. Lipton,⁴⁹ Y. Liu,⁷ Z. Liu,⁶ A. Lobodenko,⁴⁰ M. Lokajicek,¹¹ P. Love,⁴² H. J. Lubatti,⁸¹ R. Luna-Garcia,^{33,||} A. L. Lyon,⁴⁹ A. K. A. Maciel,² D. Mackin,⁷⁹ P. Mättig,²⁶ A. Magerkurth,⁶³ P. K. Mal,⁸¹ H. B. Malbouisson,³ S. Malik,⁶⁶ V. L. Malyshev,³⁶ Y. Maravin,⁵⁸ B. Martin,¹⁴ R. McCarthy,⁷¹ C. L. McGivern,⁵⁷ M. M. Meijer,³⁵ A. Melnitchouk,⁶⁵ L. Mendoza,⁸ P. G. Mercadante,⁵ M. Merkin,³⁸ K. W. Merritt,⁴⁹ A. Meyer,²¹ J. Meyer,^{22,||} J. Mitrevski,⁶⁹ R. K. Mommsen,⁴⁴ N. K. Mondal,²⁹ R. W. Moore,⁶ T. Moulik,⁵⁷ G. S. Muanza,¹⁵ M. Mulhearn,⁶⁹ O. Mundal,²² L. Mundim,³ E. Nagy,¹⁵ M. Naimuddin,⁴⁹ M. Narain,⁷⁶ H. A. Neal,⁶³ J. P. Negret,⁸ P. Neustroev,⁴⁰ H. Nilsen,²³ H. Nogima,³ S. F. Novaes,⁵ T. Nunnemann,²⁵ D. C. O'Neil,⁶ G. Obrant,⁴⁰ C. Ochando,¹⁶ D. Onoprienko,⁵⁸ J. Orduna,³³ N. Oshima,⁴⁹ N. Osman,⁴³ J. Osta,⁵⁴ R. Otec,¹⁰ G. J. Otero y Garzón,¹ M. Owen,⁴⁴ M. Padilla,⁴⁷ P. Padley,⁷⁹ M. Pangilinan,⁷⁶ N. Parashar,⁵⁵ S.-J. Park,^{22,||} S. K. Park,³¹ J. Parsons,⁶⁹ R. Partridge,⁷⁶ N. Parua,⁵³ A. Patwa,⁷² G. Pawloski,⁷⁹ B. Penning,²³ M. Perfilov,³⁸ K. Peters,⁴⁴ Y. Peters,⁴⁴ P. Pétroff,¹⁶ R. Piegaia,¹ J. Piper,⁶⁴ M.-A. Pleier,²² P. L. M. Podesta-Lerma,^{33,**} V. M. Podstavkov,⁴⁹ Y. Pogorelov,⁵⁴ M.-E. Pol,² P. Polozov,³⁷ A. V. Popov,³⁹ C. Potter,⁶ W. L. Prado da Silva,³ H. B. Prosper,⁴⁸ S. Protopopescu,⁷² J. Qian,⁶³ A. Quadt,^{22,||} B. Quinn,⁶⁵ A. Rakitine,⁴² M. S. Rangel,¹⁶ K. Ranjan,²⁸ P. N. Ratoff,⁴² P. Renkel,⁷⁸ P. Rich,⁴⁴ M. Rijssenbeek,⁷¹ I. Ripp-Baudot,¹⁹ F. Rizatdinova,⁷⁵ S. Robinson,⁴³ R. F. Rodrigues,³ M. Rominsky,⁷⁴ C. Royon,¹⁸ P. Rubinov,⁴⁹ R. Ruchti,⁵⁴ G. Safronov,³⁷ G. Sajot,¹⁴ A. Sánchez-Hernández,³³ M. P. Sanders,¹⁷ B. Sanghi,⁴⁹ G. Savage,⁴⁹ L. Sawyer,⁵⁹

T. Scanlon,⁴³ D. Schaile,²⁵ R. D. Schamberger,⁷¹ Y. Scheglov,⁴⁰ H. Schellman,⁵² T. Schliephake,²⁶ S. Schlobohm,⁸¹ C. Schwanenberger,⁴⁴ R. Schwienhorst,⁶⁴ J. Sekaric,⁴⁸ H. Severini,⁷⁴ E. Shabalina,⁵⁰ M. Shamim,⁵⁸ V. Shary,¹⁸ A. A. Shchukin,³⁹ R. K. Shivpuri,²⁸ V. Siccaldi,¹⁹ V. Simak,¹⁰ V. Sirotenko,⁴⁹ P. Skubic,⁷⁴ P. Slattery,⁷⁰ D. Smirnov,⁵⁴ G. R. Snow,⁶⁶ J. Snow,⁷³ S. Snyder,⁷² S. Söldner-Rembold,⁴⁴ L. Sonnenschein,²¹ A. Sopczak,⁴² M. Sosebee,⁷⁷ K. Soustruznik,⁹ B. Spurlock,⁷⁷ J. Stark,¹⁴ V. Stolin,³⁷ D. A. Stoyanova,³⁹ J. Strandberg,⁶³ S. Strandberg,⁴¹ M. A. Strang,⁶⁸ E. Strauss,⁷¹ M. Strauss,⁷⁴ R. Ströhmer,²⁵ D. Strom,⁵² L. Stutte,⁴⁹ S. Sumowidagdo,⁴⁸ P. Svoisky,³⁵ M. Takahashi,⁴⁴ A. Tanasijczuk,¹ W. Taylor,⁶ B. Tiller,²⁵ F. Tissandier,¹³ M. Titov,¹⁸ V. V. Tokmenin,³⁶ I. Torchiani,²³ D. Tsybychev,⁷¹ B. Tuchming,¹⁸ C. Tully,⁶⁷ P. M. Tuts,⁶⁹ R. Unalan,⁶⁴ L. Uvarov,⁴⁰ S. Uvarov,⁴⁰ S. Uzunyan,⁵¹ B. Vachon,⁶ P. J. van den Berg,³⁴ R. Van Kooten,⁵³ W. M. van Leeuwen,³⁴ N. Varelas,⁵⁰ E. W. Varnes,⁴⁵ I. A. Vasilyev,³⁹ P. Verdier,²⁰ L. S. Vertogradov,³⁶ M. Verzocchi,⁴⁹ D. Vilanova,¹⁸ P. Vint,⁴³ P. Vokac,¹⁰ M. Voutilainen,^{66,††} R. Wagner,⁶⁷ H. D. Wahl,⁴⁸ M. H. L. S. Wang,⁴⁹ J. Warchol,⁵⁴ G. Watts,⁸¹ M. Wayne,⁵⁴ G. Weber,²⁴ M. Weber,^{49,‡‡} L. Welty-Rieger,⁵³ A. Wenger,^{23,§§} M. Wetstein,⁶⁰ A. White,⁷⁷ D. Wicke,²⁶ M. R. J. Williams,⁴² G. W. Wilson,⁵⁷ S. J. Wimpenny,⁴⁷ M. Wobisch,⁵⁹ D. R. Wood,⁶² T. R. Wyatt,⁴⁴ Y. Xie,⁷⁶ C. Xu,⁶³ S. Yacoob,⁵² R. Yamada,⁴⁹ W.-C. Yang,⁴⁴ T. Yasuda,⁴⁹ Y. A. Yatsunenkov,³⁶ Z. Ye,⁴⁹ H. Yin,⁷ K. Yip,⁷² H. D. Yoo,⁷⁶ S. W. Youn,⁵² J. Yu,⁷⁷ C. Zeitnitz,²⁶ S. Zelitch,⁸⁰ T. Zhao,⁸¹ B. Zhou,⁶³ J. Zhu,⁷¹ M. Zielinski,⁷⁰ D. Zieminska,⁵³ L. Zivkovic,⁶⁹ V. Zutshi,⁵¹ and E. G. Zverev³⁸

(D0 Collaboration)

¹Universidad de Buenos Aires, Buenos Aires, Argentina²LAFEX, Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil³Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil⁴Universidade Federal do ABC, Santo André, Brazil⁵Instituto de Física Teórica, Universidade Estadual Paulista, São Paulo, Brazil⁶University of Alberta, Edmonton, Alberta, Canada;

Simon Fraser University, Burnaby, British Columbia, Canada;

York University, Toronto, Ontario, Canada

and McGill University, Montreal, Quebec, Canada

⁷University of Science and Technology of China, Hefei, People's Republic of China⁸Universidad de los Andes, Bogotá, Colombia⁹Center for Particle Physics, Charles University, Faculty of Mathematics and Physics, Prague, Czech Republic¹⁰Czech Technical University in Prague, Prague, Czech Republic¹¹Center for Particle Physics, Institute of Physics, Academy of Sciences of the Czech Republic, Prague, Czech Republic¹²Universidad San Francisco de Quito, Quito, Ecuador¹³LPC, Université Blaise Pascal, CNRS/IN2P3, Clermont, France¹⁴LPSC, Université Joseph Fourier Grenoble 1, CNRS/IN2P3, Institut National Polytechnique de Grenoble, Grenoble, France¹⁵CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille, France¹⁶LAL, Université Paris-Sud, IN2P3/CNRS, Orsay, France¹⁷LPNHE, IN2P3/CNRS, Universités Paris VI and VII, Paris, France¹⁸CEA, Ifju, SPP, Saclay, France¹⁹IPHC, Université de Strasbourg, CNRS/IN2P3, Strasbourg, France²⁰IPNL, Université Lyon 1, CNRS/IN2P3, Villeurbanne, France and Université de Lyon, Lyon, France²¹III. Physikalisches Institut A, RWTH Aachen University, Aachen, Germany²²Physikalisches Institut, Universität Bonn, Bonn, Germany²³Physikalisches Institut, Universität Freiburg, Freiburg, Germany²⁴Institut für Physik, Universität Mainz, Mainz, Germany²⁵Ludwig-Maximilians-Universität München, München, Germany²⁶Fachbereich Physik, University of Wuppertal, Wuppertal, Germany²⁷Panjab University, Chandigarh, India²⁸Delhi University, Delhi, India²⁹Tata Institute of Fundamental Research, Mumbai, India³⁰University College Dublin, Dublin, Ireland³¹Korea Detector Laboratory, Korea University, Seoul, Korea³²SungKyunKwan University, Suwon, Korea³³CINVESTAV, Mexico City, Mexico³⁴FOM-Institute NIKHEF and University of Amsterdam/NIKHEF, Amsterdam, The Netherlands³⁵Radboud University Nijmegen/NIKHEF, Nijmegen, The Netherlands³⁶Joint Institute for Nuclear Research, Dubna, Russia

³⁷*Institute for Theoretical and Experimental Physics, Moscow, Russia*³⁸*Moscow State University, Moscow, Russia*³⁹*Institute for High Energy Physics, Protvino, Russia*⁴⁰*Petersburg Nuclear Physics Institute, St. Petersburg, Russia*⁴¹*Stockholm University, Stockholm, Sweden, and Uppsala University, Uppsala, Sweden*⁴²*Lancaster University, Lancaster, United Kingdom*⁴³*Imperial College, London, United Kingdom*⁴⁴*University of Manchester, Manchester, United Kingdom*⁴⁵*University of Arizona, Tucson, Arizona 85721, USA*⁴⁶*California State University, Fresno, California 93740, USA*⁴⁷*University of California, Riverside, California 92521, USA*⁴⁸*Florida State University, Tallahassee, Florida 32306, USA*⁴⁹*Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA*⁵⁰*University of Illinois at Chicago, Chicago, Illinois 60607, USA*⁵¹*Northern Illinois University, DeKalb, Illinois 60115, USA*⁵²*Northwestern University, Evanston, Illinois 60208, USA*⁵³*Indiana University, Bloomington, Indiana 47405, USA*⁵⁴*University of Notre Dame, Notre Dame, Indiana 46556, USA*⁵⁵*Purdue University Calumet, Hammond, Indiana 46323, USA*⁵⁶*Iowa State University, Ames, Iowa 50011, USA*⁵⁷*University of Kansas, Lawrence, Kansas 66045, USA*⁵⁸*Kansas State University, Manhattan, Kansas 66506, USA*⁵⁹*Louisiana Tech University, Ruston, Louisiana 71272, USA*⁶⁰*University of Maryland, College Park, Maryland 20742, USA*⁶¹*Boston University, Boston, Massachusetts 02215, USA*⁶²*Northeastern University, Boston, Massachusetts 02115, USA*⁶³*University of Michigan, Ann Arbor, Michigan 48109, USA*⁶⁴*Michigan State University, East Lansing, Michigan 48824, USA*⁶⁵*University of Mississippi, University, Mississippi 38677, USA*⁶⁶*University of Nebraska, Lincoln, Nebraska 68588, USA*⁶⁷*Princeton University, Princeton, New Jersey 08544, USA*⁶⁸*State University of New York, Buffalo, New York 14260, USA*⁶⁹*Columbia University, New York, New York 10027, USA*⁷⁰*University of Rochester, Rochester, New York 14627, USA*⁷¹*State University of New York, Stony Brook, New York 11794, USA*⁷²*Brookhaven National Laboratory, Upton, New York 11973, USA*⁷³*Langston University, Langston, Oklahoma 73050, USA*⁷⁴*University of Oklahoma, Norman, Oklahoma 73019, USA*⁷⁵*Oklahoma State University, Stillwater, Oklahoma 74078, USA*⁷⁶*Brown University, Providence, Rhode Island 02912, USA*⁷⁷*University of Texas, Arlington, Texas 76019, USA*⁷⁸*Southern Methodist University, Dallas, Texas 75275, USA*⁷⁹*Rice University, Houston, Texas 77005, USA*⁸⁰*University of Virginia, Charlottesville, Virginia 22901, USA*⁸¹*University of Washington, Seattle, Washington 98195, USA*

(Received 4 March 2009; revised manuscript received 10 April 2009; published 24 August 2009)

We report observation of the electroweak production of single top quarks in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV based on 2.3 fb^{-1} of data collected by the D0 detector at the Fermilab Tevatron Collider. Using events containing an isolated electron or muon and missing transverse energy, together with jets originating from the fragmentation of b quarks, we measure a cross section of $\sigma(p\bar{p} \rightarrow tb + X, tqb + X) = 3.94 \pm 0.88 \text{ pb}$. The probability to measure a cross section at this value or higher in the absence of signal is 2.5×10^{-7} , corresponding to a 5.0 standard deviation significance for the observation.

DOI: 10.1103/PhysRevLett.103.092001

PACS numbers: 14.65.Ha, 12.15.Hh, 12.15.Ji, 13.85.Qk

At hadron colliders, top quarks can be produced in pairs via the strong interaction or singly via the electroweak interaction [1]. Top quarks were first observed via pair production at the Fermilab Tevatron Collider in 1995 [2].

Since then, pair production has been used to make precise measurements of several top-quark properties, including the top-quark mass [3]. Single top-quark production, on the other hand, serves as a probe of the Wtb interaction [4],

and its production cross section provides a direct measurement of the magnitude of the quark mixing matrix element V_{tb} without assuming three quark generations [5]. However, measuring the yield of single top quarks is difficult because of the small production rate and large backgrounds.

In 2007, we presented the first evidence for single top-quark production and the first direct measurement of $|V_{tb}|$ [6] using 0.9 fb^{-1} of Tevatron data at a center-of-mass energy of 1.96 TeV. This Letter describes the observation of a single top quark signal in 2.3 fb^{-1} of data. The CDF collaboration recently also reported such evidence in 2.2 fb^{-1} of data [7] and is concurrently reporting observation in 3.2 fb^{-1} of data [8].

When top quarks are produced singly, they are accompanied by a bottom quark in the s -channel production mode [9] or by both a bottom quark and a light quark in t -channel production [10,11]. We search for both of these processes at once. The sum of their predicted cross sections is $3.46 \pm 0.18 \text{ pb}$ [11] for a top-quark mass $m_t = 170 \text{ GeV}$, at which this analysis is performed. We refer to the s -channel process as “ tb ” production and the t -channel process as “ tqb .”

The analysis presented in this Letter is an improved version of the one from 2007 [6], with a larger data set. The data were collected with the D0 detector [12] using a logical OR of many trigger conditions in place of only the single-lepton plus jets triggers used previously. Several offline selection criteria, including b -jet identification requirements for double-tagged events, have been loosened. These improvements have increased the signal acceptance by 18%. The backgrounds are W bosons produced in association with jets, top-quark pair ($t\bar{t}$) production with decay into the lepton + jets and dilepton final states (when a jet or a lepton is not reconstructed), and multijet production, where a jet is misreconstructed as an electron or a heavy-flavor quark decays to a muon that passes isolation criteria. Z + jets and diboson processes form minor additional background components.

We consider events with two, three, or four jets (which allows for additional jets from initial-state and final-state radiation), reconstructed using a cone algorithm in (y, ϕ) space, where y is the rapidity and ϕ is the azimuthal angle, and the cone radius is 0.5 [6]. The highest- p_T (leading) jet must have $p_T > 25 \text{ GeV}$, and subsequent jets have $p_T > 15 \text{ GeV}$; all jets have pseudorapidity $|\eta| < 3.4$. We require $20 < \cancel{E}_T < 200 \text{ GeV}$ for events with two jets and $25 < \cancel{E}_T < 200 \text{ GeV}$ for events with three or four jets. Events must contain only one isolated electron with $p_T > 15 \text{ GeV}$ and $|\eta| < 1.1$ ($p_T > 20 \text{ GeV}$ for three- or four-jet events), or one isolated muon with $p_T > 15 \text{ GeV}$ and $|\eta| < 2.0$. The background from multijets events is kept to $\approx 5\%$ by requiring high total transverse energy and by demanding that the \cancel{E}_T is not along the direction of the lepton or the leading jet. To enhance the signal fraction, one or two of

the jets are required to originate from long-lived b hadrons using a neural network (NN) b -jet tagging algorithm [13]. After b -jet identification, we require the leading b -tagged jet to have $p_T > 20 \text{ GeV}$. To further improve the sensitivity, we split the data by lepton flavor, number of jets, and b -tagged jets, and data collection period.

We model the signal using the SINGLETOP event generator [14] interfaced to PYTHIA [15]. We assume the SM prediction for the ratio of the tb and tqb cross sections [11]. The $t\bar{t}$, W + jets, and Z + jets backgrounds are simulated using the ALPGEN leading-log MC event generator [16] and PYTHIA to model hadronization. The $t\bar{t}$ background is normalized to the predicted cross section [17]. The diboson backgrounds are modeled using PYTHIA. In the simulation of the W + jets backgrounds, we scale the ALPGEN cross sections for events with heavy-flavor jets by factors derived from calculations of next-to-leading-order effects [18]: $Wb\bar{b}$ and $Wc\bar{c}$ are scaled by 1.47, and Wcj by 1.38.

All MC events are passed through a GEANT-based simulation of the D0 detector and are reconstructed using the same software as for the data. Small differences between data and simulation in the lepton and jet reconstruction efficiencies and resolutions are corrected in the simulation as measured from separate data samples. We also correct the $\eta(\text{jets})$, $\Delta\phi(\text{jet1, jet2})$, and $\Delta\eta(\text{jet1, jet2})$ distributions in the W + jets samples to match data. The multijets background is modeled using independent data samples containing leptons that are not isolated. The multijets background, combined with the background from W + jets, is normalized to the lepton + jets data with other backgrounds subtracted, using the $p_T(\ell)$, \cancel{E}_T , and the W boson transverse mass distributions before b -jet identification is applied.

The b -tagging algorithm is modeled in simulated events by applying weights (“tag-rate functions”) measured from data that account for the probability for each jet to be tagged as a function of jet flavor, p_T , and η . After b tagging, an empirical correction of 0.95 ± 0.13 for the $Wb\bar{b}$ and $Wc\bar{c}$ fractions is derived from the b -tagged and non- b -tagged two-jet data and simulated samples.

The above selections give 4519 b -tagged lepton + jets events, which are expected to contain 223 ± 30 single top-quark events. Table I shows the event yields, separated by jet multiplicity. The percentage acceptances are $(3.7 \pm 0.5)\%$ for tb and $(2.5 \pm 0.3)\%$ for tqb .

Systematic uncertainties arise from each correction factor or function applied to the background and signal models. Most affect only the normalization, but three corrections modify in addition the shapes of the distributions; these are the jet energy scale corrections, the tag-rate functions, and the reweighting of the distributions in W + jets events. The largest uncertainties come from the jet energy scale [the normalization part is $(1.1\text{--}13.1)\%$ for signal and $(0.1\text{--}2.1)\%$ for background], the tag-rate func-

TABLE I. Number of expected and observed events in 2.3 fb^{-1} for e and μ , and 1 and 2 b -tagged analysis channels combined. The uncertainties include both statistical and systematic components.

Source	2 jets	3 jets	4 jets
$tb + tqb$ signal	139 ± 18	63 ± 10	21 ± 5
$W + \text{jets}$	1829 ± 161	637 ± 61	180 ± 18
$Z + \text{jets}$ and dibosons	229 ± 38	85 ± 17	26 ± 7
$t\bar{t}$	222 ± 35	436 ± 66	484 ± 71
Multijets	196 ± 50	73 ± 17	30 ± 6
Total prediction	2615 ± 192	1294 ± 107	742 ± 80
Data	2579	1216	724

tions [the normalization part is (2.1–7.0)% for single-tagged events and (9.0–11.4)% for double-tagged events], and the correction for jet-flavor composition in $W + \text{jets}$ events (13.7%), with smaller contributions from the integrated luminosity (6.1%), jet energy resolution (4.0%), initial-state and final-state radiation (0.6%–12.6%), b -jet fragmentation (2.0%), $t\bar{t}$ cross section (12.7%), and lepton efficiency corrections (2.5%). All other contributions have a smaller effect. The values given are the relative uncertainties on the individual sources. The total uncertainty on the background is (8–16)% depending on the analysis channel.

After event selection, we expect single top-quark events to constitute (3–9)% of the data sample. Since the uncertainty on the background is larger than the expected signal, we improve discrimination by using multivariate analysis techniques. We have developed three independent analyses based on boosted decision trees (BDT) [19], Bayesian neural networks (BNN) [20], and the matrix element (ME) method [21]. Our application of these techniques to D0's single top-quark searches is described in Ref. [6]. The analyses presented in this Letter differ from previous implementations in the choice of input variables and some detailed tuning of each technique.

The BDT analysis has reoptimized the input variables [22] into a common set of 64 variables for all analysis channels. The variables fall into five categories, single-object kinematics, global event kinematics, jet reconstruction, top-quark reconstruction, and angular correlations. Separate sets of trees are created with these variables for each channel. The BNN analysis uses the RuleFitJF algorithm [23] to select the most sensitive of these variables, then combines 18–28 of them into a single separate discriminant for each channel. The ME analysis uses only two-jet and three-jet events, divided into a $W + \text{jets}$ -dominated set and a $t\bar{t}$ -dominated set. It includes matrix elements for more background sources, adding $t\bar{t}$, WW , WZ , and ggg diagrams in the two-jet bin and $Wugg$ in the three-jet bin, to improve background rejection.

Each analysis uses the same data and background model and has the same sources of systematic uncertainty. We test

the analyses using ensembles of pseudo data sets created from background and signal at different cross sections to confirm linear behavior and thus an unbiased cross section measurement. The analyses are also checked extensively before b -tagging is applied, and using two control regions of the data, one dominated by $W + \text{jets}$ and the other by $t\bar{t}$ backgrounds, as shown in Fig. 1. These studies confirm that backgrounds are well modeled across the full range of the discriminant output.

The cross section is determined using the same Bayesian approach as in our previous studies [6]. This involves forming a binned likelihood as a product over all bins and channels, evaluated separately for each multivariate discriminant, with no cuts applied to the outputs. The central value of the cross section is defined by the position of the peak in the posterior density, and the 68% interval about the peak is taken as the uncertainty on the measurement. Systematic uncertainties, including all correlations, are reflected in this posterior interval.

We extract inclusive single top-quark cross sections $\sigma(p\bar{p} \rightarrow tb + X, tqb + X)$ of $\sigma_{\text{BDT}} = 3.74^{+0.95}_{-0.79}$ pb, $\sigma_{\text{BNN}} = 4.70^{+1.18}_{-0.93}$ pb, and $\sigma_{\text{ME}} = 4.30^{+0.99}_{-1.20}$ pb. The sensitivity of the analyses to a contribution from single top-quark production is estimated by generating an ensemble of pseudo data sets that sample the background model and its uncertainties, with no signal present. We measure a cross section from each pseudo data set, and hence obtain the probability that the SM cross section is reached. This provides expected sensitivities [stated in terms of Gaussian standard deviations, (SD)] of 4.3, 4.1, and 4.1 SD for the BDT, BNN, and ME analyses, respectively. The measured significances, obtained by counting the number of pseudo data sets above the measured cross section, are 4.6, 5.2, and 4.9 SD, respectively.

The three multivariate techniques use the same data sample but are not completely correlated: the correlation of the measured cross section using pseudo data sets with background and SM signal is BDT : BNN = 74%, BDT : ME = 60%, BNN : ME = 57%. Their combination therefore leads to increased sensitivity and a more precise measurement of the cross section. We use the three discriminant outputs as inputs to a second set of Bayesian




FIG. 1 (color online). The combination discriminant outputs for (a) $W + \text{jets}$ and (b) $t\bar{t}$ cross-check samples. H_T is the scalar sum of the p_T of the final-state objects (lepton, \cancel{E}_T , and jets).




FIG. 2 (color online). Distribution of the combination output for all 24 analysis channels combined, (a) full range, and (b) high signal region. The bins have been ordered by their expected signal to background ratio and the signal is normalized to the measured cross section. The hatched band indicates the total uncertainty on the background. For the ranked combination output >0.92 , (c) shows the distribution of lepton charge times pseudorapidity of the leading not- b -tagged jet, and (d) shows the reconstructed top-quark mass.

neural networks, and obtain the combined cross section and its signal significance from the new discriminant output. The resulting expected significance is 4.5 SD. Figure 2 illustrates the importance of the signal when comparing data to prediction. The measured cross section is $\sigma(p\bar{p} \rightarrow tb + X, tqb + X) = 3.94 \pm 0.88$ pb.

The measurement has a p value of 2.5×10^{-7} , corresponding to a significance of 5.0 SD. The expected and measured posterior densities and the background-only pseudodataset measurements are shown in Fig. 3.

We use the cross section measurement to determine the Bayesian posterior for $|V_{tb}|^2$ in the interval $[0,1]$ and ex-




FIG. 3 (color online). (a) Expected SM and measured Bayesian posterior probability densities for the $tb + tqb$ cross section. The shaded regions indicate 1 standard deviation above and below the peak positions. (b) Measured cross sections using the ensemble of background-only pseudo data sets (containing full systematics and no signal) used to measure the significance of the result.

tract a limit of $|V_{tb}| > 0.78$ at 95% C.L. within the SM [6]. When the upper constraint is removed, we measure $|V_{tb}f_1^L| = 1.07 \pm 0.12$, where f_1^L is the strength of the left-handed Wtb coupling.

In summary, we have measured the single top-quark production cross section using 2.3 fb^{-1} of data at the D0 experiment. We measure a cross section for the combined $tb + tqb$ channels of 3.94 ± 0.88 pb. Our result provides an improved direct measurement of V_{tb} . The measured single top-quark signal corresponds to an excess over the predicted background with a significance of 5.0 SD—observation of single top-quark production.

We thank the staffs at Fermilab and collaborating institutions, and acknowledge support from the DOE and NSF (USA); CEA and CNRS/IN2P3 (France); FASI, Rosatom and RFBR (Russia); CNPq, FAPERJ, FAPESP and FUNDUNESP (Brazil); DAE and DST (India); Colciencias (Colombia); CONACyT (Mexico); KRF and KOSEF (Korea); CONICET and UBACyT (Argentina); FOM (The Netherlands); STFC (United Kingdom); MSMT and GACR (Czech Republic); CRC Program, CFI, NSERC and WestGrid Project (Canada); BMBF and DFG (Germany); SFI (Ireland); The Swedish Research Council (Sweden); CAS and CNSF (China); and the Alexander von Humboldt Foundation (Germany).

- *Visitor from Augustana College, Sioux Falls, SD, USA.
- †Visitor from Rutgers University, Piscataway, NJ, USA.
- ‡Visitor from The University of Liverpool, Liverpool, United Kingdom.
- §Visitor from CERN, Geneva, Switzerland.
- ||Visitor from II. Physikalisches Institut, Georg-August-University, Göttingen, Germany.
- ¶Visitor from Centro de Investigacion en Computacion - IPN, Mexico City, Mexico.
- **Visitor from ECFM, Universidad Autonoma de Sinaloa, Culiacán, Mexico.
- ††Visitor from Helsinki Institute of Physics, Helsinki, Finland.
- ‡‡Visitor from Universität Bern, Bern, Switzerland.
- §§Visitor from Universität Zürich, Zürich, Switzerland.
- |||Deceased.

- [1] S. S. D. Willenbrock and D. A. Dicus, Phys. Rev. D **34**, 155 (1986).
- [2] F. Abe *et al.* (CDF Collaboration), Phys. Rev. Lett. **74**, 2626 (1995); S. Abachi *et al.* (D0 Collaboration), Phys. Rev. Lett. **74**, 2632 (1995).
- [3] C. Amsler *et al.*, (Particle Data Group), Phys. Lett. B **667**, 1 (2008).
- [4] A. P. Heinson *et al.* Phys. Rev. D **56**, 3114 (1997); V. M. Abazov *et al.* (D0 Collaboration), Phys. Rev. Lett. **101**, 221801 (2008).
- [5] G. V. Jikia and S. R. Slabospitsky, Phys. Lett. B **295**, 136 (1992).

- [6] V.M. Abazov *et al.* (D0 Collaboration), Phys. Rev. Lett. **98**, 181802 (2007); V.M. Abazov *et al.* (D0 Collaboration), Phys. Rev. D **78**, 012005 (2008).
- [7] T. Aaltonen *et al.* (CDF Collaboration), Phys. Rev. Lett. **101**, 252001 (2008).
- [8] T. Aaltonen *et al.* (CDF Collaboration), following Letter, Phys. Rev. Lett. **103**, 092002 (2009).
- [9] S. Cortese and R. Petronzio, Phys. Lett. B **253**, 494 (1991).
- [10] C.-P. Yuan, Phys. Rev. D **41**, 42 (1990).
- [11] N. Kidonakis, Phys. Rev. D **74**, 114012 (2006). Individual cross sections: $\sigma_{tb} = 1.12 \pm 0.05$ pb and $\sigma_{tqb} = 2.34 \pm 0.13$ pb.
- [12] V.M. Abazov *et al.* (D0 Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **565**, 463 (2006).
- [13] T. Scanlon, Ph.D. thesis, Imperial College, University of London, 2006.
- [14] E. E. Boos *et al.*, Phys. At. Nucl. **69**, 1317 (2006). We used SINGLETOP version 4.2p1.
- [15] T. Sjöstrand, S. Mrenna, and P. Skands, J. High Energy Phys. 05 (2006) 026. We used PYTHIA version 6.409.
- [16] M. L. Mangano *et al.*, J. High Energy Phys. 07 (2003) 001. We used ALPGEN version 2.11.
- [17] N. Kidonakis and R. Vogt, Phys. Rev. D **68**, 114014 (2003). We used $\sigma(p\bar{p} \rightarrow t\bar{t} + X) = 7.91^{+0.61}_{-1.01}$ pb, including an uncertainty component for the top-quark mass.
- [18] J. M. Campbell and R. K. Ellis, Phys. Rev. D **65**, 113007 (2002).
- [19] L. Breiman *et al.*, *Classification and Regression Trees* (Wadsworth, Stamford, 1984); J. A. Benitez, Ph.D. thesis, Michigan State University, 2009; D. Gillberg, Ph.D. thesis, Simon Fraser University, 2009.
- [20] R. M. Neal, *Bayesian Learning for Neural Networks* (Springer-Verlag, New York, 1996); A. Tanasijczuk, Ph.D. thesis, Universidad de Buenos Aires, 2009.
- [21] V.M. Abazov *et al.* (D0 Collaboration), Nature (London) **429**, 638 (2004); M. Pangilinan, Ph.D. thesis, Brown University, 2009.
- [22] Q.-H. Cao *et al.* Phys. Rev. D **71**, 054023 (2005); Q.-H. Cao *et al.*, Phys. Rev. D **72**, 094027 (2005).
- [23] J. H. Friedman and B. E. Popescu, Ann. Appl. Stat. **2**, 916 (2008).