

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Agronomy & Horticulture -- Faculty Publications

Agronomy and Horticulture Department

1988

Forage Quality and Performance of Yearlings Grazing Switchgrass Strains Selected for Differing Digestibility

Bruce Anderson

University of Nebraska-Lincoln, banderson1@unl.edu

J. K. Ward

University of Nebraska-Lincoln

K. P. Vogel

United States Department of Agriculture, kvogel1@unl.edu

M. G. Ward

United States Department of Agriculture

Herman J. Gorz

United States Department of Agriculture

See next page for additional authors

Follow this and additional works at: <https://digitalcommons.unl.edu/agronomyfacpub>



Part of the [Plant Sciences Commons](#)

Anderson, Bruce; Ward, J. K.; Vogel, K. P.; Ward, M. G.; Gorz, Herman J.; and Haskins, Francis A., "Forage Quality and Performance of Yearlings Grazing Switchgrass Strains Selected for Differing Digestibility" (1988). *Agronomy & Horticulture -- Faculty Publications*. 308.
<https://digitalcommons.unl.edu/agronomyfacpub/308>

This Article is brought to you for free and open access by the Agronomy and Horticulture Department at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Agronomy & Horticulture -- Faculty Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors

Bruce Anderson, J. K. Ward, K. P. Vogel, M. G. Ward, Herman J. Gorz, and Francis A. Haskins

FORAGE QUALITY AND PERFORMANCE OF YEARLINGS GRAZING SWITCHGRASS STRAINS SELECTED FOR DIFFERING DIGESTIBILITY¹

Bruce Anderson,² J. K. Ward,³ K. P. Vogel,⁴ M. G. Ward,⁵
H. J. Gorz⁴ and F. A. Haskins²

University of Nebraska, Lincoln 68583

ABSTRACT

Switchgrass (*Panicum virgatum*) is a native, perennial warm-season grass commonly used as summer pasture. Three strains of switchgrass with different in vitro DM disappearance (IVDMD) values were evaluated with yearling cattle (*Bos taurus*) in three replicated grazing trials. All strains were similar in genetic background, maturity and yield potential. 'Trailblazer' (high-IVDMD) resulted in higher ($P < .05$) ADG than 'Pathfinder' and 'low-IVDMD'. Available forage of Trailblazer had higher IVDMD than Pathfinder and low-IVDMD. The IVDMD differences among strains were similar throughout the season; IVDMD declined between .3 and .5 percentage units per day. Forage availability, CP, NDF and ADF, permanganate lignin and ash were similar among strains. This study documents that plant breeders can select switchgrass plants for higher IVDMD and that strains consisting of progeny of these plants can result in higher animal gains than strains consisting of progeny of unselected or low-IVDMD plants with similar genetic background, forage yield, maturity, morphology and chemical composition.

(Key Words: Beef Cattle, Pastures, Growth Rate, Plant Breeding, *Panicum virgatum*, In Vitro Digestibility.)

Introduction

Switchgrass (*Panicum virgatum*) is a perennial warm-season grass that complements cool-season grasses to improve season-long gains of grazing livestock (Krueger and Curtis, 1979; Matches et al., 1982; Burns et al., 1984). In vitro DM disappearance (IVDMD) can be improved through genetic selection of forage. Burton et al. (1967) increased IVDMD of bermudagrass (*Cynodon dactylon*) by over 12% by breeding, which increased ADG by grazing yearling steers by 30% (Chapman et al., 1972).

Vogel et al. (1981) developed strains of switchgrass that differed in IVDMD by 4.3 percentage units. There were no differences ($P > .05$) among the selected strains in ADF, permanganate lignin, CP or ash content (Vogel et al., 1984).

Our trials were conducted to determine if genetic improvements in IVDMD of switchgrass would increase gains of beef yearlings grazing switchgrass pastures.

Materials and Methods

Three grazing trials were conducted at the Agricultural Research and Development Center near Mead, NE on a Sharpsburg silty clay loam soil (fine, montmorillonitic, mesic, Typic Argiudoll). September through August precipitations during the trial were 960, 770 and 620 mm for 1982, 1983 and 1985, respectively, and high/low temperatures averaged within each year over June, July and August were 29/16, 32/17 and 29/15°C.

Two new switchgrass strains, 'Trailblazer' (formerly called high-IVDMD PC) and 'low-IVDMD PC', which had been developed from the same population by one cycle of divergent, restricted phenotypic selection for IVDMD, and a third cultivar, 'Pathfinder', which is similar in yield, maturity and origin (Vogel et al., 1981), were planted during May, 1981. Twelve .4-ha pastures were established in a randomized complete block design with four replicate

¹Journal paper no. 8443. Univ. of Nebraska Agric. Res. Div.

²Dept. of Agron.

³Dept. of Anim. Sci.

⁴USDA-ARS.

⁵Present address: Fort Keogh Livest. and Range Res. Sta., USDA, Miles City, MT 59301.

Received October 1, 1987.

Accepted March 24, 1988.

pastures per strain. Atrazine [6-chloro-*N*-ethyl-*N'*-(1-methylethyl)-1,3,5-triazine-2,4-diamine] at 2.2 kg/ha was applied pre-emerge to each pasture. By September, 1981, full stands had developed and many plants were forming seedstalks.

In April of each year, residue from the previous year was burned and 2.2 kg/ha of atrazine was applied. In early May, 112 kg/ha of N was applied to each pasture. Pastures were grazed continuously by three randomly allotted beef yearlings/pasture in 1982 and 1983 and by four yearlings in 1985. Angus × Hereford (AH) heifers averaging 303 kg, Simmental × Angus-Hereford (SAH) steers averaging 310 kg and SAH steers averaging 315 kg were used in 1982, 1983 and 1985, respectively.

Grazing began for the 1982 trial on June 24 when switchgrass was 80 cm tall and in the five- to six-leaf stage; grazing continued until September 1. During this period, many seedstalks developed; grass height eventually exceeded 160 cm. Yearlings were given ad libitum access to water, trace mineral-vitamin mix⁶ and salt. Yearlings were weighed following an overnight shrink without feed or water at the beginning and end of the trial. In 1983 and 1985, grazing began on June 8 and May 31, respectively, when switchgrass was 30 cm tall and in the four-leaf stage. Grazing was not conducted on one low-IVDMD pasture due to loss of stand caused by saturated soils. Grass height remained between 15 and 50 cm throughout these grazing seasons. Grazing ended August 9 and July 17, respectively, when grass height was between 15 and 20 cm and the herbage remaining was nearly all stem and leaf sheath tissue.

Available forage was estimated weekly by random placement of three 30-cm × 182-cm frames within each pasture and clipping the enclosed forage to a 2.5-cm stubble. In addition, forage samples of the top one-third of plants (tops) were collected randomly throughout each pasture concurrently with available forage samples. All forage samples were dried at 50°C in a forced-air oven, ground through a mill with a 1-mm screen and subsampled for laboratory analyses.

⁶ Each kilogram of trace mineral-vitamin mix contained the following: NaCl, 110 g (minimum), 130 g (maximum); Ca, 110 g (minimum), 130 g (maximum); P, 120 g (minimum); I, 10 mg (minimum); Vitamin A, 45 USP.

The IVDMD was measured by the Tilley and Terry (1963) two-stage (48 h each) technique modified to include 1 g of urea per liter of buffer. The ruminal fluid was a mixture of equal parts collected from two fistulated crossbred steers, one fed a diet of alfalfa (*Medicago sativa*)-smooth brome (*Bromus inermis*) hay and the other fed ground corn cobs (*Zea mays*). Ruminal fluid was mixed with McDougall's buffer (1948) in a 30:70 (v:v) ratio. Crude protein of the switchgrass was determined as 6.25 × Kjeldahl-N (AOAC, 1975). The NDF, ADF, permanganate lignin and ash were determined sequentially (Van Soest and Robertson, 1980).

Yearling gains were analyzed using analysis of variance with individual pastures as the experimental unit, and plant analyses were evaluated as a split-plot in time with pastures as whole plots and sampling period as sub-plots (Steel and Torrie, 1980).

Results and Discussion

Daily gains of yearlings grazing Trailblazer switchgrass averaged 24 and 14% higher ($P < .05$) than of yearlings grazing Pathfinder and low-IVDMD, respectively. Average seasonal gains from the three strains were 351, 284 and 299 kg/ha, respectively, over the three grazing trials (Table 1).

Gains during 1982, when grazing began as switchgrass was jointing, were substantially less than during 1983 and 1985. Yearling cattle demonstrated a distinct preference for top grazing of switchgrass plants prior to the boot stage. They were able to consume forage at nearly the same rate as it was produced, preventing seedstalk development until after grazing was terminated. Switchgrass was in the early boot stage of development when grazing began in 1982, and yearlings were unable to consume the new growth as rapidly as it developed, allowing many seedheads to develop. After panicle emergence, visual observations showed that yearlings almost completely avoided grazing reproductive culms, preferring vegetative culms, new tillers and leaf blades. As a result, forage IVDMD and CP were lower in 1982 than in 1983 and 1985 and NDF, ADF and permanganate lignin content were higher (Table 2). Anderson and Matches (1983) reported that IVDMD and CP concentration of switchgrass declined rapidly as plants matured to the heading stage and suggested that grazing should begin prior to the boot stage. Thus, low

TABLE 1. PERFORMANCE OF BEEF YEARLINGS GRAZING SWITCHGRASS STRAINS IN 1982, 1983 AND 1985

Strain	1982 ^a		1983 ^b		1985 ^c		3-yr mean	
	Gain/ha	ADG	Gain/ha	ADG	Gain/ha	ADG	Gain/ha	ADG
Trailblazer	235	.45	456	.98	366	.77	351	.73
Pathfinder	143	.27	371	.80	345	.73	284	.59
Low-IVDMD	208	.41	354	.76	326	.68	299	.64
P <	.05	.05	.12	.12	.01	.01	.05	.05
SEM	19	.09	29	.16	6	.03	13	.07

kg

^a Three yearling heifers per .4 ha pasture, duration June 24 to Sept. 1 (69 d).

^b Three yearling steers per .4 ha pasture, duration June 8 to Aug. 9 (62 d).

^c Four yearling steers per .4 ha pasture, duration May 31 to July 17 (47 d).

gains in 1982 probably were due to offering yearlings a lower-quality diet, although gain potential probably was lower for AH heifers than for SAH steers.

Grazing earlier in 1983 and 1985 than in 1982 also reduced forage availability (Table 3). Yearlings had twice as much forage available for grazing in 1982 as in 1983 or 1985. Although increased forage availability often enhances selective grazing, the lower gains in 1982 vs other years suggest that quality of forage was not sufficient to produce as rapid gain in 1982.

Mean forage availability over the grazing season did not differ among switchgrass strains except in 1985, when Pathfinder averaged 23% more forage available than Trailblazer and low-IVDMD. In addition, leaf:stem ratios of all three strains were similar in small plots during the vegetative growth stage in early June through anthesis in August (data not shown). Thus, performance differences among strains probably was not due to differences in forage availability.

The IVDMD of all strains declined between .3 and .5 percentage units per day during all three trials (Figure 1); these were similar to declines reported for switchgrass from clipping studies (Anderson and Matches, 1983). Available forage of Trailblazer averaged 1.8 and 2.5 percentage units higher in IVDMD than Pathfinder and low-IVDMD, respectively, and IVDMD averaged 2.5 and 3.7 percentage units higher for tops of Trailblazer than of the other strains (Table 2). When available forage in small plots was harvested at early heading, Trailblazer averaged 4.0 and 4.3 percentage units higher in IVDMD than Pathfinder and low-IVDMD, respectively (Vogel et al., 1984).

Although IVDMD differences were smaller in these grazing trials than in previous clipping studies, Trailblazer was superior consistently to the other strains. Only in 1983 did IVDMD of available forage of Trailblazer fail to be superior ($P < .10$) to forage of other strains. The superiority of Trailblazer for IVDMD was consistent throughout the grazing season of all three trials (Figure 1), but all other measured forage components remained similar among strains within each grazing trial (Table 2).

Top grazing was considerable during 1983 and 1985, so forage samples of the top one-third of plants should have more closely resembled the diet than available forage. Tops generally had higher IVDMD and CP and lower NDF, ADF and ash than did available forage

TABLE 2. FORAGE QUALITY OF THREE SWITCHGRASS STRAINS DURING THREE GRAZING SEASONS

Strain	Available forage, %						Tops, %					
	IVDMD ^a	CP	NDF	ADF	Permanganate lignin	Ash	IVDMD ^a	CP	NDF	ADF	Permanganate lignin	Ash
	1982											
Trailblazer	49.4	9.1	72.1	41.7	7.4	1.3	55.0	11.0	70.1	38.0	5.9	.9
Pathfinder	47.2	8.4	73.3	43.4	8.0	1.3	52.9	10.9	71.0	39.0	6.2	1.0
Low-IVDMD	47.9	8.6	72.7	42.3	7.3	1.4	52.3	10.8	71.0	38.6	5.9	1.1
P <	.02	.09	.19	.19	.32	.18	.01	.81	.32	.30	.73	.12
SEM	1.07	.50	1.09	1.61	.93	.14	.61	.64	1.19	1.26	.98	.22
	1983											
Trailblazer	63.6	11.3	66.3	34.5	4.2	1.9	66.6	12.3	66.8	34.5	4.3	1.0
Pathfinder	62.1	11.2	66.0	34.8	4.2	1.4	62.1	11.4	67.3	35.1	4.6	1.1
Low-IVDMD	61.3	11.4	66.0	35.3	5.0	1.4	62.8	12.3	66.3	33.6	4.7	1.1
P <	.11	.94	.81	.80	.22	.66	.01	.09	.78	.37	.68	.54
SEM	1.83	.51	.92	1.16	.44	1.35	1.16	.67	1.64	1.26	.89	.75
	1985											
Trailblazer	61.1	9.6	69.6	37.6	4.6	.7	59.1	10.5	68.1	36.2	3.9	1.1
Pathfinder	59.3	8.9	70.8	39.0	5.2	.9	58.1	10.6	67.8	36.5	4.0	.9
Low-IVDMD	57.4	9.4	70.7	38.7	5.1	1.0	54.6	9.9	68.9	37.5	4.0	1.3
P <	.01	.07	.28	.37	.28	.51	.07	.15	.09	.11	.88	.73
SEM	.91	.44	1.17	1.52	.61	.32	1.75	.44	.46	.68	.61	.75

^aIVDMD = in vitro dry matter disappearance.

TOPS

AVAILABLE FORAGE




Figure 1. In vitro dry matter disappearance (IVDMD) of available herbage and the top one-third of switchgrass plants in Trailblazer (●), Pathfinder (■) and low-IVDMD (▲) pastures during 1982, 1983 and 1985. SE = standard error of the seasonal mean.

TABLE 3. MEAN FORAGE AVAILABILITY OVER THE GRAZING SEASON IN SWITCHGRASS PASTURES DURING 1982, 1983 AND 1985

Strains	Mean forage availability			
	1982	1983	1985	3-yr mean
	kg/ha			
Trailblazer	5,150	2,590	2,520	3,420
Pathfinder	4,970	2,130	3,040	3,380
Low-IVDMD	4,800	2,280	2,410	3,160
P <	.78	.42	.04	.61
SEM	143	103	71	38

due to higher proportions of stem material in available forage samples. The relationships among strains for forage quality, though, were nearly identical for available forage and tops.

Gain responses by yearlings grazing the three switchgrass strains were consistent even though the three trials were conducted under diverse conditions. Neither sex (heifers in 1982, steers in 1983 and 1985), breed cross (AH in 1982 and SAH in 1983 and 1985), stocking density (three yearlings/4 ha in 1982 and four yearlings in 1985) nor stage of grass maturity at initial grazing (boot in 1982, jointing in 1983 and 1985) altered the relationship among the three strains to cattle gains. Only during 1982, when low-IVDMD resulted in higher gains than Pathfinder, were gain relationships inconsistent.

Forage IVDMD was the only measured forage component to differ consistently among the three switchgrass strains. Because available forage and yield potential did not differ among the strains, intake presumably was similar. Previous research with bermudagrass showed that digestibility (Burton et al., 1967) and animal gains (Chapman et al., 1972) can be improved by plant breeding. Although in vivo and in vitro measurements of switchgrass digestibility may differ substantially with cattle, the present study documents that switchgrass selected for higher IVDMD can produce plant progeny that result in higher cattle gains than progeny of unselected or low-IVDMD strains with similar genetic background, yield potential, morphology, maturity and forage quality traits such as CP, NDF, ADF and lignin.

Literature Cited

Anderson, Bruce and A. G. Matches. 1983. Forage

yield, quality and persistence of switchgrass and caucasian bluestem. *Agron. J.* 75:119.

AOAC. 1975. *Official Methods of Analysis*. Association of Official Analytical Chemists, Washington, DC.

Burns, J. C., R. D. Mochrie and D. H. Timothy. 1984. Steer performance from two perennial Pennisetum species, switchgrass and a fescue - 'Coastal' bermudagrass system. *Agron. J.* 76: 795.

Burton, G. W., R. H. Hart and R. L. Lowrey. 1967. Improving forage quality in bermudagrass by breeding. *Crop Sci.* 7:329.

Chapman, H. D., W. H. Marchant, P. R. Utley, R. E. Hellwig and W. G. Monson. 1972. Performance of steers on Pensacola bahiagrass, coastal bermudagrass and coastcross-1 bermudagrass pastures and pellets. *J. Anim. Sci.* 34:373.

Krueger, C. R. and D. C. Curtis. 1979. Evaluation of big bluestem, indiagrass, sideoats grama and switchgrass pastures with yearling steers. *Agron. J.* 71:480.

Matches, A. G., F. A. Martz, D. P. Mowrey and S. Bell. 1982. Integration of subtropical perennial grasses into forage-animal systems. *Agron. Abstr.*, Am. Soc. of Agron., Madison, WI. p 150.

McDougall, E. I. 1948. Studies on ruminant saliva. I. The composition and output of sheep's saliva. *Biochem. J.* 43:99.

Steel, R.G.D. and J. H. Torrie. 1980. *Principles and Procedures of Statistics*. McGraw-Hill Book Co., New York.

Tilley, J.M.S. and R. A. Terry. 1963. A two-stage technique for the in vitro digestion of forage crops. *J. Br. Grassl. Soc.* 18:108.

Van Soest, P. J. and J. B. Robertson. 1980. Systems of analysis for evaluating fibrous feeds. In: W. J. Pigden, C. C. Balch and M. Graham (Ed.) *Standardization of Analytical Methodology for Feeds*. pp 49-60. Int. Div. Res. Ctr., Ottawa, Can.

Vogel, K. P., R. Britton, H. J. Gorz and F. A. Haskins. 1984. In vitro and in vivo analyses of hays of switchgrass strains selected for high and low in vitro dry matter digestibility. *Crop Sci.* 24:977.

Vogel, K. P., F. A. Haskins and H. J. Gorz. 1981. Divergent selection for in vitro dry matter digestibility in switchgrass. *Crop Sci.* 21:39.