

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

USGS Northern Prairie Wildlife Research Center Wildlife Damage Management, Internet Center for

2013

Age Structure of Moose (*Alces alces*) Killed by gray Wolves (*Canis lupus*) in Northeastern Minnesota, 1967–2011


L. David Mech

USGS Northern Prairie Wildlife Research Center, david_mech@usgs.gov

Michael E. Nelson

U.S. Geological Survey

Follow this and additional works at: <https://digitalcommons.unl.edu/usgsnpwrc>

 Part of the [Animal Sciences Commons](#), [Behavior and Ethology Commons](#), [Biodiversity Commons](#), [Environmental Policy Commons](#), [Recreation, Parks and Tourism Administration Commons](#), and the [Terrestrial and Aquatic Ecology Commons](#)

Mech, L. David and Nelson, Michael E., "Age Structure of Moose (*Alces alces*) Killed by gray Wolves (*Canis lupus*) in Northeastern Minnesota, 1967–2011" (2013). *USGS Northern Prairie Wildlife Research Center*. 356.

<https://digitalcommons.unl.edu/usgsnpwrc/356>

This Article is brought to you for free and open access by the Wildlife Damage Management, Internet Center for at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in USGS Northern Prairie Wildlife Research Center by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Age Structure of Moose (*Alces alces*) Killed by Gray Wolves (*Canis lupus*) in Northeastern Minnesota, 1967–2011

L. DAVID MECH¹ and MICHAEL E. NELSON²

U.S. Geological Survey, Northern Prairie Wildlife Research Center, 8711–37th Street SE, Jamestown, North Dakota 58401-7317 USA

¹Mailing address: 1920 Fitch Avenue, University of Minnesota, St. Paul, Minnesota 55108 USA

²Mailing address: 7332 Tracy Road, Duluth, Minnesota 55803 USA

Mech, L. David, and Michael E. Nelson. 2013. Age structure of Moose (*Alces alces*) killed by Gray Wolves (*Canis lupus*) in northeastern Minnesota, 1967–2011. *Canadian Field-Naturalist* 127(1): 70–71.

The ages of 77 adult Moose (*Alces alces*) killed by Gray Wolves (*Canis lupus*) during the period 1967–2011 in northeastern Minnesota were significantly older than those of a sample of 17 585 Moose killed by hunters in nearby Ontario. Our findings support those of earlier studies of protected Moose populations in national parks that found that Gray Wolves tend to kill disproportionately more older Moose.

Key Words: *Canis lupus*, Gray Wolf, *Alces alces*, Moose, predation, hunting, Minnesota, Ontario.

The age structure of Moose (*Alces alces*) killed by Gray Wolves (*Canis lupus*) is available from only two national parks in the United States where hunting by people is not allowed and from three areas in Alaska where Moose are hunted (Mech 1966; Peterson et al. 1984; Ballard et al. 1987; Mech et al. 1998). The samples of Moose killed by Gray Wolves from each hunted area are relatively small (47–117), given that Moose live to 20 or more years (Passmore et al. 1955). This article adds age data from another 77 Moose killed by Gray Wolves from a fourth (lightly) human-hunted area and assesses the age structure of all the samples.

This study was conducted as part of a radio-tracking study of Gray Wolves in the Superior National Forest in northeastern Minnesota (48°N, 92°W) (Mech 2009). There, Gray Wolves prey on both White-tailed Deer (*Odocoileus virginianus*) and Moose. The point estimates of the Moose population in the region varied from 2140 to 8840 between 1971 and 2011 (J. Giudice, Minnesota Department of Natural Resources, personal communication and M. Lenarz, Minnesota Department of Natural Resources, personal communication). Since 1971, Moose hunting on a lottery basis has been allowed, and about 200 Moose per year have been harvested (Edwards et al. 2004).

Of 107 Moose killed by Gray Wolves that we located by aerially radio-tracking Gray Wolves from 1967 through 2011, we examined 85 that we could sex; 45 were females and 40 were males. We collected teeth from 37 cows and 26 bulls from among the 85 and from 14 Moose from among the 22 of unknown sex (total 77 Moose). Mattson's Laboratory (Milltown, Montana) aged the teeth by cementum analysis (calves were probably underrepresented because there are fewer of their remains to find). The age structures of the two sexes were not significantly different (Kolmogorov-Smirnov test; $P > 0.40$). We therefore pooled them and added the unknowns that we could age. The ages of our sam-

ple varied from calves >6 months old to bulls 13 years old and a cow 19 years old (Figure 1).

The best, and about the only, age-structure data available for a hunted Moose population are those for 17 585 Moose killed by hunters in north-central Ontario, presumably a representative estimate of the Moose population at large (Timmermann and Rempel 1998). The age structure of our sample killed by Gray Wolves was significantly different from that of the Ontario population (Kolmogorov-Smirnov; $P < 0.0001$).

Peterson et al. (1984) plotted ages of Moose killed by Gray Wolves from various areas in Alaska in 6-year age classes (1–6, 7–12, and 13+). Of our sample of adult Moose killed by Gray Wolves, only 34% were 1–6 years old, similar to the 27% in the sample in Alaska in Peterson et al. (1984) (no significant difference), whereas 76% of the Ontario population was of this age class ($\chi^2 = 68.55$; $P < 0.0001$). Assuming that the




FIGURE 1. Age distribution of 77 Moose (*Alces alces*) killed by Gray Wolves (*Canis lupus*) in northeastern Minnesota between 1967–2011. (Calves are probably underrepresented.)

age structure of the Ontario Moose population at large reasonably approximates that of the study population in Superior National Forest, this is strong evidence that Moose 1–6 years of age are the least vulnerable to Gray Wolves in this area.

We have long known that Gray Wolves tend to kill a disproportionate number of older Moose and calves and that a disproportionate number of younger Moose > 1-year of age survive Gray Wolf predation (Mech 1966; Peterson 1977; Haber 1977; Peterson et al. 1984; Ballard et al. 1987; Mech et al. 1998). However, the actual age when disproportionate vulnerability begins may vary. Peterson et al. (1998) showed that Moose ≥ 9 years of age were more vulnerable on Isle Royale and those >12 years of age on the Kenai Peninsula were more vulnerable (Peterson et al. 1984).

In our sample, Moose ≥ 9 years old comprised 39% of the Moose >1 year of age killed by Gray Wolves, whereas Moose ≥ 9 years old comprised only 10% of the Ontario sample of Moose >1 year of age killed by hunters ($P < 0.0001$; $\chi^2 = 68.29$; 1 d.f.). Thus our study tends to confirm the findings on Isle Royale (Peterson et al. 1998) and adds to the general conclusion that Gray Wolves tend to kill older Moose, whether in populations hunted by people or not.

Acknowledgements

This study was funded by the U.S. Department of the Interior and the U.S. Forest Service. Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government. We thank numerous pilots for safe flying and R. Moen, P. Cross, and an anonymous reviewer for critiquing this manuscript and offering helpful suggestions for improvement.

Literature Cited

Ballard, W. B., J. S. Whitman, and C. L. Gardner. 1987. Ecology of an exploited wolf population in south-central Alaska. *Wildlife Monographs* 98.

Edwards, A. J., M. Schrage, and M. Lenarz. 2004. North-eastern Minnesota moose management – a case study in cooperation. *Alces* 40: 23–31.

Haber, G. C. 1968. The social structure and behavior of an Alaskan wolf population. Master's thesis, Northern Michigan University, Marquette, Michigan.

Haber, G. C. 1977. Socio-ecological dynamics of wolves and prey in a subarctic ecosystem. Ph.D. dissertation, University of British Columbia, Vancouver, British Columbia. 824 pages.

Mech, L. D. 1966. The Wolves of Isle Royale. National Parks Fauna Series No. 7. U.S. Government Printing Office. 210 pages.

Mech, L. D. 2009. Long-term research on wolves in the Superior National Forest. Pages 15–34 in *Recovery of Gray Wolves in the Great Lakes Region of the United States: an Endangered Species Success Story*. Edited by A. P. Wydeven, E. J. Heske, and T. R. Van Deelen. Springer, New York, New York. 350 pages.

Mech, L. D., L. G. Adams, T. J. Meier, J. W. Burch, and B. W. Dale. 1998. The Wolves of Denali. University of Minnesota Press, Minneapolis, Minnesota. 227 pages.

Passmore, R. C., R. L. Peterson, and A. T. Cringan. 1955. Appendix A: A study of mandibular tooth wear as an index to age of moose. Pages 223–246 in *North American Moose*. Edited by R. L. Peterson. University of Toronto Press, Toronto, Ontario.

Peterson, R. O. 1977. Wolf ecology and prey relationships on Isle Royale. National Park Service Science Monograph Series No. 11. Government Printing Office, Washington, D.C.

Peterson, R. O., J. D. Woolington, and T. N. Bailey. 1984. Wolves of the Kenai Peninsula, Alaska. *Wildlife Monographs* 88.

Peterson, R. O., N. J. Thomas, J. M. Thurber, J. A. Vucetich, and T. A. Waite. 1998. Population limitation and the wolves of Isle Royale. *Journal of Mammalogy* 79: 828–841.

Timmermann, H. R., and R. S. Rempel. 1998. Age and sex structure of hunter harvested moose under two harvest strategies in northcentral Ontario. *Alces* 34: 21–30.

Accepted 13 February 2013

Received 15 April 2013