

2005

# Cryptic plasmid pSKU146 from the wall-less plant pathogen *Spiroplasma kunkelii* encodes an adhesin and components of a type IV translocation-related conjugation system

Robert E. Davis

Ellen L. Dally

Rasa Jomantiene

Yan Zhao

Bruce Roe

*See next page for additional authors*

Follow this and additional works at: <http://digitalcommons.unl.edu/usdaarsfacpub>

 Part of the [Agricultural Science Commons](#)

Davis, Robert E.; Dally, Ellen L.; Jomantiene, Rasa; Zhao, Yan; Roe, Bruce; Lin, ShaoPing; and Shao, Jonathan, "Cryptic plasmid pSKU146 from the wall-less plant pathogen *Spiroplasma kunkelii* encodes an adhesin and components of a type IV translocation-related conjugation system" (2005). *Publications from USDA-ARS / UNL Faculty*. 345.  
<http://digitalcommons.unl.edu/usdaarsfacpub/345>

This Article is brought to you for free and open access by the U.S. Department of Agriculture: Agricultural Research Service, Lincoln, Nebraska at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Publications from USDA-ARS / UNL Faculty by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

---

**Authors**

Robert E. Davis, Ellen L. Dally, Rasa Jomantiene, Yan Zhao, Bruce Roe, ShaoPing Lin, and Jonathan Shao



ELSEVIER

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

SCIENCE @ DIRECT®

Plasmid 53 (2005) 179–190

PLASMID

[www.elsevier.com/locate/yplas](http://www.elsevier.com/locate/yplas)

## Cryptic plasmid pSKU146 from the wall-less plant pathogen *Spiroplasma kunkelii* encodes an adhesin and components of a type IV translocation-related conjugation system

Robert E. Davis<sup>a,\*</sup>, Ellen L. Dally<sup>a</sup>, Rasa Jomantiene<sup>a,b</sup>, Yan Zhao<sup>a</sup>,  
Bruce Roe<sup>c</sup>, ShaoPing Lin<sup>c</sup>, Jonathan Shao<sup>a</sup>

<sup>a</sup> Molecular Plant Pathology Laboratory, USDA-Agricultural Research Service, Beltsville, MD 20705, USA

<sup>b</sup> Microbial Molecular Biology Group, Fitovirus Laboratory, Institute of Botany, Vilnius 2021, Lithuania

<sup>c</sup> Advanced Center for Genome Technology, Department of Chemistry and Biochemistry, University of Oklahoma, Norman, OK 73019, USA

Received 17 March 2004, revised 21 June 2004

Communicated by K. Smalla

### Abstract

A cryptic plasmid of the wall-less plant pathogenic mollicute, *Spiroplasma kunkelii* CR2-3X, was cloned and its sequence analyzed. The 14,615 bp plasmid, designated pSKU146, has a nucleotide content of 28 mol% G + C, and contains 18 potential protein-coding regions (open reading frames, ORFs), of which six encode proteins that exhibit similarity to virulence-associated proteins involved in cell-to-cell adhesion or conjugal DNA transfer. One ORF encodes a 96 kDa protein, SkARP1, that is highly similar to SARP1 adhesin involved in attachment of *Spiroplasma citri* to insect vector gut membrane. Five ORFs encode proteins similar to TraE and Mob in walled bacteria, and to ORFs found in the integrative, conjugative element (ICEF) of *Mycoplasma fermentans*, respectively. Presence of domains similar to proteins of the Type IV secretion system in pathogenic bacteria suggests that spiroplasma possesses a related translocation system. Plasmid pSKU146 also contains two identical *oriT* regions each containing a nick sequence characteristic of the IncP conjugative plasmid family, as well as a 58 bp palindromic sequence, palSK1. Features in pSKU146 suggest that the plasmid functions as a mobile genetic element in conjugative transmission of spiroplasma pathogenicity-related genes.

Published by Elsevier Inc.

**Keywords:** Virulence; Mobilization; Conjugation; Pathogenicity; Vir domain

\* Corresponding author.

E-mail address: [davisr@ba.ars.usda.gov](mailto:davisr@ba.ars.usda.gov) (R.E. Davis).

## 1. Introduction

Spiroplasmas are helical, motile, cell wall-less prokaryotes that are classified, along with members of genus *Mycoplasma* and other wall-less bacteria, in class Mollicutes. In their descent from the Gram-positive, low G+C *Bacillus*–*Clostridium* group of walled bacteria, spiroplasmas underwent massive evolutionary genome reduction, while acquiring capabilities for arthropod parasitism, and in some cases plant pathogenicity. First discovered in association with plant disease (Davis and Worley, 1973; Davis et al., 1972), spiroplasmas comprise broadly diverse species, most of which parasitize a variety of insects or ticks (Gasparich, 2002). As in other bacteria, plasmids presumably have played key roles in spiroplasma evolution, and contributed to the diversity of *Spiroplasma* species, through horizontal exchange of genes encoding virulence factors, niche adaptation factors, and factors conferring competitive advantages on the spiroplasma cell. In the present work, we characterized a plasmid from a plant pathogen, *Spiroplasma kunkelii*.

*Spiroplasma kunkelii* is the causative agent of corn stunt disease, a major factor limiting maize production in the Americas (Davis and Worley, 1973; Davis et al., 1972; Whitcomb and Williamson, 1975). This spiroplasma is transmitted between maize plants exclusively by phloem-feeding leafhopper insects, in which it multiplies. In maize plants, the spiroplasma resides specifically in sieve cells of the plant's phloem tissue, and induces symptoms including general stunting, broad chlorotic stripes, poor filling of ears, sterility of tassels, and plant death. Although plasmids have been reported in *S. kunkelii* (Gasparich et al., 1993), none has previously been sequenced. Here we present the structural analysis of a previously unreported *S. kunkelii* plasmid, designated pSKU146, harboring regions potentially encoding proteins exhibiting significant homologies, respectively, with a spiroplasma adhesion-related protein and proteins involved in conjugal mating pair formation and DNA transfer.

## 2. Materials and methods

### 2.1. Cloning and sequencing of DNA

*Spiroplasma kunkelii* strain CR2-3X was grown in broth medium LD8A3 as described (Lee and Davis, 1989). Spiroplasma cells were embedded in agarose and high molecular weight DNA was prepared according to published procedures (Birren et al., 1999; Peterson et al., 2000). The high molecular weight DNA was partially digested with *Hind*III (New England BioLabs, Beverly, MA) and cloned in vector pBeloBac11 (Kim et al., 1996) according to Peterson et al. (2000). The cloned plasmid DNA was partially sequenced using automated sequencing and the data were combined with whole genome shotgun data from the Spiroplasma Genome Sequencing Project web site at [www.genome.ou.edu/spiro.html](http://www.genome.ou.edu/spiro.html). The sequence data from these two sources were assembled in silico using the SeqMan program of the sequence analysis software suite Lasergene (DNA-STAR, Madison, WI).

### 2.2. Nucleotide and amino acid sequence analyses

BLAST searches (Altschul et al., 1990), open reading frame (ORF) analysis, and COGNITOR searches (Tatusov et al., 2001) were carried out at the National Center for Biotechnology Information (NCBI) web site (<http://www.ncbi.nlm.nih.gov>). In silico restriction mapping and multiple sequence alignments were performed using the MapDraw and Megalign programs, respectively, of the Lasergene sequence analysis software suite, and conserved regions were defined as common segments in the alignment. A signal sequence search was done using SignalP (Nielsen et al., 1997) (<http://www.cbs.dtu.dk/services/SignalP>), and the Simple Modular Architecture Research Tool SMART (Letunic et al., 2002; Schultz et al., 1998) (<http://smart.embl-heidelberg.de>) was used to search for signal sequences, transmembrane regions, and regions of low complexity. Protein mass and isoelectric points were estimated using the GeneQuest option of the Lasergene suite. A search for gene orthologs in completely sequenced gen-

omes of *Mycoplasma* species was carried out by using the ortholog table construction option at the Whitehead Institute Center for Genome Research (<http://www-genome.wi.mit.edu/annotation/microbes/methanosarcina/keggmap.html>).

### 3. Results and discussion

The complete nucleotide sequence of plasmid pSKU146 has been deposited in the GenBank database under GenBank Accession No. AY528560. Plasmid pSKU146 is a circular molecule of 14,615 bp, within the size range of naturally occurring extrachromosomal DNAs thus far reported in *Spiroplasma* spp. and similar to the size of extrachromosomal DNAs reported in *S. kunkelii* (Gasparich and Hackett, 1994; Ranhand et al., 1980; Razin et al., 1987; Salvado et al., 1989). Plasmid pSKU146 has a G + C content of 28 mol%. A search for open reading frames (ORFs) larger than 30 codons resulted in 18 potential protein-coding regions that were compared with current databases (Table 1). The ORFs were located in all three reading frames on the same DNA strand and had ATG as the translational start codon. The plasmid sequence was numbered arbitrarily starting at the first base of the inverted repeat in a palindromic sequence designated palSK1 (Fig. 1). The deduced amino acid sequences encoded by 13 ORFs displayed some similarities to known sequences (Table 1). The overall organization of the plasmid includes genes encoding a putative plasmid partitioning protein (ParA) and a putative adhesin (SkARP1), a region encoding putative Type IV secretion system-like components of conjugation machinery (ICEF-IA ORF15-like, ICEF-IA ORF16-like, and TraE protein with VirB4 domain similar to ICEF-II ORF17 of *Mycoplasma fermentans*), and a duplicated origin of transfer (*oriT*) region. These features will be discussed in detail below.

#### 3.1. Putative plasmid partitioning protein

ORF6 (bases 1602–2369) encodes a putative protein with ATPase activity related to Soj family proteins (COG1192) that are involved in chromo-

some and plasmid partitioning in other bacteria. A Pfam:ParA domain is predicted to occur at residues 82–193. ParA family ATPases are involved in bacterial plasmid partitioning, and an analysis of the similarity of ORF6 with described proteins in GenBank revealed similarity with the ATPase involved in chromosome partitioning in *Corynebacterium glutamicum* (NP\_600639; 29% identity over 236 amino acids), and the ParA protein of *Leptospira interrogans* (NP\_714527; 33% identity over 209 amino acids).

#### 3.2. *Spiroplasma* adhesin gene pathogenicity factor

ORF9 (bases 3740–6337) encodes a putative protein homolog of the SARP1 adhesion protein (P89) of *Spiroplasma citri*. The ORF9 putative protein, designated SkARP1, is a new member of the sarpin family of proteins described by Berg et al. (2001) and may function similarly as an adhesin of *S. kunkelii*. A putative signal peptide of 23 amino acids is present at the N-terminus of the preprotein. The predicted mature protein (SkARP1), after cleavage of the signal peptide, comprised 842 amino acids with a calculated molecular mass of 96 kDa. This contrasted with the mature SARP1 (here termed ScARP1) protein (GenBank Accession No. AJ297706) of *S. citri*, with a mass of 86 kDa (Berg et al., 2001). The SkARP1 signal peptide was the same in size and similar in amino acid sequence to that found in the ScARP1 preprotein. Low sequence similarity of the N-termini accounts for much of the size difference between the two mature proteins. The mature proteins also differed in numbers and identities of charged amino acids and have different isoelectric points (6.3 for SkARP1 and 5.5 for ScARP1).

It is known that ScARP1 is involved in the attachment of *S. citri* cells to gut cells of the insect vector, *Circulifer tenellus*, during early stages of infection in this host (Berg et al., 2001), and therefore it is quite likely that the *S. kunkelii* SkARP1 ortholog similarly is involved in attachment of *S. kunkelii* cells to gut tissue during infection of its insect vectors, which include *Dalbulus* spp. We postulate that amino acid sequence variations and differences in isoelectric points may reflect dif-

Table 1  
Properties of plasmid pSKU146 ORFs and their deduced products

ORF	Endpoints (nt)	G + C %	Ribosome-binding site/ start codon <sup>a</sup>	Product (aa/kDa)	Best BLAST hit <sup>b</sup> % identity (over aa)	Predicted function/ similar protein	Related COG <sup>c</sup> and gene name	Functional <sup>d</sup> category
ORF1	140–733	31.8	GAGTGGTTGGT <b><u>AAAGGAG</u></b> TTGATATT/ATG	197/23.6	NP_814286 28 (164)	Hypothetical protein	No related COG	
ORF2	789–893	12.4	AATTTGAAACAAT <b><u>ACGGAG</u></b> TAAAAT/ATG	34/4.0	No significant hits	Hypothetical protein	No related COG	
ORF3	895–1071	17.5	AAAAATGACATAAA <b><u>AAAGGGTAAA</u></b> /ATG	58/7.2	NP_703879 33 (56)	Hypothetical protein	No related COG	
ORF4	1034–1516	28.4	N.A. <sup>e</sup> /ATG	160/19.5	CAD58576 26 (125)	Hypothetical protein (TriL-like, MobC-like)	No related COG	N
ORF5	1095–1190	28.1	N.A./ATG	31/3.8	No significant hits	Hypothetical protein	No related COG	
ORF6	1602–2369	23.8	AATATATAAG <b><u>AAAGGA</u></b> AAACACAAA/ATG	255/29.7	NP_600639 29 (236)	Soj protein, PFAM ParA	COG1192 <i>Soj</i>	D
ORF7	2372–2626	26.7	TAGAACAAA <b><u>AAGGAGT</u></b> GATATAAAC/ATG	84/9.8	NP_228535 30 (82)	Hypothetical protein, RepA-like region	No related COG	
ORF8	2774–3088	23.5	TTAAAATAAATAT <b><u>AAGGAG</u></b> TAAATA/ATG	104/12.4	CAD20867 23 (89)	Hypothetical protein	No related COG	
ORF9	3740–6337	30.1	TTATTGTT <b><u>CAGAAAGGA</u></b> AAACAACG/ATG	865/98.4	CAC10363 71 (536)	SkARPI adhesion protein	No related COG <i>Skarp1</i>	
ORF10	6337–6606	37.4	AAAACAAC <b><u>AAGGGC</u></b> GGTGATGAATA/ATG	89/9.4	AAN85225 31 (69)	Hypothetical protein (ICEF-1A ORF15-like)	No related COG	
ORF11	6618–8180	32.1	CAATCACAAGTT <b><u>AGGCCGG</u></b> TGTAATT/ATG	520/57.6	CAC10364 87 (82)	Hypothetical protein (ICEF-1A ORF16-like)	No related COG	
ORF12	8170–10,692	28.2	AAAAAAGAT <b><u>AAAGGA</u></b> GACAAAAACA/ATG	840/96.9	AA074893 94 (840)	TraE protein with VirB4 domain <sup>f</sup>	COG3451 <i>TraE</i>	N
ORF13	10,696–11,034	28.6	TAACCTAGG <b><u>AAAGGA</u></b> AATATAGTTT/ATG	112/13.3	NP_040346 33 (95)	Hypothetical protein	No related COG	
ORF14	11,136–12,650	27.5	TCTTAAAAG <b><u>AAAGTG</u></b> GGTTTTTTAA/ATG	504/58.3	NP_114056 23 (425)	Mob protein with VirD4 domain	COG3505, 0433 <i>Mob</i>	N
ORF15	12,833–12,955	21.1	N.A./ATG	40/5.0	No significant hits	Hypothetical protein, membrane associated	No related COG	
ORF16	13,002–13,133	25.8	N.A./ATG	43/5.0	No significant hits	Hypothetical protein, membrane associated	No related COG	
ORF17	13,491–14,135	30.1	ACTATTATT <b><u>AAGGAG</u></b> GACAAATAAT/ATG	214/25.5	NP_701124 26 (143)	Hypothetical protein	No related COG	
ORF18	14,213–14,338	23.8	N.A./ATG	41/5.1	No significant hits	Hypothetical protein, membrane associated	No related COG	

<sup>a</sup> Bold underlined nucleotides denote purine-rich region within the putative ribosomal-binding site (RBS).

<sup>b</sup> GenBank accession number or protein ID of the best BLAST hit, followed by the percent identity between the query and the best hit.

<sup>c</sup> COG stands for Cluster of Orthologous Groups.

<sup>d</sup> Functional classification based on the result of COG search.

<sup>e</sup> N.A., not applicable; putative RBS not identified.

<sup>f</sup> Similar to ICEF-II ORF17 of *Mycoplasma fermentans*.




Fig. 1. Physical and genetic map of plasmid pSKU146. Putative protein-coding regions (open reading frames, ORFs) and direction of transcription are represented by open arrows. Overlapping ORFs are indicated by arrows in inner concentric circle. Most ORFs are numbered. ORF6, putative *Soj* plasmid partitioning protein. ORF9, SkARP1 adhesion-related protein. ORF10 and ORF11, hypothetical proteins (ICEF-1A ORF15-like and ORF16-like, respectively). ORF12, TraE-like protein with VirB4 domain. ORF13, hypothetical protein 4–Spiroplasma virus 1. ORF14, Mob protein with VirD4 domain. ORF7, similarity with replication proteins. ORF1-ORF5, ORF8, ORF15, ORF16, ORF17, and ORF18, hypothetical proteins, no related COGs. palSK1, 58 bp palindromic sequence (bases 1–29 abutting inverted inverted repeat, bases 14,587–14,615). IRSK, inverted repeat units potentially forming a hairpin structure downstream of the Mob gene (ORF14). Putative *oriTA* and *oriTB* regions are indicated by broken arrows. Cleavage sites of selected restriction endonucleases are indicated.

ferences in the ligand attachment sites of ScARP1 and SkARP1 in their respective insect hosts.

Based on work with *Enterococcus faecalis* (Hirt et al., 2000; Olmsted et al., 1991), it is possible that putative virulence factors SkARP1 and ScARP1 could be involved in formation of spiroplasma mating aggregates for conjugal DNA transfer. Studies of the aggregation substance (AS) surface protein encoded by *E. faecalis* virulence plasmid pCF10 indicate that AS is a virulence factor involved in the adhesion of bacterial cells to host tissues (Hirt et al., 2000), as well as a conjugation protein involved in the formation of a mating pair,

enhancing conjugal transfer of the plasmid (Olmsted et al., 1991). A homologous protein is encoded by *E. faecalis* plasmid pAD1 (Francia et al., 2001). It would be interesting to know whether the SkARP1 and ScARP1 putative virulence factors likewise are involved in the formation of spiroplasma mating aggregates leading to conjugal DNA transfer.

### 3.3. Components of conjugation machinery, Type IV secretion system-like proteins

Plasmid pSKU146 contains ORFs that encode proteins having similarity to conjugative elements and components of the bacterial Type IV secretion system. The deduced amino acid sequence encoded by ORF10 possesses a signal peptide (residues 1–34) and a single transmembrane region (residues 62–80). ORF10 displayed similarity with ICEF-IA ORF15 of *M. fermentans* (GenBank Accession No. AAN85225; 31% identity over 69 amino acids). The ICEF of *M. fermentans* is a non-self replicating, chromosomally integrating DNA that encodes proteins similar to known conjugation proteins and that is capable of being transferred during a conjugation-like event (Calcutt et al., 2002).

ORF11 (bases 6618–8180) encodes a hypothetical protein having similarity with a hypothetical protein of *S. citri* (GenBank Accession No. CAC10364; 87% identity over 82 amino acids) and similarity with several other proteins, including the integrative conjugal element ICEF-IA ORF16 of *M. fermentans* (GenBank Accession No. AAN85226; 27% identity over 147 amino acids). The deduced amino acid sequence encoded by ORF11 contains eight transmembrane regions (residues 36–58, 99–121, 153–175, 185–207, 260–282, 287–306, 327–349, and 353–372), and two regions of low complexity at residues 461–471 and 503–520, respectively. This predicted protein also shares topological similarities with membrane-trafficking proteins, in particular the bacterial Family 1 export proteins that export signal peptide-lacking proteins through the membrane.

ORF12 (bases 8170–10,692) encodes a TraE-like protein, a putative membrane-bound ATPase with a VirB4 domain related to Type IV secretory

pathway COG3451. This putative conjugal gene transfer protein is among the first to be found in class Mollicutes.

The protein encoded by ORF12 contains a signal peptide (residues 1–49), two transmembrane regions (50–72 and 79–93), and a low complexity region (244–253) predicted by SMART. The presence of a signal peptide and transmembrane regions is uncharacteristic of VirB4 proteins and may indicate that the ORF12 protein carries out additional functions. This protein has similarity with a putative membrane-bound ATPase of *S. kunkelii* (GenBank Accession No. [AAO74893](#); 94% identity over 840 amino acids), as well as to other GenBank listed proteins that include a TRSE-like protein of *Mycoplasma pulmonis* (GenBank Accession No. [NP\\_326214](#); 27% identity over 795 amino acids); protein ICEF-II ORF17 of the integrative, conjugative element of *M. fermentans* (ICEF) (GenBank Accession No. [AAN85276](#); 23% identity over 728 amino acids); a transfer complex protein TrsE of *Lactococcus lactis* plasmid pMRCO1 (GenBank Accession No. [NP\\_047296](#); 22% identity over 601 amino acids); a putative ATPase TraE of *Staphylococcus epidermis* (GenBank Accession No. [NP\\_765045](#); 21% identity over 522 amino acids); a putative pilus assembly protein of *Pseudomonas resinovorans* (GenBank Accession No. [NP\\_758682](#); 19% identity over 660 amino acids); and a sex pilus assembly protein of *Vibrio cholerae* (GenBank Accession No. [AAL59681](#); 19% identity over 409 amino acids).

Thus, the gene arrangement of pSKU146 ORFs 10 through 12 parallels the gene arrangement in ORFs 15 through 17 in the ICEF of *M. fermentans*, consistent with the concept that pSKU146 is also involved in DNA transfer. The results also raise the possibility that pSKU146 and the ICEF shared a common ancestor or donor that contributed pSKU146 ORF10-, ORF11-, and ORF12-like sequences to both elements.

ORF14 (bases 11,136–12,650) encodes another putative protein exhibiting significant similarity with bacterial conjugation proteins. Domains predicted in the deduced amino acid sequence were a signal peptide (residues 1–41), a transmembrane domain (residues 64–86), and a Pfam:TRAG domain (residues 103–491). The putative ORF14-encoded

protein exhibits the same domain architecture as conjugal transfer protein VirD4 (GenBank Accession No. [CAC15172](#)) of *Agrobacterium tumefaciens* and has similarity with the Mob protein (GenBank Accession No. [NP\\_114056](#)) of *Streptococcus mutans* (23% identity over 425 amino acids) and COGs related to VirD4, Type IV secretion pathway COG3505, and predicted ATPase COG0433. A putative Walker A nucleotide-binding site, <sup>139</sup>GT TGS<sup>146</sup>GKT<sup>146</sup>, is present in the deduced amino acid sequence of the pSKU146-encoded Mob.

Since bacterial Mob proteins are involved in plasmid mobilization and transfer to a recipient cell, and the TraG-TraD family proteins and VirD4 are bacterial coupling proteins involved in passage of conjugally transferred DNA through the mating channel and in protein translocation via the Type IV secretion system (Christie, 2001), the presence of ORFs encoding domains similar to Vir and Tra proteins in pSKU146 suggests that *Spiroplasma* spp. utilize components related to the Type IV system for macromolecule transfer. A recent report of a putative pilus-like structure in electron micrographs of spiroplasma cells (Ozbek et al., 2003) is consistent with this concept. We therefore hypothesize that pSKU146 can be conjugally transferred to recipient cells through a Type IV-related secretory machinery involving proteins encoded by the plasmid. The possibility that additional proteins related to components of a Type IV translocation system are encoded by the *S. kunkelii* genome is currently under investigation.

Compared to the Gram-negative *A. tumefaciens*, we anticipate that the *S. kunkelii* genome ([www.genome.ou.edu/spiro.html](http://www.genome.ou.edu/spiro.html)) encodes fewer Type IV secretion system-related putative conjugation proteins, but presumably encodes all that it requires for a functional translocation system. Notably, only four homologues of Type IV secretory pathway components have been previously found to be encoded by conjugative elements from unicellular Gram-positive bacteria (Grohmann et al., 2003). *Spiroplasma* is a member of the Firmicutes, Gram-positive low G + C bacteria. As noted previously (Waters, 1999), Gram-positive bacteria, lacking an outer membrane, may not require as many components as Gram-negative bacteria for formation of a conjugative bridge. *Spiroplasma*, lacking a wall and

bounded only by a single membrane, may require fewer components than are required for DNA transfer among bacteria having an outer membrane.

#### 3.4. Type IV secretion system-related genes in *Mollicutes*

As discussed above, the Type IV secretion system is an intercellular macromolecule transfer machinery that is evolutionarily related to conjugation systems of Gram-negative bacteria and also is implicated in bacterial pathogenesis for its capacity to deliver virulence effectors to eukaryotic host cells (Christie, 2001). In addition to *Spiroplasma* ([www.genome.ou.edu/spiro.html](http://www.genome.ou.edu/spiro.html) and this communication), new evidence has emerged indicating that Type IV secretion-related systems might exist among other Gram-positive, cell wall-less bacteria, since genes encoding components of such a system are found in the genomes of *Mycoplasma* species, and include the *M. pulmonis* TRSE-like protein that has similarity with the pSKU146 ORF12 putative protein as noted above. The TraE/TrsE family NTPase protein gene (Calcutt et al., 2002) and a putative TraG family NTPase gene in the integrative conjugal element (ICEF) of *M. fermentans* (GenBank Accession No. AY168953) encode proteins similar to the deduced amino acid sequence encoded by ORF12 of pSKU146 that contains a region similar to the Type IV secretion system-related VirB4 domain. A search for VirD4-related sequences among the completely sequenced genomes of Mollicutes reveals a sequence, related to the C-terminal portion of the plasmid pSKU146 putative Mob protein, in *Mycoplasma pneumoniae* that encodes a putative VirD4-like amino acid sequence (GenBank Accession No. MPN513). These findings seem to indicate that Vir-related proteins, Type IV translocation system-related elements, may be more common among wall-less bacteria than previously realized.

#### 3.5. Ribosomal-binding sites and coordinate regulation

Several of the ORFs in pSKU146 are in juxtaposition or overlap. The start codon of ORF10, encoding a hypothetical protein exhibiting similarity to

the *M. fermentans* mobile genetic element ICEF-IA ORF15, overlaps the termination codon of ORF9, which encodes a putative adhesin (SkARP1). The termination codon of ORF10 and the start of ORF11, encoding a protein with similarity to ICEF-IA ORF16, are separated by only 12 bases. The start of ORF12, encoding a TraE-like protein, overlaps the 3'-end of ORF11 by 10 bases. ORF13 begins 4 bases after the termination codon of ORF12, and ORF14, encoding a putative Mob protein, begins only 2 bases after the termination codon of ORF13. A putative ribosome-binding site (RBS) could be located upstream of each of the above ORFs, and in some cases is within the 3'-end of the respective ORF immediately upstream. Departure of some RBS from strict homology with the 3'-end of 16S rRNA may reflect translational regulation through modulating the formation of the translation initiation complex.

Based on the relative positions of the juxtaposed and overlapping ORFs, it is likely that at least some of these genes are coordinately expressed. It is possible that the *tra* (transfer)-related genes (ORFs 10, 11, 12, and 14) are co-transcribed with *skarp1* as a single operon. Plasmid R1 contains a polycistronic *tra*-operon (Koraimann and Högenauer, 1989), and co-transcription has been demonstrated for *tra* genes of plasmid pIP501 from *Streptococcus agalactiae* (Kurenbach et al., 2002). The *tra* genes in pSKU146 may be expressed under the influence of the *skarp1* adhesin gene (ORF9) promoter. This hypothesis suggests a possible relationship between the *skarp1* gene and the genes encoding components of the transfer machinery. In addition, the possible coordinate expression could imply that the adhesin protein functions not only in spiroplasma cell–host cell adhesion, but also in spiroplasma cell-to-cell contact for conjugation, as reported earlier in the case of plasmid-encoded adhesin of *E. faecalis* involved in contact between mating cells (Hirt et al., 2000; Olmsted et al., 1991; Wirth, 1994).

#### 3.6. Origin of transfer (*oriT*) and palindromic sequences

Plasmid pSKU146 contains two identical 175 bp direct repeat units, designated Repeat

Alpha (bases 377–551) and Repeat Beta (bases 1160–1334), in each of which is embedded the sequence ATCCTG, a sequence that is conserved at the *oriT* nick region among conjugative plasmids (Furuya and Komano, 2000; Herrera-Cervera et al., 1998). The ATCCTG in pSKU146 is part of an imperfect inverted repeat, that can potentially form a hairpin structure important for relaxosome formation and endonuclease protein binding that results in a single-stranded nick required for plasmid mobilization during conjugation. The ATCCTG putative nick region sequence in pSKU146 is identical to nick regions found in the IncP conjugative plasmid family (Waters, 1999) and is preceded 5 bases upstream by a 10-base AT-rich sequence. The putative nick site (*nic*) within this region likely lies between bases 494 and 495 and between bases 1277 and 1278 in *oriTA* and *oriTB*, respectively (Fig. 2). Repeat Alpha and Repeat Beta thus represent two putative *oriT* regions (*oriTA* and *oriTB*, respectively) that probably resulted from sequence duplication. Two *oriT* sequence regions have been reported in other plasmids (Avila et al., 1996; Becker and Meyer, 2003; Herrera-Cervera et al., 1998), including the conjugative virulence plasmid pAD1 of *E. faecalis* (Francia et al., 2001).

In pSKU146, *oriTA* is embedded in putative protein-coding region ORF1 (bases 140–733). *oriTB* is embedded in a putative protein-coding region (ORF4, bases 1034–1516) and spans most of ORF5 (bases 1095–1190). The region spanned by the two *oriT* regions and the sequence upstream of *oriTA* is characterized by a cluster of direct and inverted repeat sequences. The two *oriT* regions

each contain at least seven inverted repeat sequences (IR), including the putative nick region (nick) at IR5 (Fig. 2). Four of these inverted repeats are located upstream and two are located downstream of the putative nick region, resulting in a structural order as follows: *oriTA* IR1 $\alpha$ -IR2 $\alpha$ -IR3 $\alpha$ -IR4 $\alpha$ -*nick*-IR6 $\alpha$ -IR7 $\alpha$ -ORF2-ORF3-*oriTB* (IR1 $\beta$ -IR2 $\beta$ -IR3 $\beta$ -IR4 $\beta$ -*nick*-IR6 $\beta$ -IR7 $\beta$ ).

Interestingly, the A + T content of pSKU146 *oriTA* and *oriTB* (69%) is lower than the A + T content of the intervening sequence between the two *oriT* regions (76%) and of sequences flanking the putative double *oriT* region, as well as being lower than the A + T content of the plasmid (72%) and that of the spiroplasma genome (74%). A similar observation has been made for the *oriT* region of conjugative element pRS01 from *L. lactis* subsp. *lactis* ML3 (Mills et al., 1998). In pSKU146, the A + T content of sequences flanking the *oriTA*–*oriTB* region is 86% for bases 14,245–14,483 (located upstream of *oriTA*), 71% for bases 1–378 (from beginning of the second inverted repeat in palSK1 to the beginning of *oriTA*), 76% from the end of *oriTB* to the start of the *soj* gene, and 76% in the *soj* gene.

A prominent feature of pSKU146 is a palindromic sequence, palSK1, of 58 bp (bases 1–29 abutting 14,587–14,615) located upstream of ORF1 and consisting of two abutted, inverted repeats having the sequence GGTAGGTTGTTT ATTGGTTGTATTTATTGCAATAAATACAA CCAATAACAACCTACC. This sequence is similar in size to the palA (48 bp) and palB (49 bp) palindromes of *E. faecalis* conjugative plasmid pRE25 (GenBank Accession No.




Fig. 2. Nucleotide sequence and structure of putative *oriTA* region in plasmid pSKU146. Inverted repeat (IR) sequences are represented by opposing arrows. Paired bases are underlined. Direct repeats (DR) are indicated by non-opposing arrows. The putative nick region within the *oriT* region is in bold and double underlined. The putative nick site is indicated by a vertical arrow. An AT-rich sequence is indicated upstream of the putative nick region. *oriTA* and *oriTB* are separated by a 608 bp sequence and are identical in sequence (not shown).

X92945). A second prominent structural feature, designated IRSK, is located at base positions 12,810–12,855 and consists of two 20 bp inverted repeat units that are separated by 6 bp and have the sequence ACTAAAAAATAAAAACACCT ttatgAGGTGTTTTTATTTTTTAGT. The IRSK sequence potentially forms a hairpin structure 160 bases downstream of the Mob gene (ORF14). The start codon for ORF15 is located within the six bases separating the inverted repeats of IRSK; this ORF may not be functional, since a putative RBS could not be identified. Whether palSK1 or IRSK is involved in plasmid mobilization, as noted for some palindromes (Priebe and Lacks, 1989), remains to be determined.

No ORFs with significant sequence similarity to any *rep* genes previously associated with plasmid origins of replication in Gram-positive bacteria were identified. However, a low scoring PFAM HTH\_4 domain (amino acid positions 46–83) was identified in the putative protein product of ORF7, through a domain search using SMART. A subsequent BLASTP search, using the region spanning amino acids 46–83 as a query against the SMART protein database, revealed similarity between the 84-amino acid ORF7 protein and the 49-amino acid replication protein (RepA) (NP\_862029) from *L. lactis* plasmid pSH72 (39% identity in 38 a.a. overlap; 68% similarity in 38 a.a. overlap) as well as the 52-amino acid RepA protein (CAA52852) from *Lactobacillus fermentum* (28% identity in 38 a.a. overlap; 71% similarity in 38 a.a. overlap). In addition, a sequence similar to the plasmid pGT5/pC194 rolling circle plasmid family double-stranded origin (dso) sequence TTATCTTGATA occurs twice with a single base mismatch (bases 11,312–11,322, within ORF14 on the forward strand, and bases 13,324–13,334 on the reverse strand), and the core sequence CTTGATA is exactly matched four times (three times on the forward strand at bases 739–745, at 12,242–12,248 within TTATcaaCTTGATA in ORF14, and at 13,737–13,743 in ORF16, and once on the reverse strand at bases 13,803–13,809), in pSKU146. Finally, the putative *Salmonella* NTP1 plasmid origin of replication sequence AGGCGTT occurs once in pSKU146 at bases 4035–4041, within ORF9.

### 3.7. Hypothetical proteins and spiroplasma virus protein homolog

The putative protein product of ORF1 exhibited similarity (28% identity over 164 residues) to a hypothetical protein from *E. faecalis* V583 (GenBank Accession No. NP\_814286). Both ORF1 and ORF4 protein products exhibited similarities (ORF1 28% over 164 residues and ORF4 29% over 151 residues) to an unknown protein from the *E. faecalis* conjugative virulence plasmid pAD1 (GenBank Accession No. AAL59457), raising the possibility that these two ORFs arose from gene duplication. The ORF4 product also exhibited similarity (26% identity over 125 residues) to the TriL protein (GenBank Accession No. CAD58576) encoded by a cryptic plasmid that encodes a conjugative transfer system in *Yersinia enterocolitica* (Goelz et al., 2003; Strauch et al., 2003). This degree of similarity raises the possibility that the proteins encoded by these ORFs are the result of horizontal gene transfer and may be involved in horizontal transfer of plasmid pSKU146.

The ORF13 (bases 10,696–11,034) putative protein has similarities (36% identity over 95 residues, see Table 1) to a hypothetical protein of spiroplasma virus 1 (GenBank Accession No. NP\_040346). However, of the remaining ORFs (ORF2, ORF3, ORF8, and ORF17), the deduced amino acid sequences displayed little or no significant similarity to previously deposited GenBank sequences using BLASTP searches. Although they most likely are expressed since they are preceded by putative ribosome-binding sites (RBS) (Table 1), any function(s) associated with these putative proteins thus remain cryptic. The putative products of the only remaining ORFs (ORF5, ORF15, ORF16, and ORF18) exhibited no similarity to known sequences, and since clear ribosomal-binding sites were not observed, these ORFs may not be translated into functional proteins. However, there is mounting evidence that several organisms, including *M. pneumoniae* express genes that lack clear ribosomal-binding sites (Weiner III et al., 2000). Thus, since the putative protein products of ORF15, ORF16, and ORF18 each contain a transmembrane segment (ORF15

product residues 5–27, ORF16 product residues 4–23, and ORF18 product residues 7–29) predicted by SMART, it is likely that these ORFs result in expressed proteins that are membrane associated. It would be interesting to know whether they function, or have functioned, in macromolecule transfer in the spiroplasma or a progenitor.

In summary, plasmid pSKU146 harbors genes encoding putative proteins similar to known bacterial virulence factors and conjugative elements including Mob family proteins, origins of plasmid transfer, Vir domains, ICEF ORF-like proteins, and the ScARP1 adhesin of *S. citri*, implicating a role of pSKU146-encoded genes in conjugal cell-to-cell communication and pathogenesis. Conjugative and mobilizable plasmids have not been previously described in genus *Spiroplasma*, making the putative conjugative element pSKU146 the first of its type to be described in Spiroplasmataceae.

The finding of putative conjugative plasmid sequences in *S. kunkelii* raises some important questions. For example, it will be of interest to learn whether other *Spiroplasma* species carry the array of putative pathogenicity factors observed in pSKU146 and to understand how pSKU146 and similar plasmids may influence the biology of pathogenic spiroplasmas and contribute to an evolving spiroplasma diversity. One may also ask whether conjugative plasmids in spiroplasma may mediate genetic exchange with other genera of organisms, including phytoplasmas and *Liberobacter* spp.; phytoplasmas and *Liberobacter* spp. are wall-less and walled plant pathogens, respectively, that inhabit sieve cells of plant phloem tissue just as do spiroplasmas, but cannot be isolated in artificial culture. Some phytoplasmas even share both insect vector and plant host species with a spiroplasma. For example, maize bushy stunt (MBS) phytoplasma shares plant and insect hosts with *S. kunkelii*, and periwinkle virescence phytoplasma (beet leafhopper transmitted virescence agent, VR) shares plant and insect hosts with *S. citri* (Barros et al., 2001; Nault, 1980; Oldfield, 1984). Moreover, since *S. kunkelii* is a member of the spiroplasma-*Mycoplasma mycoides* clade (Weisburg et al., 1989), which contains *M. mycoides* and *Mycoplasma capricolum*, one may ask

whether similar plasmids play a role in the biology of mammalian pathogenic *Mycoplasma* spp., including the development and spread of antibiotic resistance.

## Acknowledgments

*Spiroplasma kunkelii* gene sequence data were provided by the *Spiroplasma kunkelii* Genome Sequencing Project funded by U.S. Department of Agriculture, Agricultural Research Service Project Number: 1275-22000-144-02. The data were made available by Bruce Roe, ShaoPing Lin, HongGui Jia, HongMin Wu, and Doris Kupfer (University of Oklahoma, Department of Chemistry and Biochemistry, Norman, Oklahoma 73019) and Robert E. Davis (U.S. Department of Agriculture-Agricultural Research Service, Molecular Plant Pathology Laboratory, Beltsville, MD 20705). We thank J. Fletcher (Oklahoma State University, Stillwater) for kindly supplying *Spiroplasma kunkelii* strain CR2-3X.

## References

- Altschul, S.F., Gish, W., Miller, W., Myers, E.W., Lipman, D.J., 1990. Basic local alignment search tool. *J. Mol. Biol.* 215, 403–410.
- Avila, P., Nunez, B., de la Cruz, F., 1996. Plasmid R6K contains two functional oriTs which can assemble simultaneously in relaxosomes in vivo. *J. Mol. Biol.* 261, 135–143.
- Barros, T.S.L., Davis, R.E., Resende, R.O., Dally, E.L., 2001. Design of a polymerase chain reaction for specific detection of corn stunt spiroplasma. *Plant Dis.* 85, 475–480.
- Becker, E.C., Meyer, R.J., 2003. Relaxed specificity of the R1162 nickase: a model for evolution of a system for conjugative mobilization of plasmids. *J. Bacteriol.* 185, 3538–3546.
- Berg, M., Melcher, U., Fletcher, J., 2001. Characterization of *Spiroplasma citri* adhesion related protein SARP1, which contains a domain of a novel family designated sarpin. *Gene* 275, 57–64.
- Birren, B., Green, E.D., Klapholz, S., Myers, R.M., Riethman, J., Roskams, J., 1999. *Genome Analysis: A Laboratory Manual*, Vol. 3, Cloning Systems. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, p. 648.
- Calcutt, M.J., Lewis, M.S., Wise, K.S., 2002. Molecular genetic analysis of ICEF, an integrative conjugal element that is present as a repetitive sequence in the chromosome of

- Mycoplasma fermentans* PG18. J. Bacteriol. 184, 6929–6941.
- Christie, P.J., 2001. Type IV secretion: intercellular transfer of macromolecules by systems ancestrally related to conjugation machines. Mol. Microbiol. 40, 294–305.
- Davis, R.E., Worley, J.F., 1973. Spiroplasma: motile, helical microorganism associated with corn stunt disease. Phytopathology 63, 403–408.
- Davis, R.E., Worley, J.F., Whitcomb, R.F., Ishijima, T., Steere, R.L., 1972. Helical filaments produced by a mycoplasmalike organism associated with corn stunt disease. Science 176, 521–523.
- Francia, M.V., Haas, W., Wirth, R., Samberger, E., Muscholl-Silberhorn, A., Gilmore, M.S., Ike, Y., Weaver, K.E., An, D.B., Clewell, D.B., 2001. Completion of the nucleotide sequence of the *Enterococcus faecalis* conjugative plasmid pAD1 and identification of a second transfer origin. Plasmid 46, 117–127.
- Furuya, N., Komano, T., 2000. Initiation and termination of DNA transfer during conjugation of IncI1 plasmid R64: roles of two sets of inverted repeat sequences within *oriT* in termination of R64 transfer. J. Bacteriol. 182, 3191–3196.
- Gasparich, G., 2002. Spiroplasmas: evolution, adaptation and diversity. Front. Biosci. 7, d619–d640. Available from: <<http://bioscience.igh.cnrs.fr/2002/v7/d/gaspari/fulltext.htm>> .
- Gasparich, G.E., Hackett, K.J., 1994. Characterization of a cryptic extrachromosomal element isolated from the mollusc *Spiroplasma taiwanense*. Plasmid 32, 342–343.
- Gasparich, G.E., Hackett, K.J., Clarke, E.A., Renaudin, J., Whitcomb, R.F., 1993. Occurrence of extrachromosomal deoxyribonucleic acids in spiroplasmas associated with plants, insects, and ticks. Plasmid 29, 81–93.
- Goelz, G., Knabner, D., Appel, B., Strauch, E., 2003. A conjugal type IV transfer system in *Yersinia enterocolitica* strains. Adv. Exp. Med. Biol. 149, 51–53.
- Grohmann, E., Muth, G., Espinosa, M., 2003. Conjugative plasmid transfer in Gram-positive bacteria. Microbiol. Mol. Biol. Rev. 67, 277–301.
- Herrera-Cervera, J.A., Sanjuan-Pinilla, J.M., Olivares, J., Sanjuan, J., 1998. Cloning and identification of conjugative transfer origins in the *Rhizobium meliloti* genome. J. Bacteriol. 180, 4583–4590.
- Hirt, H., Erlandsen, S.L., Dunny, G.M., 2000. Heterologous inducible expression of *Enterococcus faecalis* pCF10 aggregation substance *asc10* in *Lactococcus lactis* and *Streptococcus gordonii* contributes to cell hydrophobicity and adhesion to fibrin. J. Bacteriol. 182, 2299–2306.
- Kim, U.-J., Birren, B.W., Slepak, T., Mancino, V., Boysen, C., Kang, H.L., Simon, M.I., Shizuya, H., 1996. Construction and characterization of a human bacterial artificial chromosome library. Genomics 34, 213–218.
- Koraimann, G., Högenauer, G., 1989. A stable core region of the *tra* operon mRNA of plasmid R1-19. Nucleic Acids Res. 17, 1283–1297.
- Kurenbach, B., Grothe, D., Farias, M.E., Szewzyk, U., Grohmann, E., 2002. The *tra* region of the conjugative plasmid pIP501 is organized in an operon with the first gene encoding the relaxase. J. Bacteriol. 184, 1801–1805.
- Lee, I.-M., Davis, R.E., 1989. Serum-free media for cultivation of spiroplasmas. Can. J. Microbiol. 35, 1092–1099.
- Letunic, I., Goodstadt, L., Dickens, N.J., Doerks, T., Schultz, R., Mott, R., Ciccarelli, F., Copley, R.R., Ponting, C.P., Bork, P., 2002. Recent improvements to the SMART domain-based sequence annotation resource. Nucleic Acids Res. 30, 242–244.
- Mills, D.A., Phister, T.G., Dunney, G.M., McKay, L.L., 1998. An origin of transfer (*oriT*) on the conjugative element pRS01 from *Lactococcus lactis* subsp. *lactis* ML3. Appl. Environ. Microbiol. 64, 1541–1544.
- Nault, L.R., 1980. Maize bushy stunt and corn stunt: a comparison of disease symptoms, pathogen host ranges, and vectors. Phytopathology 70, 659–662.
- Nielsen, H., Engelbrecht, J., Brunak, S., von Heijne, G., 1997. Identification of prokaryotic and eukaryotic signal peptides and prediction of their cleavage sites. Protein Eng. 10, 1–6.
- Oldfield, G.N., 1984. Field ecology of *Spiroplasma citri* in western North America. Isr. J. Med. Sci. 20, 998–1001.
- Olmsted, S.B., Kao, S.M., van Putte, L.J., Gallo, J.C., Dunny, G.M., 1991. Role of the pheromone-inducible surface protein *Asc10* in mating aggregate formation and conjugal transfer of the *Enterococcus faecalis* plasmid pCF10. J. Bacteriol. 173, 7665–7672.
- Ozbek, E., Miller, S.A., Meulia, T., Hogenhout, S.A., 2003. Infection and replication sites of *Spiroplasma kunkelii* (class Mollicutes) in midgut and malpighian tubules of the leafhopper *Dalbulus maidis*. J. Invertebr. Pathol. 82, 167–175.
- Peterson, D.G., Tomkins, J.P., Frisch, D.A., Wing, R.A., Peterson, A.H., 2000. Construction of plant bacterial artificial chromosome (BAC) libraries: an illustrated guide. J. Agr. Genom. 5, Available from: <<http://www.ncgr.org/research/jag>> .
- Priebe, S.D., Lacks, S.A., 1989. Region of the streptococcal plasmid pMV158 required for conjugative mobilization. J. Bacteriol. 171, 4778–4784.
- Ranhand, J.M., Mitchell, W.O., Popkin, T.J., Cole, R.M., 1980. Covalently closed circular deoxyribonucleic acids in spiroplasmas. J. Bacteriol. 143, 1194–1199.
- Razin, S., Nur, I., Glaser, G., 1987. Spiroplasma plasmids. Isr. J. Med. Sci. 23, 678–682.
- Salvado, J.-C., Barroso, G., Labarère, J., 1989. Involvement of a *Spiroplasma citri* plasmid in erythromycin-resistance transfer. Plasmid 22, 151–159.
- Schultz, J., Milpetz, F., Bork, P., Ponting, C.P., 1998. SMART, a simple modular architecture research tool: identification of signaling domains. Proc. Natl. Acad. Sci. USA 95, 5857–5864.
- Strauch, E., Goelz, G., Knabner, D., Konietzny, A., Lanka, E., Appel, B., 2003. A cryptic plasmid of *Yersinia enterocolitica* encodes a conjugative transfer system related to the regions of CloDF13 Mob and IncX Pil. Microbiology 149, 2829–2845.

- Tatusov, R.L., Natale, D.A., Garkavtsev, I.V., Tatusova, T.A., Shankavaram, U.T., Rao, B.S., Kiryutin, B., Galperin, M.Y., Federova, N.D., Koonin, E.V., 2001. The COG database: new developments in phylogenetic classification of proteins from complete genomes. *Nucleic Acids Res.* 29, 22–28.
- Waters, V.L., 1999. Conjugative transfer in the dissemination of beta-lactam and aminoglycoside resistance. *Front. Biosci.* 4, d416–d439.
- Weiner III, J., Herrmann, R., Browning, G.F., 2000. Transcription in *Mycoplasma pneumoniae*. *Nucleic Acids Res.* 28, 4488–4496.
- Weisburg, W.G., Tully, J.G., Rose, D.L., Petzel, J.P., Oyaizu, D., Yang, D., Mandelco, L., Sechrest, J., Lawrence, T.G., Van Etten, J. et al., 1989. A phylogenetic analysis of the mycoplasmas: basis for their classification. *J. Bacteriol.* 171, 6455–6467.
- Whitcomb, R.F., Williamson, D.L., 1975. Helical wall-free prokaryotes in insects: multiplication and pathogenicity. *Ann. N. Y. Acad. Sci.* 266, 260–275.
- Wirth, R., 1994. The sex pheromone system of *Enterococcus faecalis*. More than just a plasmid-collection mechanism?. *Eur. J. Biochem.* 222, 235–246.