University of Nebraska - Lincoln
DigitalCommons@University of Nebraska - Lincoln

Cornhusker Economics | Agricultural Economics Department

4-30-2008

The Economic Optimal Weight for the Replacement Beef Female

Matt Stockton
University of Nebraska - Lincoln, mstockton2@unl.edu

Roger K. Wilson
rwilson6@unl.edu, rwilson6@unl.edu

Follow this and additional works at: http://digitalcommons.unl.edu/agecon_cornhusker

Part of the Agricultural and Resource Economics Commons

Stockton, Matt and Wilson, Roger K., "The Economic Optimal Weight for the Replacement Beef Female" (2008). *Cornhusker Economics.* Paper 361.
http://digitalcommons.unl.edu/agecon_cornhusker/361

This Article is brought to you for free and open access by the Agricultural Economics Department at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Cornhusker Economics by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

CORNHUSKER ECONOMICS

University of Nebraska–Lincoln Extension

Nebraska Lincoln

April 30, 2008

Institute of Agriculture & Natural Resources
Department of Agricultural Economics
http://www.agecon.unl.edu/Cornhuskereconomics.html

The Economic Optimal Weight for the Replacement Beef Female

Market Report	Yr Ago	4 Wks Ago	4/25/08
Livestock and Products, Weekly Average			
Nebraska Slaughter Steers, 35-65% Choice, Live Weight	$96.26	$87.93	$92.70
Nebraska Feeder Steers, Med. & Large Frame, 550-600 lb.	128.21	121.20	123.56
Nebraska Feeder Steers, Med. & Large Frame 750-800 lb.	108.57	99.75	105.64
Choice Boxed Beef, 600-750 lb. Carcass	161.54	139.99	154.49
Western Corn Belt Base Hog Price Carcass, Negotiated	69.52	54.02	71.34
Feeder Pigs, National Direct 50 lbs, FOB	72.76	46.31	50.00
Pork Carcass Cutout, 185 lb. Carcass, 51-52% Lean	74.26	56.50	70.71
Slaughter Lambs, Ch. & Pr., Heavy, Wooled, South Dakota, Direct	91.75	92.95	83.50
National Carcass Lamb Cutout, FOB	243.47	257.11	250.92
Crops, Daily Spot Prices			
Wheat, No. 1, H.W. Imperial, bu.	4.73	9.52	7.95
Corn, No. 2, Yellow Omaha, bu.	3.44	5.27	5.49
Soybeans, No. 1, Yellow Omaha, bu.	6.73	11.82	12.81
Grain Sorghum, No. 2, Yellow Dorchester, cwt.	5.55	9.04	9.32
Oats, No. 2, Heavy Minneapolis, MN , bu.	2.78	3.54	3.96
Hay			
Alfalfa, Large Square Bales, Good to Premium, RFV 160-185 Northeast Nebraska, ton.	135.00	135.00	*
Alfalfa, Large Rounds, Good Platte Valley, ton.	92.50	85.00	*
Grass Hay, Large Rounds, Premium Nebraska, ton.	90.00	*	*

* No market.

Determining the optimal weight for breeding replacement heifers has been a goal of researchers for decades. It is an important question because replacing cows in the beef herd is expensive.

Finding the answer is a complex problem because of the inter-related nature of the economic and physiological variables. Patterson et al. (1992) may have expressed the nature of the problem best when he said, "Although puberty in the female and the events that precede its onset may be fixed from a physiological standpoint, the animal and management environment in which it is placed are in constant transition." This indicates that the optimum depends on economic, as well as the biological factors.

Recent studies conducted by University of Nebraska researchers (Funston and Deutscher 2004, and Martin et al. 2007) at the West Central Research and Extension Center (WCREC) have shown the fertility rates between groups of heifers fed to different average sizes prior to breeding were not statistically different, while the feed costs for the heavier heifers was greater than that of the lighter ones. Thus, the economic analyses in these studies were based on the development cost of cattle groups. This study also suggested that optimal heifer size was less than the traditionally accepted 64 to 65 percent mature body weight (PMBW).

The data collected in the above WCREC studies was used to conduct a rigorous economic analysis of optimal breeding weight using classical economic optimization theory. A profit equation was the central tool, as suggested by Fuez (1991). This profit equation consisted of five revenue equations and three cost equations, all of which were affected by the heifer's pre-breeding size, PMBW. The complex nature of the profit equation required a numerical methodology to derive the optimal PMBW.

The revenue equations included sale of non-pregnant heifers, sale of cows that did not have a calf after the calving season, sale of cows that did not rebreed, sale of calves born to the heifers, and the value of pregnant three year-old cows (see Figure 1).

The three cost equations included the initial costs of securing the heifers, the cost of feeding them to the appropriate PMBW,

IANR

Extension is a Division of the Institute of Agriculture and Natural Resources at the University of Nebraska–Lincoln cooperating with the Counties and the U.S. Department of Agriculture.

University of Nebraska Extension educational programs abide with the non-discrimination policies of the University of Nebraska–Lincoln and the United States Department of Agriculture.

and the cost of dystocia (calving difficulty) as seen in Figure 2. Other costs, such as opportunity cost and reduced pregnancies, were implicitly included in the revenue calculation.

These cost and revenue equations were developed using both ordinary least squares (OLS) and limited dependant variable (PROBIT) regressions. For instance, the relationship between PMBW and expected pregnancy rates, which decreased as PMBW decreased, was used to predict the number of heifers sold at the first pregnancy check.

Of all relationships among the many factors, only a few will be mentioned here. A complete report can be obtained from the authors.

Smaller PMBWs reduced the sale weights of all subsequent groups. The lower sale weights at first pregnancy check, and fall calf sales produced from these smaller cows increased the price per pound received, since lighter weight animals usually sell for a premium on a pound basis, but resulted in decreased revenue. A lower PMBW increased the incidence of dystocia among first calf heifers. It also led to a lower second pregnancy rate, resulting in increased cost.

The optimal biological reproductive rate was achieved at 62.9 PMBW, which is lower than the traditionally accepted recommendation. This difference may be due to the way PMBW is calculated. While others used an average herd size for each PBMW calculation, we used the actual mature weights for individual animals.

When the economic factors of revenue and cost are considered, the profit maximizing PMBW falls to 59.7. The implication is that the cost of obtaining a larger animal (62.8) is less than the value returned for that size, giving the smaller heifer the economic advantage (see Figure 3).







Matt Stockton,
(308) 696-6713
Assistant Professor
and Extension Economist
mstockton2@unl.edu

Roger Wilson,
(308) 696-6738
Research Analyst
rwilson6@unl.edu

West Central Research &
Extension Center
University of Nebraska–Lincoln

References Cited:

Feuz, D.M. 1991. "The Optimal Prebreeding Target Weight for Replacement Beef Heifers." Abstract. *Western Journal of Agricultural Economics*. 16(2):461.

Funston, R.N. and G.H. Deutscher. 2004. "Comparison of Target Breeding Weight and Breeding Date for Replacement Beef Heifers and Effects on Subsequent Reproduction and Calf Performance." *Journal of Animal Science*. 82:3094-3099.

Martin, J.L., K.W. Creighton, J.A. Musgrave, T.J. Klopfenstein, R.T. Clark, D.C. Adams and R.N. Funston. 2008. "Effect of Pre-breeding Body Weight or Progestin Exposure before Breeding on Beef Heifer Performance through the Second Breeding Season." *Journal of Animal Science*. 86:451-459.

Patterson, D.J., R.C. Perry, G.H. Kiracofe, R.A. Bellows, R.B. Staigmiller and L.R. Corah. 1992. "Management Considerations in Heifer Development and Puberty." *Journal of Animal Science*. 70:4018-4035.

Patterson, T., R. Clark, R. Salverson, W. Fahsholtz and T. Line. 2005. "Heifer Development: Revisiting Target Weights and Management Approaches." *Proceedings: The Range Beef Cow Symposium XIX*. December 6-8, 2005. Rapid City, South Dakota.