

2014

Grow them and we will come for the feast

Tiffany Heng-Moss

University of Nebraska-Lincoln, thengmoss2@unl.edu

Jeff Bradshaw

University of Nebraska-Lincoln, jbradshaw2@unl.edu

Kyle Koch

University of Nebraska-Lincoln

Travis Prochaska

University of Nebraska-Lincoln

Teresa Donze-Reiner

University of Nebraska-Lincoln

See next page for additional authors

Follow this and additional works at: <http://digitalcommons.unl.edu/entomologyfacpub>

Heng-Moss, Tiffany; Bradshaw, Jeff; Koch, Kyle; Prochaska, Travis; Donze-Reiner, Teresa; and Sarath, Gautam, "Grow them and we will come for the feast" (2014). *Faculty Publications: Department of Entomology*. 365.

<http://digitalcommons.unl.edu/entomologyfacpub/365>

This Article is brought to you for free and open access by the Entomology, Department of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Publications: Department of Entomology by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors

Tiffany Heng-Moss, Jeff Bradshaw, Kyle Koch, Travis Prochaska, Teresa Donze-Reiner, and Gautam Sarath

Grow them and we will come for the feast



Tiffany Heng-Moss, Jeff Bradshaw, Kyle Koch, Travis Prochaska, and Teresa Donze-Reiner, University of Nebraska at Lincoln, Lincoln, NE, USA
Gautam Sarath, Forage and Bioenergy Research Unit, USDA-ARS, Lincoln, NE, USA
Contact: Gautam.Sarath@ars.usda.gov

Sustainable production of switchgrass and other bioenergy grasses will require effective pest management. Identification of potential insect pests and detailed characterization of the plant-insect interaction will better enable us to address emergent insect pests in production fields. An added uncertainty is how manipulation of plants for improved quality (e.g. lower lignin) will affect plant resistance to insect herbivory. Plants can utilize different mechanisms to defend against chewing versus piercing-sucking insects; however, some basal plant responses appear to be shared across diverse biotic stressors. The range and modulation of these responses are beginning to be addressed for several of the temperate, perennial, warm-season grasses that are designated as biomass crops. Other systems have demonstrated the need to become proactive in these studies. For instance, buffalograss, *Buchloë dactyloides* (Nuttall) Engelm., was first introduced as a low-maintenance turf species. However, a number of important insect pests were documented within a relatively short time frame.^{1,2} Similarly, several recent reports have been published for switchgrass (*Panicum virgatum* L.) and Miscanthus (*Miscanthus x giganteus*) that indicate the presence of insect pests in production fields, and the overall susceptibility of these species to insect herbivory.³⁻⁹

A number of aphid species can also feed on these perennial grasses.¹⁰ A more detailed evaluation of the suitability and categories of resistance present in tetraploid switchgrasses to feeding by two potential aphid pests have been conducted recently.^{11,12} Collectively, these studies demonstrate that the upland switchgrass cultivar, Summer, was generally susceptible to both the greenbug (*Schizaphis graminum*) and to the yellow sugarcane aphid (*Sipha flava*). The lowland switchgrass cultivar, Kanlow was highly resistant to both aphids, indicative of a strong antibiosis response. Interestingly, a population derived

Sustainable production of switchgrass and other bioenergy grasses will require effective pest management

from crossing Kanlow x Summer plants exhibited tolerance to the yellow sugarcane aphid. These results demonstrate that multiple resistance mechanisms are present in switchgrasses and understanding these mechanisms could be important to the development of new germplasm with enhanced resistance to a broad array of insect pests. One of the most effective and sustainable strategies for controlling insect pests has been the development of insect-resistant plants. Studies cited here, and those in progress in many other research laboratories, should provide a foundation for development of sustainable pest management strategies that offer improved environmental safety and minimal human health risk.

References

1. Baxendale FP, Heng-Moss TM and Riordan TP, *Blissus occiduus* (Hemiptera : Lygaeidae): A chinch bug pest new to buffalograss turf. *J Econ Entomol* **92**:1172–1176 (1999).
2. Heng-Moss TM, Baxendale FP, Riordan TP and Foster JE, Evaluation of buffalograss germplasm for resistance to *Blissus occiduus* (Hemiptera : Lygaeidae). *J Econ Entomol* **95**:1054–1062 (2002).
3. Dowd PF and Johnson ET, Differential resistance of switchgrass *Panicum virgatum* L. lines to fall armyworms *Spodoptera frugiperda* (J. E. Smith). *Genet Resour Crop Evol* **56**:1077–1089 (2009).
4. Dowd PF, Sarath G, Mitchell RB, Saathoff AJ and Vogel KP, Insect resistance of a full sib family of tetraploid switchgrass *Panicum virgatum* L. with varying lignin levels. *Genet Resour Crop Evol* **60**:975–984 (2013).
5. Mitchell R, Vogel KP and Sarath G, Managing and enhancing switchgrass as a bioenergy feedstock. *Biofuel Bioprod Bioref* **2**:530–539 (2008).
6. Nability PD, Orpet R, Miresmailli S, Berenbaum MR and Delucia EH, Silica and nitrogen modulate physical defense against chewing insect herbivores in bioenergy crops *Miscanthus x giganteus* and *Panicum virgatum* (Poaceae). *J Econ Entomol* **105**:878–883 (2012).
7. Prasifka JR, Bradshaw JD, Boe AA, Lee D, Adamski D and Gray ME, Symptoms, Distribution and abundance of the stem-boring caterpillar, *Blastobasis reartella* (Dietz), in switchgrass. *Bioenerg Res* **3**:238–242 (2010).
8. Prasifka JR, Bradshaw JD and Gray ME, Potential biomass reductions to *Miscanthus x giganteus* by stem-boring caterpillars. *Environ Entomol* **41**:865–871 (2012).
9. Prasifka JR, Bradshaw JD, Meagher RL, Nagoshi RN, Steffey KL and Gray ME, Development and feeding of fall armyworm on *Miscaninus x giganteus* and switchgrass. *J Econ Entomol* **102**:2154–2159 (2009).
10. Burd JD, Prasifka JR and Bradshaw JD, Establishment and host effects of cereal aphids on switchgrass (*Panicum virgatum* L.) cultivars. *Southwest Entomol* **37**:115–122 (2012).
11. Koch KG, Fithian R, Heng-Moss TM, Bradshaw JD and Sarath G, Evaluation of tetraploid switchgrass populations (*Panicum virgatum* L.) for host suitability and differential resistance to four cereal aphids. *J Econ Entomol* (in press).
12. Koch KG, Heng-Moss TM, Bradshaw JD and Sarath G, Categories of resistance to greenbug and yellow sugarcane aphid (Homoptera: Aphididae) in three tetraploid switchgrass populations. *Bioenerg Res* (in press).

Gautam Sarath

Forage and Bioenergy Research Unit,
USDA-ARS, Lincoln, NE, USA

DOI: 10.1002/bbb.1477