

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

USDA National Wildlife Research Center - Staff
Publications

U.S. Department of Agriculture: Animal and Plant
Health Inspection Service

February 2004

Response of Canada Geese to a Dead Goose Effigy

Thomas W. Seamans

USDA APHIS Wildlife Services, National Wildlife Research Center

Glen E. Bernhardt

USDA APHIS Wildlife Services, National Wildlife Research Center

Follow this and additional works at: http://digitalcommons.unl.edu/icwdm_usdanwrc



Part of the [Environmental Sciences Commons](#)

Seamans, Thomas W. and Bernhardt, Glen E., "Response of Canada Geese to a Dead Goose Effigy" (2004). *USDA National Wildlife Research Center - Staff Publications*. 384.

http://digitalcommons.unl.edu/icwdm_usdanwrc/384

This Article is brought to you for free and open access by the U.S. Department of Agriculture: Animal and Plant Health Inspection Service at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in USDA National Wildlife Research Center - Staff Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Response of Canada Geese to a Dead Goose Effigy

Thomas W. Seamans and Glen E. Bernhardt

USDA APHIS Wildlife Services, National Wildlife Research Center, Ohio Field Station, Sandusky, Ohio

ABSTRACT: The North American Canada goose population increased at a rate of 10.5% per year, 1966 - 2001. Canada geese rank as the third most hazardous species in regards to collisions with aircraft. Sound Canada goose management tools are critical for a safer airport environment. We conducted field evaluations of a Canada goose effigy during the breeding season with territorial pairs and in late summer with post-fledging flocks to determine if geese were deterred by the effigy. No difference in territorial pairs was found between pretreatment and treatment periods for Canada geese when goose effigies were placed within their territories. In post-fledging flocks, the mean number of geese observed during pretreatment (74.9 ± 12.9), treatment (14.8 ± 4.5), and posttreatment (53.6 ± 14.2) periods differed ($P < 0.01$). There was no difference ($P = 0.56$) between the mean number of geese observed during a second round of 5-day pretreatment (58.7) and 5-day second round treatment (43.7) periods. By itself, the goose effigy was not effective as a Canada goose deterrent after approximately 5 days. However, this effigy may have some potential in an integrated goose control program conducted outside of the breeding season. Further evaluation of the effigy as part of an integrated Canada goose control program is recommended.

KEY WORDS: bird damage control, *Branta canadensis*, Canada goose, deterrents, effigy

Proc. 21st Vertebr. Pest Conf. (R. M. Timm and W. P. Gorenzel, Eds.)
Published at Univ. of Calif., Davis. 2004. Pp. 104-106.

INTRODUCTION

Long term population trends from North American Breeding Bird Survey (BBS) data (1966 - 2002) show an increase of 10.4% per year ($P < 0.01$) for Canada geese (*Branta canadensis*) populations in North America (Sauer et al. 2003). The giant Canada goose (*B. c. maxima*) population in the Mississippi flyway has increased from about 800,000 in 1993 to about 1.5 million in 2000 (U.S. Fish and Wildlife Service 2000). Ankney (1996) noted that it is not possible to predict when the giant Canada goose population will stop increasing.

Wildlife-strikes cause serious safety hazards to aircraft. Wildlife strikes cost civil aviation at least \$489.8 million annually in the United States (Cleary et al. 2003). Canada geese rank as the third most hazardous species in regards to collisions with aircraft (Dolbeer et al. 2000). From 1990 to 2002, geese were involved in 1,027 strikes with civil aircraft and caused \$351 million in total costs (Cleary et al. 2003). In September 1995, 24 people were killed and a \$190-million aircraft was destroyed when an AWACS aircraft crashed on takeoff at Elmendorf Air Force Base, Alaska, after striking Canada geese (Wright 1997). Sound management techniques that reduce goose numbers in and around airports are therefore critical for safe airport operations.

Large-scale killing of nuisance birds is often undesirable or impractical (Dolbeer 1986, 1998; Dornbush et al. 1996, Smith et al. 1999); thus, there is considerable demand for effective nonlethal techniques to deter bird use of problem sites. Numerous harassment and frightening techniques for reducing conflicts involving birds are available (Solman 1994, Cleary 1994, Dolbeer et al. 1995). Many of these techniques are expensive, ineffective, require multiple years to achieve desired results, produce temporary results, or have not been evaluated quantitatively. Realistic dead bird effigies of gulls (*Larus* spp.) and turkey vultures (*Cathartes aura*)

have shown promise as species-specific frightening devices (Saul 1967, Stout et al. 1975, Stout and Schwab 1979, Stout and Schwab 1980, Seamans et al. 2000, Tillman et al. 2002). Currently, a device called the Dead Goose Decoy is marketed as a non-lethal method to scare geese away from designated areas. This device consists of a plastic Canada goose decoy that has the form and appearance of a dead goose. No studies on the efficacy of the device have been published in peer-reviewed journals or proceedings. Our goal was to evaluate the efficacy of this Canada goose effigy.

METHODS

Territorial Pairs

This study was conducted from March to April 2001 on the 2,200-ha National Aeronautics and Space Administration's Plum Brook Station (PBS) in Erie County, Ohio. Eight territorial pairs of Canada geese were located on 8 separate ponds (≤ 0.4 ha) on PBS. Counts of geese were conducted for 7 days at about the same time each day to establish the consistent use of each pond by at least one pair of Canada geese. Four of the 8 ponds were then randomly selected to receive 2 goose effigies. Counts of geese on each pond were again conducted as during the pretreatment period for 7 days.

Because territorial Canada geese maintain their territory and generally do not leave their territory for another occupied territory, the control and treated pairs may be considered as independent. The change in numbers of geese using the ponds was compared using *t*-tests.

Post-fledging flocks

During August through September 2002, we located 6 ponds (0.4 - 2.0 ha) in Erie and Huron Counties, Ohio that were actively used by Canada geese. We counted geese on each pond or within 25 m of the pond between 1300 and 1600 hrs for 5 consecutive days (pretreatment). Two

days following the last pretreatment count, at least 2 effigies per 0.4 ha were placed between 0800 and 1100 hrs in each pond as per the manufacturer's suggestions. Counts were conducted as during pretreatment for 5 consecutive days (treatment). At the end of the 5-day treatment period, effigies were removed and geese were counted on the ponds for 5 consecutive days (posttreatment). The mean number of geese using all ponds was compared between periods using Kruskal-Wallis analysis of variance (Statistix7 2000).

Following the posttreatment period, 4 ponds were selected to receive effigies for a second time. At the 2 ponds not retested, goose use had become too inconsistent to effectively test the control technique. Counts were conducted as during the earlier portions of the study but continued until Canada goose numbers were similar to the posttreatment numbers. The change in numbers of geese using the ponds during this portion of the test was compared using *t*-tests.

RESULTS

Territorial Pairs

There was no difference ($t=0.66$; 49 df; $P=0.51$) at the 4 control ponds in the mean (\pm SE) Canada goose numbers between pretreatment (2.1 ± 0.5) and treatment (2.7 ± 0.7) periods. At the 4 treated ponds, there also was no difference ($t=0.52$; 51 df; $P=0.61$) in mean Canada goose numbers between pretreatment (1.1 ± 0.1) and treatment (0.9 ± 0.2) periods.

Post-fledging flocks

The mean number (\pm SE) of geese observed on the 6 ponds during pretreatment (74.9 ± 12.9), treatment (14.8 ± 4.5), and posttreatment (53.6 ± 14.2) periods differed ($W=17.65$; $P<0.01$). There was no difference ($Z=0.58$; $P=0.56$) between the mean number of geese observed on the 4 ponds during the second round 5-day pretreatment (58.7 ± 20.7) and 5-day second round treatment (43.7 ± 15.6) periods.

DISCUSSION

Territorial pairs of Canada geese showed no response to the goose effigies. The manufacturer claims that geese will abandon nests and eggs when a decoy is placed near a pair's nest. We did not observe this behavior at any of our 4 treatment ponds. However, during the post-fledging period, the presence of goose effigies had an initial repellent effect at all sites tested. Canada geese were observed either flying towards treated ponds and then flaring away, or landing on the water only to flush off of the pond within 30 seconds of landing. By the end of the first 5-day treatment period, geese were generally returning to the pond but were staying at least 25 m away from the effigies. During the second 5-day treatment period, geese were observed swimming next to or between the pairs of effigies within 1 to 3 days of effigy placement.

Effective Canada goose management programs generally require an integrated approach to be ultimately successful (Booth 1994, Smith et al. 1999). The presence of goose effigies may enhance other control techniques, such as pyrotechnics that simulate gunfire (danger), lasers

(Blackwell et al. 2002), and chemical repellents (Dolbeer et al. 1998). The short-term (1 week) use of effigies at the start of an integrated control program to disperse Canada geese from an airfield or other site should prove useful.

Further experiments with goose effigies may include use of pyrotechnics and lasers to determine whether, if used in combination, the effectiveness of these techniques might be enhanced. Also, the use of lethal control could be added to see if a combination of all 4 techniques would create effective control.

ACKNOWLEDGEMENTS

This study was supported by the Federal Aviation Administration. Opinions expressed in this study do not necessarily reflect current FAA policy decisions governing the control of wildlife on or near airports. We thank Firelands College at Bowling Green State University, K. Conger, S. Deehr, J. Giese, the Ohio Veterans Home, and G. Palm for access to study sites. We thank R. Bush, L. Brohl, Z. Patton and R. White for field support.

LITERATURE CITED

- ANKNEY, C. D. 1996. An embarrassment of riches: too many geese. *J. Wildl. Manage.* 60:217-223.
- BLACKWELL, B. F., G. E. BERNHARDT, AND R. A. DOLBEER. 2002. Lasers as non-lethal avian repellents. *J. Wildl. Manage.* 66:250-258.
- BOOTH, T. H. 1994. Bird dispersal techniques. Pp. E19-E23 *in*: S. E. Hygnstrom, R. M. Timm, and G. E. Larson (Eds.), *Prevention and Control of Wildlife Damage*. Cooperative Extension, Institute of Agriculture and Natural Resources, University of Nebraska, Lincoln, NE.
- CLEARY, E. C. 1994. Waterfowl. Pp. E139-E155 *in*: S. E. Hygnstrom, R. M. Timm, and G. E. Larson (Eds.), *Prevention and Control of Wildlife Damage*. Cooperative Extension, Institute of Agriculture and Natural Resources, University of Nebraska, Lincoln, NE.
- CLEARY, E. C., S. E. WRIGHT, AND R. A. DOLBEER. 2003. Wildlife strikes to civilian aircraft in the United States, 1990-2002. Special Serial Report No. 9, National Wildlife Strike Database, Federal Aviation Administration, Washington, D.C. June 2003. 60 pp.
- DOLBEER, R. A. 1986. Current status and potential of lethal means of reducing bird damage in agriculture. *Int. Ornithol. Congr.* 19:474-483.
- DOLBEER, R. A. 1998. Population dynamics: the foundation of wildlife damage management for the 21st Century. *Proc. Vertebr. Pest Conf.* 18:2-11.
- DOLBEER, R. A., N. R. HOLLER, AND D. W. HAWTHORNE. 1995. Identification and control of wildlife damage. Pp. 474-506 *in*: T. A. Bookhout (Ed.), *Research and Management Techniques for Wildlife and Habitats*. The Wildlife Society, Bethesda, MD.
- DOLBEER, R. A., T. W. SEAMANS, B. F. BLACKWELL, AND J. L. BELANT. 1998. Anthraquinone formulation (Flight Control) shows promise as avian feeding repellent. *J. Wildl. Manage.* 62:1558-1564.
- DOLBEER, R. A., S. E. WRIGHT, AND E. C. CLEARY. 2000. Ranking the hazard level of wildlife species to aviation using the National Wildlife Strike Database. *Wildl. Soc. Bull.* 28:372-378.
- DORNBUSH, C., G. FEIGELSON, D. GRUSKIN, B. HEDGES, AND A. TURNER. 1996. Non-lethal controls for "resident" Canada

- geese. A report presented by the executive committee of the Canada Geese Citizens Advisory Committee, Rockland County, New York.
- SAUL, E. K. 1967. Birds and aircraft: a problem at Auckland's new international airport. *J. Royal Aero. Soc.* 71:366-375.
- SAUER, J. R., J. E. HINES, AND J. FALLON. 2003. The North American breeding bird survey, results and analysis 1966 - 2002. Version 2003.1, U. S. Geological Survey, Patuxent Wildlife Research Center, Laurel, MD.
- SEAMANS, T. W., S. W. YOUNG, AND J. D. CEPEK. 2000. Response of roosting turkey vultures to a hanging vulture effigy. Federal Aviation Administration Interim Report DTFA03-99-X-90001, Task 3, Experiment 5, Atlantic City, NJ.
- SMITH, A. E., S. R. CRAVEN, AND P. D. CURTIS. 1999. Managing Canada geese in urban environments. Jack Berryman Institute Publication 16, and Cornell University Cooperative Extension, Ithaca, NY.
- SOLMAN, V. E. F. 1994. Gulls. Pp. E49-E52 *in*: S. E. Hygnstrom, R. M. Timm, and G. E. Larson (Eds.), *Prevention and Control of Wildlife Damage*. Cooperative Extension, Institute of Agriculture and Natural Resources, University of Nebraska, Lincoln, NE.
- STATISTIX7. 2000. *Statistix 7 User's Manual*. Analytical Software, Tallahassee, FL.
- STOUT, J. F., W. H. GILLETT, J. L. HAYWARD, JR., AND C. J. AMLANDER, JR. 1975. Dispersal of seagulls in an airdrome environment. Air Force Weapons Laboratory Final Report AFWL-TR-74-324, Kirtland Air Force Base, NM.
- STOUT, J. F., AND E. R. SCHWAB. 1979. Behavioral control of seagulls at Langley Air Force Base. Pp. 96-110 *in*: W. B. Jackson, S. S. Jackson, and B. A. Jackson (Eds.), *Proc. Eighth Bird Control Seminar*, Bowling Green St. University, Bowling Green, OH.
- STOUT, J. F., AND E. R. SCHWAB. 1980. Telemetry of heart rate as a measure of the effectiveness of dispersal inducing stimuli in seagulls. Pp. 603-610 *in*: C. J. Amlaner, Jr., and D. W. Macdonald (Eds.), *A Handbook of Biotelemetry and Radio Tracking*. Pergamon Press, Oxford.
- TILLMAN, E. A., J. S. HUMPHREY, AND M. L. AVERY. 2002. Use of vulture carcasses and effigies to reduce vulture damage to property and agriculture. *Proc. Vertebr. Pest Conf.* 20:123-128.
- U.S. FISH AND WILDLIFE SERVICE. 2000. Waterfowl population status, 2000. U. S. Department of the Interior, Washington, D.C.
- WRIGHT, S. L. 1997. Canada geese: flying elephants we must avoid! *FAA Aviation News* 36:1-5.

