

1-1-2010

Dryland Performance of Sweet Sorghum and Grain Crops for Biofuel in Nebraska

Charles S. Wortmann

University of Nebraska - Lincoln, cwortmann2@unl.edu

Adam Liska

University of Nebraska - Lincoln, aliska2@unl.edu

Richard B. Ferguson

University of Nebraska - Lincoln, rferguson1@unl.edu

Drew J. Lyon

University of Nebraska--Lincoln, dlyon1@unl.edu

R. N. Klein

University of Nebraska - Lincoln, rklein1@unl.edu

See next page for additional authors

Follow this and additional works at: <http://digitalcommons.unl.edu/agronomyfacpub>

 Part of the [Plant Sciences Commons](#)

Wortmann, Charles S.; Liska, Adam; Ferguson, Richard B.; Lyon, Drew J.; Klein, R. N.; and Dweikat, Ismail M., "Dryland Performance of Sweet Sorghum and Grain Crops for Biofuel in Nebraska" (2010). *Agronomy & Horticulture -- Faculty Publications*. Paper 387.

<http://digitalcommons.unl.edu/agronomyfacpub/387>

This Article is brought to you for free and open access by the Agronomy and Horticulture Department at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Agronomy & Horticulture -- Faculty Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors

Charles S. Wortmann, Adam Liska, Richard B. Ferguson, Drew J. Lyon, R. N. Klein, and Ismail M. Dweikat



Dryland Performance of Sweet Sorghum and Grain Crops for Biofuel in Nebraska

C. S. Wortmann,* A. J. Liska, R. B. Ferguson, D. J. Lyon, R. N. Klein, and I. Dweikat

ABSTRACT

Sweet sorghum [SS; *Sorghum bicolor* (L.) Moench] is a potential biofuel crop for the Great Plains. Sweet sorghum was compared with corn [*Zea mays* (L.)] and grain sorghum for potential ethanol yield, energy use efficiency, and greenhouse gas (GHG) emissions at seven dryland site-years in Nebraska. Seasonal rainfall ranged from approximately 340 to 660 mm. Soils were deep with medium texture at all site-years. The effects of seeding rate, N rate, and cultivar on SS performance were evaluated. Sweet sorghum sugar yield was not affected by seeding rate and N application at six of seven site-years, but yield was increased by 19% at one site-year. Calculated ethanol yield and net energy yield were 33 and 21% more, respectively, with the grain crops compared with SS, but mean net energy yield of an earlier-maturing SS cultivar was comparable with the grain crops. The mean ratio of energy produced in ethanol per total energy invested was 23% less for grain crops compared with SS. Mean life cycle GHG emissions were 53% and 66 to 69% less compared with gasoline for SS and grain crops, respectively. Very efficient use of the ethanol coproducts was assumed for the grain crops while SS bagasse was assumed to be returned to the field. At least one SS cultivar is competitive with grain crops for some biofuel criteria, but SS is not competitive with grain crops for total or net liquid transportation fuel produced per hectare.

SWEET SORGHUM is a sugar crop with biofuel potential and has been found to be competitive with corn for theoretical ethanol yield with less energy invested (Smith et al., 1987; Smith and Buxton, 1993; Hunter and Anderson, 1997). Ethanol production from sugar does not require energy to depolymerize carbohydrates, as is required for grain starch. Smith et al. (1987) reported total sugar yield of SS ranging from 4 to 10.7 Mg ha⁻¹ for the continental United States and up to 12 Mg ha⁻¹ for Hawaii, while Smith and Buxton (1993) reported sugar yields at 6 Mg ha⁻¹ in Iowa and Colorado. Vermerris et al. (2007) reported total sugar concentrations of the juice ranging from 9 to 15%. Ricaud et al. (1979) found sugar concentration in juice at the grain soft dough stage (Warrick, 2009) to range from 12.8 to 16.6%. Calculated ethanol yield estimates for SS have generally ranged from 3000 to 4000 L ha⁻¹ (Lueschen et al., 1991), with one report estimating the ethanol yield potential to be as high as 8000 L ha⁻¹ (Hunter and Anderson, 1997). This amount is equivalent to ethanol produced from approximately 20 Mg ha⁻¹ of corn grain.

Biomass yield from SS has been found to be similar across a wide plant population range (Lueschen et al., 1991; Ferraris and Charles-Edwards, 1986). Sucrose concentration in the juice was 7% less but fermentable sugar yield was 14% more with a plant population of 140,000 compared with 70,000 plant ha⁻¹ (Broadhead and Freeman, 1980), and water soluble carbohydrate yield was greater with 160,000 compared with 80,000 plant ha⁻¹ (Martin and Kelleher, 1984).

Response to applied N has varied with location. Nitrogen rate did not affect fermentable sugar yield (Smith and Buxton, 1993), total and stalk dry matter yield at harvest (Barbanti et al., 2006), or fermentable carbohydrate and ethanol yield (Lueschen et al., 1991). In Louisiana, biomass yield has been shown to increase by 140% with application of 100 kg ha⁻¹ N, but yield was not increased with an additional 100 kg ha⁻¹ N (Ricaud and Arenneaux, 1990). In the same study, total sugar yield was increased by 150% by applying 100 kg ha⁻¹ N, with an increase of only 4% from an additional 100 kg ha⁻¹ N. Sweet sorghum has been found to require approximately 36% of the fertilizer N needed for similar ethanol yield levels in corn (Geng et al., 1989). Total dissolved solid concentration in stalk juice was shown to decrease with increased N rate (Wiendenfeld, 1984).

Life cycle GHG emissions are restricted for biofuels under the Energy Independence and Security Act of 2007 (available at http://frwebgate.access.gpo.gov/cgi-bin/getdoc.cgi?dbname=110_cong_bills&docid=f:h6enr.txt.pdf, verified 3 Dec. 2009). Advanced biofuels need to have >50% reduction in GHG emissions relative to gasoline. Ethanol from SS

279 Plant Sci., Univ. of Nebraska, Lincoln, NE 68583-0915. A contribution of the Univ. of Nebraska Agric. Res. Div. supported in part by funds provided through the Hatch Act. Additional support was provided by the U.S. Agency for International Development, under the terms of grant No. LAG-G-00-96-900009-00 to the International Sorghum and Millet Collaborative Research Support Program (INTSORMIL). Received 8 July 2009. *Corresponding author (cwortmann2@unl.edu).

Published in Agron. J. 102:319–326 (2010)
Published online 11 Dec. 2009
doi:10.2134/agronj2009.0271

Copyright © 2010 by the American Society of Agronomy, 677 South Segoe Road, Madison, WI 53711. All rights reserved. No part of this periodical may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording, or any information storage and retrieval system, without permission in writing from the publisher.



Abbreviations: BESS, Biofuel Energy Systems Simulator; CO₂e, carbon dioxide equivalent of greenhouse gases including carbon dioxide, methane, and nitrous oxide; GHG, greenhouse gas; HPAL, High Plains Agricultural Laboratory; NCTA, Nebraska College of Technical Agriculture; SCAL, South Central Agricultural Laboratory; SOC, soil organic carbon; SS, sweet sorghum; WCREC, West Central Research and Extension Center.

Table 1. Soil and agronomic information for sweet sorghum research at seven Nebraska site-years.

Site, soil, practice	HPAL07†	WCREC07	SCAL07	HPAL08	NCTA08	SCAL08	Havelock08
Location and elevation	41°13' N 103°0' W 1316 m	41°02' N 100°45' W 919 m	40°34' N 98°07' W 550 m	41°14' N 103°0' W 1292 m	40°39' N 100°31' W 824 m	40°34' N 98°08' W 552 m	40°51' N 96°37' W 357 m
Soil series and slope	Alliance silt loam, 0–1%	Hall silt loam, 0–1%	Crete silt loam, 0–1%	Keith loam, 0–1%	Hall silt loam, 0–1%	Hastings silt loam, 0–1%	Aksarben silty clay loam, 2–6%
Taxonomic class	fine-silty, mixed, mesic Aridic Argiustolls	fine-silty, mixed, superactive, mesic Pachic Argiustolls	fine, smectitic, superactive, mesic Pachic Argiustolls	fine-silty, mixed, mesic Aridic Argiustolls	fine-silty, mixed, mesic Pachic Argiustolls	fine, smectitic, superactive, mesic Udic Argiustolls	fine, smectitic, mesic Typic Argiudolls
Previous crop	wheat	wheat	wheat	wheat	wheat	soybean	soybean
Planting date							
Corn	12 May	16 May	22 May	15 May	9 May	2 May	13 May
Grain sorghum	6 June	6 June	29 May	31 May	3 June	1 June	13 May
Sweet sorghum	8 June	6 June	29 May	31 May	3 June	15 May	13 May
Harvest date‡	2 Oct.	1 Oct.	4 Sept.	21 Oct.	Oct.‡	29 Oct.‡	8 Oct.‡
Soil pH _{1:1}	6.55	5.45	5.66	5.61	6.04	7.03	4.62
SOM, kg kg ⁻¹ §	20.9	18.0	27.6	17.7	24.2	26.0	36.6
Bray-1 P, mg kg ⁻¹	17.2	71.3	32.6	7.7	18.3	11.5	13.4
Soil K, mg kg ⁻¹	3250	668	529	741	331	385	479
Soil NO ₃ -N, mg kg ⁻¹	5.98	7.59	3.68	7.28	14.56	6.49	3.33

† HPAL, High Plains Agricultural Laboratory; NCTA, Nebraska College of Technical Agriculture; SCAL, South Central Agricultural Laboratory; WCREC, West Central Research and Extension Center; 07 and 08, 2007 and 2008.

‡ Harvest at NCTA in 2008 spanned several days because of wet weather. Sweet sorghum 'Simon' was harvested 6 Sept., 27 Aug., and 25 Aug. at NCTA, SCAL, and Havelock, respectively, in 2008.

§ SOM, soil organic matter. Soil properties were from the 0- to 20-cm depth, except NO₃-N which was from the 0- to 120-cm depth.

would likely be in this category. Bioenergy yield and energy invested in producing the crop and converting it to ethanol are important factors in determining associated life cycle GHG emissions (Liska et al., 2009) as well as net energy yield (energy yield relative to energy invested). Low N rate requirements compared with grain crops and avoiding the need to convert starch to sugar favor SS relative to grain crops for more efficient ethanol production. To compare the ethanol-equivalent yield, CO₂-equivalent (CO₂e) emissions, and energy balance of crop-based ethanol production systems, the Biofuel Energy Systems Simulator (BESS model; Liska et al., 2009; available at www.bess.unl.edu; verified 3 Dec. 2009) provides a platform for analysis.

The objectives of this research were to compare the yield and efficiency of ethanol production from SS compared with corn or grain sorghum across a range of dryland environments and to develop N management and planting guidelines for SS production in Nebraska. These field results were used to estimate the direct life cycle GHG emissions associated with ethanol production from SS and the two grain crops (Liska et al., 2009).

MATERIALS AND METHODS

Research was conducted at seven site-years in Nebraska in 2007–2008 between 40°30' N to 41°15' N latitude and 350 to 1320 m elevation (Table 1). The soils were deep silt loam, loam, or silty clay loam. The previous crop was either soybean or winter wheat. All sites were no-till. A soil sample was taken for each block from the 0- to 20-cm depth before planting for basic analysis, and from the 0- to 120-cm depth for NO₃-N analysis. Fertilizer P was broadcast applied without incorporation as triple superphosphate per University of Nebraska-Lincoln recommendations (Ferguson, 2000). Growing season rainfall and

minimum and maximum temperature were recorded within 2 km of each site (Table 2).

Treatments, Experimental Design, and Procedures

The SS trials consisted of a randomized complete block design with a split-strip plot arrangement of treatments consisting of N rate, plant population, and cultivar. The N rates were 0, 45, and 90 kg ha⁻¹ surface applied before planting with urease inhibitor treated [N-(n-butyl) thiophosphoric triamide; Agrotain International, LCC, St. Louis, MO] urea. The plant population treatments were 75,000, 125,000, and 175,000 viable seed ha⁻¹. The N rate and plant population treatments were applied in eight-row main plots that were split in a nonrandom manner into four-row subplots for the two cultivars, resulting in a split-strip plot arrangement of treatments. The SS cultivars were Keller and M81E in 2007, and Simon and M81E in 2008; Keller lodged badly at two locations in 2007 and was replaced with Simon in 2008. These cultivars were selected based on performance in eastern Nebraska cultivar trials. Only Simon was planted at the Nebraska College of Technical Agriculture (NCTA) in 2008 because the M81E seed was lost.

The row spacing for the eight-row main plots was 76 cm. Plot length was 12 m. There were four replications for each treatment. To the outside of each SS replication, main plots of corn and grain sorghum were planted for comparison of SS with grain crops. The main plots of corn and grain sorghum were split to have either no N applied or N applied at the recommended rate, which ranged from 67 to 112 kg ha⁻¹ dependent on crop, yield goal, and predicted soil N availability (Ferguson, 2000). The main plots of corn and grain sorghum were eight rows wide with 76-cm spacing, 24 m long, and were split into 12 m subplots for N rates. The crop species were separated by at least four rows or 3 m of each crop to minimize border effects.

Table 2. Monthly rainfall and mean minimum and maximum temperature for seven Nebraska site-years in 2007 and 2008.

Site-year†	May			Jun			Jul			Aug			Sep		
	Rain	Min	Max	Rain	Min	Max	Rain	Min	Max	Rain	Min	Max	Rain	Min	Max
	mm	°C		mm	°C		mm	°C		mm	°C		mm	°C	
HPAL07	75	6.6	21.8	50	10.8	27.8	52	15.9	32.1	44	15.3	32.4	24	8.9	26.2
WCREC07	132	8.8	23.1	65	13.4	27.1	94	17.2	30.8	22	17.2	31.6	52	10.2	26.8
SCAL07	116	11.1	23.6	45	14.4	27.3	98	17.7	30.2	98	18.5	30.8	57	10.8	25.8
HPAL08	66	3.8	19.7	76	9.3	26.5	72	14.2	32.8	43	13.5	28.4	56	6.8	24.1
NCTA08	277	7.0	20.9	164	12.6	26.9	35	17.1	31.6	49	15.3	28.2	30	9.0	25.0
SCAL08	156	8.1	21.3	85	14.1	28.2	80	15.8	30.6	10	12.9	28.1	42	10.1	24.1
Havelock08	125	9.2	21.9	207	15.9	28.4	94	19.1	31.3	38	15.9	29.9	99	11.8	24.7

† HPAL, High Plains Agricultural Laboratory; NCTA, Nebraska College of Technical Agriculture; SCAL, South Central Agricultural Laboratory; WCREC, West Central Research and Extension Center; 07 and 08, 2007 and 2008.

Grain yield of grain sorghum at the Havelock Agronomy Farm was not determined because of severe bird damage.

Sweet sorghum harvest consisted of cutting the stalks at 8 to 10 cm above the soil surface in a single, 3-m length of row and counting harvested stalks. Harvest occurred at the soft dough stage for the earlier-maturing Simon, but was only near half-bloom or later stage and shortly before or after a killing frost for the later-maturity M81E and Keller (Table 1). Sweet sorghum biomass harvest was done in early September 2007 at South Central Agricultural Laboratory (SCAL07) when the crop was in the boot to half bloom growth stage because of severe lodging caused by strong winds, but Brix was measured later from standing plants at approximately the half bloom stage.

Fresh biomass weight was obtained at harvest. Leaves and panicles were removed from six stalks per treatment in one block. These six stalks were weighed with and without leaves and panicles to determine the mean conversion factor for converting all fresh biomass weights to fresh stalk weight. Brix is a measure of dissolved sugar to water mass ratio of a liquid. Brix was determined for six stalk segments per plot, with segments taken along the length of the stalks. Ten to 12 stalk segments of approximately 20-cm length, with leaves, were cut from stalks randomly selected from the biomass samples and at different positions to obtain samples representative of stalk length. These samples were weighed wet, oven-dried, and weighed dry to determine the water content after correcting for leaf weight. Juice and sugar yield were calculated in three steps:

$$CSY = (FSY - DSY) \times \text{Brix} \times 0.75;$$

$$JY, 80\% \text{ extracted} = [FSY - (DSY - CSY)] \times 0.8;$$

$$SY = JY \times \text{Brix} \times 0.75;$$

where CSY is conservative sugar yield (Mg ha^{-1}), FSY is fresh stalk yield (Mg ha^{-1}), DSY is dry stalk yield (Mg ha^{-1}), JY is juice yield (Mg ha^{-1}), and SY is sugar yield (Mg ha^{-1}). Sugar concentration of juice is 75% of Brix expressed in g kg^{-1} sugar juice.

Corn and grain sorghum yield were determined by either ear or panicle harvest, respectively, from two rows 6 m in length. Subsamples of six ears or panicles were weighed and the shelling or threshing percentage determined. Grain water content was corrected to 155 g kg^{-1} .

The ANOVAs for SS agronomic data by site-year were conducted using Statistix 9 (Analytical Software, Tallahassee, FL).

The ANOVA for the comparison of SS with corn and grain sorghum was combined across site-years for 2007, but was done by site-year for 2008 due to missing crop data at NCTA and Havelock. Site-year and replication effects were considered random and all treatment effects were considered fixed. Differences were considered significant at $P \leq 0.05$.

Estimation of Ethanol-Equivalent Yield, Greenhouse Gas Emissions, and Energy Balance

The BESS model (Liska et al., 2009) was used to calculate ethanol-equivalent yield, CO_2e emissions, and energy balance using state-of-the-art crop production and processing technology. The parameters and conversion efficiencies used in BESS are well documented in Liska et al. (2009) and most are not repeated in this paper. Emissions of N_2O were determined based on the 2006 IPCC Guidelines for National Greenhouse Gas Inventories from the Intergovernmental Panel on Climate Change (available at <http://www.ipcc-nggip.iges.or.jp/public/2006gl/index.html>; verified 3 Dec. 2009); BESS model equations were modified to account for N_2O emission for SS crop residues. Emission of N_2O from bagasse, stover, and roots was estimated as 1% of the N content, with N concentrations of 6 and 7 g kg^{-1} for above- and belowground biomass, respectively, and with root biomass equal to 25% of aboveground biomass. Ethanol yields from grain and fermentable sugar were based on industry survey data and theoretical yields, respectively (Table 3). Fermentable sugar yield of SS was estimated as 75% of the Brix value, with 80% of the sugar extracted from stalks and 95% of the extracted sugar converted to ethanol.

The BESS calculations were made assuming no-till production. Fuel use varied according to the crop and yield level (Table 3). Nitrogen rate used in the calculations varied according to recommendations for corn (Ferguson et al., 2000) and grain sorghum (Wortmann et al., 2006), while no N application was considered for SS production since SS yield was increased by N application in only one of seven site-years. Fertilizer P and K were applied at estimated removal rates for the grain crops, but calculations were done with no P and K application for SS, assuming that these nutrients would be returned to the fields in the bagasse. Application of 5 kg ha^{-1} of herbicide and no other pesticide was assumed for all crops. Wet distillers grains fed to beef cattle were given an energy and GHG coproduct credit that was attributed to corn and grain sorghum in the offset of other energy-intensive feeds. It was assumed that corn and grain sorghum stover stayed in the field and SS bagasse was returned to the field with no energy

Table 3. Parameter values used for estimation of ethanol yields, GHG emissions, and energy balances of corn, grain sorghum, and sweet sorghum in Nebraska.

Parameter†	Unit	Value
Crop production and harvest, diesel		
No-till production	L ha ⁻¹	4
Grain harvest, <8 (>8) Mg ha ⁻¹	L ha ⁻¹	9 (13)
SS harvest and crushing	L Mg ⁻¹ fresh stalks	0.3
Sugar-to-ethanol yield		
Brix to fermentable sugar concentration	%	75
Stalk juice expressed	%	80
Fermentable sugar converted to alcohol	%	95
Biorefinery yields and energy inputs		
Grain-to-ethanol conversion	L Mg ⁻¹	423
Sugar-to-ethanol conversion	L Mg ⁻¹	665
Grain ethanol natural gas input‡	MJ L ⁻¹ ethanol	5.44
SS juice fermentation and distillation	MJ L ⁻¹ ethanol	3.33

† All parameter values not shown above are reported in the BESS2008.3.1 User's Guide (www.bess.unl.edu; verified 2 Dec. 2009) developed from analysis of the corn-ethanol life cycle (Liska et al., 2009).

‡ Natural gas biorefinery in Nebraska producing wet distillers grains (BESS model default No. 5). The only cropping inputs for SS production besides fossil fuels were herbicide (5 kg ha⁻¹) and seed (2 kg ha⁻¹). Grain and SS juice transportation costs were assumed equivalent because of comparatively localized SS processing of higher masses. Biorefinery electricity use rate was assumed the same for all systems.

value given and no change in soil organic carbon (SOC). It was assumed that fuel consumed for transport of SS juice was similar to consumption for grain transport with shorter mean transport distance compensating for the greater SS mass transported. It was assumed that energy required for heating the juice for fermentation and distillation for SS was 3.3 MJ L⁻¹ of ethanol for mean outdoor air temperature < 10°C during these processes (Table 3).

RESULTS AND DISCUSSION

Overall measured or estimated SS means were 15.2 stalks m⁻² harvested, 109 g g⁻¹ fermentable sugar concentration in juice assuming fermentable sugar equal to 0.75 Brix, 51.4 Mg ha⁻¹ fresh stalk yield, 27.3 Mg ha⁻¹ juice yield assuming 80% extraction efficiency, 3.38 Mg ha⁻¹ fermentable sugar yield, and 2248 L ha⁻¹ ethanol yield assuming 95% of fermentable sugar is converted to ethanol (Table 4). Mean grain and calculated ethanol yield, respectively, were 7.94 Mg ha⁻¹ and 3361 L ha⁻¹ for corn, and 6.16 Mg ha⁻¹ and 2612 L ha⁻¹ for grain sorghum (Table 5).

Effects of Plant Population, Nitrogen Rate, and Cultivar

Variables of SS were not affected by two- and three-way interactions of N rate, plant population, and SS cultivars. Plant population effects on SS variables were not significant except for effects on stalk m⁻² at five site-years and juice, sugar, and ethanol yield at NCTA (Table 4). Stalk m⁻² increased by 32 to 43%, with increased plant population at High Plains Agricultural Laboratory (HPAL) and West Central Research and Extension Center (WCREC) in 2007 and at SCAL, NCTA, and Havelock in 2008. Yields of fresh stalks, juice, sugar, and ethanol were 14 to 19% greater at NCTA in 2008, with 17.5 seed m⁻² compared with the lower seed rates. Fresh stalk yield was decreased but Brix was increased with the high seeding rate compared with the lower rates at Havelock in 2008. The general lack of plant population effect on sugar yield is consistent with Lueschen et al. (1991) and Ferraris and Charles-Edwards

(1986), who reported no plant population effects on SS yield and sugar concentration.

Sweet sorghum was not affected by N rate at four of seven site-years (Table 4). Sugar concentration (Brix) increased by 5% at HPAL in 2007 but decreased by 5% at NCTA and by 7% at SCAL in 2008 with increased N rate. Fresh biomass, juice, sugar, and ethanol yield increased 19 to 27% with increasing N rate at NCTA in 2008, the site-years where sugar yield was greatest compared with other site-years. This occurred even though preplant soil nitrate N at this site-year (Table 1) was high compared with most site-years. Brix was 3% higher with 40 compared with 80 kg N ha⁻¹ applied at SCAL in 2008. Corn and grain sorghum yield were increased by means of 19 and 10%, respectively, with N application (data not reported). While substantial SS yield increase occurred with N application in Louisiana (Ricaud and Arenneaux, 1990), lack of SS response to applied N is common (Wiendenfeld, 1984; Lueschen et al., 1991; Barbanti et al., 2006). Based on other data collected by the authors of this paper (unpublished data, 2008), this lack of greater response of SS to N application is attributed to less N uptake, a more gradual rate of nutrient uptake, and N uptake later in the season compared with the grain crops.

An unmeasured observation was that Keller was the cultivar most susceptible to lodging with severe damage observed at SCAL and WCREC in 2007, while M81E was only moderately susceptible. Simon was earlier to mature and reached dough stage by early September, while most stalks of M81E and Keller were in preanthesis to milk growth stages by October. The performance of the SS cultivars Keller and Simon, relative to M81E, was inconsistent across site-years for sugar content and yield (Table 4). Because of later maturity, Keller and M81E were harvested before the soft dough stage of grainfill was reached, the stage when sugar concentration is expected to peak (Broadhead, 1973; Broadhead, 1974; Tarpley et al., 1994).

Ethanol-Equivalent Yield, Greenhouse Gas Emissions, and Energy Balance

Corn grain yield (Y_C) ranged from 4.6 to 11.8 Mg ha⁻¹ with a mean of 8.4 Mg ha⁻¹. Grain sorghum grain yield (Y_{GS}) ranged from 4.2 to 9.2 Mg ha⁻¹, with a mean of 6.2 Mg ha⁻¹ and was related to Y_C as $Y_{GS} = 0.67 Y_C + 0.70$ ($R^2 = 0.81$). Staggenborg et al. (2008) reported corn to have a grain yield advantage compared with grain sorghum under water-deficit conditions when corn grain yields exceed 6.4 Mg ha⁻¹. Estimated ethanol yield of SS varied across site-years independently of corn and grain sorghum grain yield (Fig. 1). Mean ethanol yield was 52% more with corn and 18% more with grain sorghum compared with SS (Table 5). Planting of all crops, especially grain sorghum and SS, was often later than intended due to weather conditions and competing priorities. The yields of the sorghum crops may have been higher with earlier planting, although the planting dates were not late compared with farmers' practice.

Calculated ethanol yield for SS was less than the yield for the two grain crops at all site-years, with the exception of NCTA in 2008. Ethanol yield with the three crops was less at HPAL, where in-season rainfall was <350 mm and elevation was higher, which causes a shorter growing season, compared with the other site-years. Ethanol yield for the three crops, however,

Table 4. Plant population, N rate, and cultivar effects on seven sweet sorghum properties at seven Nebraska site-years.

Site-year†	Stalks m ⁻²	Fresh stalk yield	Dry stalk yield	Juice yield, 80% expressed	Sugar content, 75% Brix	Sugar yield	Ethanol yield
			Mg ha ⁻¹		g kg ⁻¹	Mg ha ⁻¹	L ha ⁻¹
HPAL07							
Population	***	NS‡	NS	NS	NS	NS	NS
N rate	*	NS	NS	NS	*	NS	NS
Keller	13.4b‡	34.24b§	11.99	19.84a	114.4a	2.27	1509
M81E	14.8a	38.00a	12.51	22.37b	96.9b	2.17	1441
HPAL08							
Population	NS	NS	NS	NS	NS	NS	NS
N rate	NS	NS	NS	NS	NS	NS	NS
M81E	12.7	27.57	7.51b	17.39	83.6b	1.45a	967a
Simon	12.3	27.32	8.43a	17.14	133.9a	2.29b	1526b
WCREC07							
Population	***	NS	NS	NS	NS	NS	NS
N rate	NA	NS	NS	NS	NS	NS	NS
Keller	24.9a	59.45b	22.57	32.52a	102.3a	3.33	2212
M81E	17.8b	69.91a	23.36	40.36b	83.8b	3.38	2249
NCTA08, 'Simon'							
7.5 seed m ⁻²	14.1	92.98b	45.03	43.07a	122.3	5.27a	3503a
12.5 seed m ⁻²	16.3	91.55b	42.34	44.05a	119.0	5.24a	3486a
17.5 seed m ⁻²	18.9	104.78a	46.47	52.20b	119.0	6.21b	4131b
Population	**	*	NS	*	NS	*	*
N, 0 kg ha ⁻¹	15.5	86.32b	40.14	41.54a	124.3a	5.16a	3433a
N, 40 kg ha ⁻¹	17.1	96.58b	45.91	45.32a	118.0b	5.35a	3556a
N, 80 kg ha ⁻¹	16.8	106.42a	47.78	52.45b	118.1b	6.19b	4119b
SCAL07							
Population	NS	NS	NS	NS	NS	NS	NS
N rate	NS	NS	NS	NS	NS	NS	NS
Keller	10.6b	47.95	15.30	29.07	112.90	3.28	2182
M81E	12.8a	54.44	17.73	32.79	116.40	3.82	2538
SCAL08							
Population	**	NS	NS	NS	NS	NS	NS
N rate	NS	NS	NS	NS	*	NS	NS
M81E	12.7a	51.12	14.16a	33.33b	119.5b	5.31b	3530b
Simon	14.2b	50.95	18.08b	28.82a	96.6a	3.71a	2465a
Havelock08							
Population	***	**	NS	NS	*	NS	NS
N rate	NS	NS	NS	NS	NS	NS	NS
M81E	19.6	48.41b	20.80b	24.53a	110.60b	2.71a	1804a
Simon	19.1	75.61a	34.02a	37.43b	124.90a	4.67b	3109b

* Significant at the 0.05 probability level.

** Significant at the 0.01 probability level.

*** Significant at the 0.001 probability level.

† HPAL, High Plains Agricultural Laboratory; NCTA, Nebraska College of Technical Agriculture; SCAL, South Central Agricultural Laboratory; WCREC, West Central Research and Extension Center; 07 and 08, 2007 and 2008.

‡ NS refers to not significant at the 0.05 probability level.

§ Treatment means for site-year within columns with different letters are significantly different at the 0.05 probability level. Means separation is provided for cultivars but cultivars were planted in subplot strips and not randomly assigned to subplots.

was not related to rainfall amount for site-years where in-season rainfall ranged from 430 to 660 mm. Juice yield and Brix accounted for 83 and 15% of the variation in calculated ethanol yield of SS across site-years, respectively.

Calculated crop energy use was greatest for corn and least for SS (Table 5). Sweet sorghum had the highest estimates for transportation fuel consumption because of high harvest and juice extraction requirements (data not shown). Estimated total energy use (MJ L⁻¹) was 41% more with corn and grain sorghum compared with SS. Net energy yield was 49% more with corn and 18% more with grain sorghum compared with SS cultivars Keller and M81E, but 12% more with corn and 12% less with grain sorghum compared with SS cultivar Simon. Energy yield per energy invested was 9% more for the mean of the grain crops compared with SS.

Life cycle GHG emissions per liter of ethanol produced were 16 and 24% less for corn and grain sorghum, respectively, compared with SS, primarily due to higher N₂O emissions from crop biomass left to decompose on the field (Table 5) and the lack of a SS coproduct credit (Table 6). Approximately 15% of the CO₂e emission associated with SS production was estimated to consist of CO₂ emissions, while 85% of emissions were N₂O from residue decomposition. Estimated emission of CO₂e per unit of ethanol produced associated with crop production were 9% less with corn and 31% less with grain sorghum compared with SS (Table 5). While N was not applied to SS, 60% of the CO₂e emission associated with corn production was N₂O emission because of fertilizer N application and plant residue decomposition. Whereas life cycle GHG emissions for ethanol from grain crops averaged between 66 and 69% less

Table 5. Mean yields and calculated CO₂e emissions for grain and sugar produced (kg Mg⁻¹), ethanol produced (kg Mg⁻¹), and energy balances of corn, grain sorghum, and sweet sorghum ('M81E', 'Keller', and 'Simon') at seven Nebraska site-years. Grain crops included a standard energy and GHG coproduct credit, while no coproduct was included for sweet sorghum.

	2007, n = 3				2008, n = 4			
	Corn	Grain sorghum	Sweet sorghum		Corn	Grain sorghum	Sweet sorghum	
			M81E	Keller			M81E	Simon
Grain or sugar yield, Mg ha ⁻¹	7.36	5.73	3.12	2.96	8.38	6.62	3.16	4.06
N rate, kg ha ⁻¹	97	48	0	0	115	52	0	0
Ethanol yield, L ha ⁻¹	3113	2422	2077	1968	3547	2801	2099	2700
Crop† energy use MJ ha ⁻¹	8411	5594	3291	3282	9323	5939	3133	3466
Crop† CO ₂ emission, kg Mg ⁻¹	78.7	68.3	78.2	79.8	76.6	63.7	89.3	65.0
Crop† CH ₄ emission, kg Mg ⁻¹	0.08	0.07	0.06	0.06	0.08	0.07	0.07	0.05
Crop† N ₂ O emission, kg Mg ⁻¹	0.37	0.27	0.83	0.81	0.38	0.25	0.79	0.94
Crop† CO ₂ e‡ emission, kg Mg ⁻¹	191	149	328	323	192	141	327	345
Crop† CO ₂ e emission, g MJ ⁻¹	21.4	16.7	23.4	23.0	21.5	15.8	23.3	24.6
Life cycle CO ₂ e emission, g MJ ⁻¹	31.1	28.7	41.9	40.4	31.2	28.2	41.8	43.1
Energy use rate, MJ L ⁻¹	11.0	10.5	7.65	7.39	10.9	10.4	7.90	7.37
Energy yield, GJ ha ⁻¹	72.6	56.0	43.8	41.5	82.5	64.5	44.3	57.0
Energy consumed, GJ ha ⁻¹	34.1	25.5	15.7	14.6	38.5	29.0	15.7	19.6
Net energy yield, GJ ha ⁻¹	38.5	30.5	28.1	26.9	44.0	35.5	28.6	37.3
Net energy ratio	2.13	2.20	2.76	2.86	2.13	2.22	2.70	2.87
CO ₂ e reduction, %§	66.1	68.8	52.9	54.5	66.1	69.4	53.0	51.6

† Energy use and emissions associated with crop production. These and other estimates were calculated using the Biofuel Energy Systems Simulator (BESS; available at www.bess.unl.edu; confirmed 2 Dec. 2009).

‡ CO₂e, total global warming potential expressed as CO₂ equivalent.

§ CO₂e reduction associated with ethanol production and use compared with gasoline assuming 92 g CO₂e emission MJ⁻¹ for gasoline.

than gasoline, ethanol from SS was found to reduce emissions by 52 to 54% on average (Table 5, Fig. 2). The reduction in GHG emissions was more variable with SS compared with the

grain crops; the coefficients of variation were 7 to 14% for SS, but <3% for the grain crops.

These results show that SS will likely qualify as an *advanced* biofuel (per the Energy Independence and Security Act of 2007), with >50% reduction in GHG emission compared with gasoline, under the Energy Independence and Security Act of 2007. Life cycle GHG assessment requirements administered by the USEPA, however, have not yet been formalized. These regulations could impose additional indirect emissions in the life cycle due to land use change, which could systematically shift the GHG emissions results downward for most biofuels (Liska and Perrin, 2009). Performing better than SS, the




Fig. 1. Calculated ethanol yield of corn, grain sorghum, and sweet sorghum at three sites in 2007 and four sites in 2008 in Nebraska. The ANOVA was combined for sites in 2007 and the LSD 0.05 was 507 L ha⁻¹. The 2008 sites were analyzed separately because of sites with a missing crop or cultivar; the Y-bars indicate the LSD 0.05 for each site. HPAL, High Plains Agricultural Laboratory; NCTA, Nebraska College of Technical Agriculture; SCAL, South Central Agricultural Laboratory; WCREC, West Central Research and Extension Center.




Fig. 2. Average life cycle net energy yield and reduction of greenhouse gas emissions compared with gasoline of ethanol produced from sweet sorghum and grain crops in a study conducted at seven site-years in Nebraska. Error bars indicate the standard deviation based on data from seven site-years. Calculations did not account for potential N₂O emissions resulting from use of grain and grain coproducts, and assumed the most efficient grain biorefinery configuration.

dryland corn-ethanol and grain sorghum-ethanol systems with efficient use of wet distillers grains described in this study are considered near the upper end of the performance efficiency for grain-based biofuel systems (Liska et al., 2009). Based on other studies, GHG emissions from SS are comparable with the average for corn-ethanol systems, including less efficient systems with less efficient use of coproducts compared with the system used in this study, which reduce GHG emissions by roughly 50% compared with gasoline.

If crop residues from SS are removed from the field, the anticipated declines in SOC will result in increased net C emissions. With complete residue removal, loss of SOC is estimated to be approximately 0.8 Mg ha⁻¹ yr⁻¹ (Anderson-Teixeira et al., 2009) which would be an additional emission of 63 g CO₂e MJ⁻¹ of ethanol. This would increase net life cycle emissions to 18% greater compared with gasoline, based on the average scenario (Table 6). This suggests that SS bagasse should be returned to the field to maintain SOC, despite causing substantial N₂O emissions, unless either the feeding or a bioenergy use of bagasse results in sufficient life cycle reduction in CO₂e emissions to compensate for loss of SOC.

The crop N₂O emission estimates used in this study do not account for grain N that is eventually emitted as N₂O during ethanol production and handling of the coproducts and the amount that would occur from excrement of animals fed the coproducts. The coproducts, used as a substitute for corn and protein sources, often make up 20 to 40% of beef cattle rations. The resulting rations have more protein than needed for efficient cattle growth. This results in more manure N excretion than occurs with traditional rations and likely causes more N₂O emission during storage and handling, and following land application of manure compared with traditional beef cattle rations. There was no attempt, however, to estimate the N₂O emission from grain N in this study, resulting in estimates of crop N₂O emission that are likely low for the grain crops.

Estimated SS sugar yield and concentration in this study were low compared with concentrations reported by Vermeris et al. (2007) and Ricaud et al. (1979). Energy yield and energy balance would have been increased, and CO₂e emission reduced, for SS harvested at a higher sugar concentration. The low sugar concentration observed in this study for Keller and M81E is attributed to the many SS stalks that had not produced a panicle and to the large number of panicles that had not reached the dough stage by harvest time. The SS cultivar Simon did reach the dough stage, and sugar yields may have been higher if SS harvest dates had been selected based on monitoring for peak sugar concentration. The low SS sugar content resulted in relatively low ethanol yield in L ha⁻¹ and L Mg⁻¹ of biomass. Energy use efficiency for SS in this study was also low compared with potentially higher SS sugar contents per the reasons stated above. The low SS concentrations resulted in more fuel and energy consumption for biomass harvest and juice extraction, transport, fermentation, and distillation per liter of ethanol produced. Emissions of CO₂e L⁻¹ ethanol produced would decrease with increased sugar concentration.

When comparing SS to grain crops for ethanol production, the progress made in improving the efficiency of ethanol production from grain needs to be considered (Liska et al., 2009).

Table 6. Life cycle greenhouse gas emissions inventory of ethanol production from grain of corn and grain sorghum, and sweet sorghum ('M81E') sugar for the WCREC, 2007 site-years which had production levels near the seven Nebraska site-year average.

GHG emission category	Corn grain	Sorghum grain	Sweet sorghum
	-g CO ₂ e MJ ⁻¹ of ethanol produced†-		
Crop production			
Nitrogen fertilizer	4.12	2.27	0
Phosphorus fertilizer	1.0	1.0	0
Potassium fertilizer	0.596	0.604	0
Herbicides	1.56	2.16	2.68
Seed	0.40	0.097	0.034
Diesel	0.753	0.797	2.66
Depreciable capital	0.297	0.411	0.509
N ₂ O emissions	13.1	8.70	22.1
Total	21.8	16.0	26.0
Biorefinery			
Natural gas input	15.4	15.4	9.43
Electricity input	6.53	6.53	6.53
Depreciable capital	0.458	0.458	0.458
Grain transportation	2.11	2.11	2.11
Total	24.5	24.5	18.5
Coproduct credit			
Diesel	0.301	0.301	0
Urea production	-2.69	-2.69	0
Corn production	-10.3	-7.56	0
Enteric fermentable.-CH ₄	-3.69	-3.69	0
Total	-16.4	-13.6	0
Life-cycle net GHG emissions	31.3	28.3	45.9
GHG reduction relative to gasoline, %	65.9	69.2	50.1

† CO₂e, total global warming potential expressed as CO₂ equivalent.

Full credit was given in this study for the energy and GHG value of grain distillers coproducts. These coproducts are in much demand by beef cattle feeders because they are highly nutritious, result in improved growth rates, and provide feed use efficiency (Farran et al., 2006; Klopfenstein et al., 2008). Substituting distillers' coproducts for other feeds in beef rations is estimated to reduce CO₂e emissions by 28 to 30% in the corn-ethanol life cycles (Liska et al., 2009), assuming that feeding coproducts does not result in increased N₂O emission from excreted N. The grain-ethanol process benefits from the tremendous genetic and management gains that have occurred in Nebraska corn production. To a lesser extent, similar gains have occurred in grain sorghum production.

For the purposes of this study, an assumption was made that only 80% of the sugar was extracted from the stalks. More efficient extraction technology would increase ethanol yield and efficiency of SS. It was also assumed that fermentable sugars were 75% of the Brix reading; however, the concentration relative to Brix may be higher in many cases. Energy value as a fuel or animal feed might have been given to the SS bagasse, but it was assumed it would be land applied and was only credited for the return of the P and K although the C, N, and other nutrients that also would be returned to the land have energy value and contribute to reduced CO₂e emissions. Energy required for heating the juice for efficient fermentation and for distillation, which in this study was assumed to occur mostly in late fall, should be less with higher air temperatures and with the application of efficient heat exchange technology. Sweet sorghum productivity in Nebraska can probably be greatly improved through plant breeding; cultivars used in this

study are relatively old and, although selected for performance from a wider set of cultivars, not particularly well-adapted to Nebraska conditions.

CONCLUSION

Sweet sorghum sugar yield was improved by increasing sowing rate above 75,000 viable seed ha⁻¹ and with N application only at one of seven site-years, while corn and grain sorghum yields were increased by N application in most site-years. In consideration of unpublished data, the lack of more frequent response of SS to applied N is attributed to a lower rate of peak N uptake, proportionally more N uptake occurring later in the growing season, and to less total N uptake compared with corn and grain sorghum.

Mean calculated ethanol yield and net energy yield were less with SS than with the grain crops, but mean net energy yield of one of the SS cultivars evaluated, Simon, was comparable with the grain crops. The grain crops were much more efficient in terms of ethanol produced per petroleum invested as ethanol yields were higher and less fuel was used for grain crops compared with SS. Higher SS sugar concentrations would result in improved fuel use efficiency for harvest, crushing, and transport. Ethanol production with SS was more energy use efficient compared with the grain crops, with 25% more energy produced per MJ of energy invested.

The efficiency of SS might be improved by better use of bagasse that was assumed to be returned to the field in this study. The efficient use of the ethanol coproducts of corn and grain sorghum for feeding beef cattle as wet distillers grain, a common practice in Nebraska, contributed greatly to the efficiency of grain crops for ethanol production.

The later-maturing SS cultivars Keller and M81E both had >50% of the stalks that did not reach the dough stage at any of the seven site-years. This resulted in sugar concentrations that were lower than what might have occurred with a longer growing season. The cultivar Simon has a maturity time more appropriate for southern Nebraska and was observed to be less susceptible to lodging.

Sweet sorghum for ethanol production is considered competitive with grain crops for some criteria, and only with SS cultivar Simon in the environments tested. However, even this cultivar is not competitive with corn and grain sorghum for total or net liquid transportation fuel produced per hectare. Based on GHG emissions, SS may be competitive compared with average corn systems. Gains in SS yield, energy efficiency, and reduced GHG emissions caused by changed management practices are likely to be small. Cultivars with higher sugar yield and/or efficient use of the bagasse (while considering SOC impacts) will be needed to make SS competitive with corn and grain sorghum in ethanol production.

ACKNOWLEDGMENTS

The authors are grateful to J. Golus, R. Higgins, S. Melvin, J. Petersen, and G. Slater for their technical assistance.

REFERENCES

Anderson-Teixeira, K.J., S.C. Davis, M.D. Masters, and E.H. Delucia. 2009. Changes in soil organic carbon under biofuel crops. *Glob. Chan. Biol. Bioenergy* 1:75–96.

- Barbanti, L., S. Grandi, A. Vecchi, and G. Venturi. 2006. Sweet and fiber sorghum (*Sorghum bicolor* (L.) Moench), energy crops in the frame of environmental protection from excessive nitrogen loads. *Eur. J. Agron.* 25:30–39.
- Broadhead, D.M. 1973. Effects of deheading on stalk yield and juice quality of Rio sweet sorghum. *Crop Sci.* 13:395–397.
- Broadhead, D.M. 1974. Effect of planting date and maturity on juice quality of Rio sweet sorghum. *Agron. J.* 64:389–390.
- Broadhead, D.M., and K.C. Freeman. 1980. Stalk and sugar yield of sweet sorghum as affected by spacing. *Agron. J.* 72:523–524.
- Farran, T.B., G.E. Erickson, T.J. Klopfenstein, C.N. Macken, and R.U. Lindquist. 2006. Wet corn gluten feed and alfalfa hay levels in dry-rolled corn finishing diets: Effects on finishing performance and feedlot nitrogen mass balance. *J. Anim. Sci.* 84:1205–1214.
- Ferguson, R.B., G.W. Hergert, and E.J. Penas. 2000. Corn. In R.B. Ferguson (ed.) *Nutrient management for agronomic crops in Nebraska*. EC 01-155-S. Univ. of Nebraska Coop. Ext., Lincoln.
- Ferraris, R., and D.A. Charles-Edwards. 1986. A comparative analysis of the growth of sweet and forage sorghum crops. II. Accumulation of soluble carbohydrates and nitrogen. *Aust. J. Agric. Res.* 37:513–522.
- Geng, S., F.J. Hills, S.S. Johanson, and R.N. Sah. 1989. Potential yields and on-farm ethanol cost of corn, sweet sorghum, fodder beet and sugarbeet. *J. Agron. Crop Sci.* 162:21–29.
- Hunter, E.L., and I.C. Anderson. 1997. Sweet sorghum. *Hortic. Rev.* 21:73–104.
- Klopfenstein, T.J., G.E. Erickson, and V.R. Bremer. 2008. Board-invited review: Use of distillers byproducts in the beef cattle feeding industry. *J. Anim. Sci.* 86:1223–1231.
- Liska, A.J., and R.K. Perrin. 2009. Indirect land use emissions in the life cycle of biofuels: Regulations vs. science. *Biofuels Bioproducts Biorefining* 3:318–328.
- Liska, A.J., H.S. Yang, V.R. Bremer, T.J. Klopfenstein, D.T. Walters, G.E. Erickson, and K.G. Cassman. 2009. Improvements in life cycle energy efficiency and greenhouse gas emissions of corn-ethanol. *J. Ind. Ecol.* 13:58–74.
- Lueschen, W.E., D.H. Putnam, B.K. Kanne, and T.R. Hoversted. 1991. Agronomic practices for production of ethanol from sweet sorghum. *J. Prod. Agric.* 4:619–625.
- Martin, P.M., and F.M. Kelleher. 1984. Effects of row spacing and plant population on sweet sorghum yield. *Aust. J. Exp. Agric. Anim. Husb.* 24:386–390.
- Ricaud, R., and A. Arenneaux. 1990. Sweet sorghum for biomass and sugar production in 1990. Manuscript Rep. Exp. Stn., Louisiana State Univ., St. Gabriel.
- Ricaud, R., B. Cochran, A. Arenneaux, and G. Newton. 1979. Sweet sorghum for sugar and biomass production in Louisiana. Manuscript Rep. Exp. Stn., Louisiana State Univ., St. Gabriel.
- Smith, G.A., M.O. Bagby, R.T. Lewellan, D.L. Doney, P.H. Moore, F.J. Hills, L.G. Campbell, G.J. Hogaboam, G.E. Coe, and K. Freeman. 1987. Evaluation of sweet sorghum for fermentable sugar production potential. *Crop Sci.* 27:788–793.
- Smith, G.A., and D.R. Buxton. 1993. Temperate zone sweet sorghum ethanol production potential. *Bioresour. Technol.* 43:71–75.
- Staggenborg, S.A., K.C. Dhuyvetter, and W.B. Gordon. 2008. Grain sorghum and corn comparisons: Yield, economics, and environmental responses. *Agron. J.* 100:1600–1604.
- Tarpley, L., S.E. Lingle, D.M. Vietor, D.L. Andrews, and F.R. Miller. 1994. Enzymatic control of nonstructural carbohydrate concentrations in stems and panicles of sorghum. *Crop Sci.* 33:446–452.
- Vermerris, W., C. Rainbolt, D. Wright, and Y. Newman. 2007. Production of biofuel crops in Florida: Sweet sorghum. Available at <http://edis.ifas.ufl.edu/AG298>; [cited 6 Oct. 2009; verified 25 Nov. 2009]. Univ. of Florida.
- Warrick, B.E. 2009. How a sorghum plant develops. Available at <http://sanangelo.tamu.edu/agronomy/sorghum/sorghum.htm>; [cited 6 Oct. 2009; verified 25 Nov. 2009]. Texas Coop. Ext., Texas A&M Univ., San Angelo.
- Wiendenfeld, R. 1984. Nutrient requirement and use efficiency by sweet sorghum. *Energy Agr.* 3:49–59.
- Wortmann, C.S., A.R. Dobermann, R.B. Ferguson, G.W. Hergert, C.A. Shapiro, and D. Tarkalson. 2006. Fertilizer suggestions for grain sorghum. NebGuide G1669, Univ. of Nebraska, Lincoln.