

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Historical Materials from University of
Nebraska-Lincoln Extension

Extension

1994

NF94-158 Escherichia coli 0157:H7

Susan S. Sumner

Julie A. Albrecht

University of Nebraska--Lincoln, jalbrecht1@unl.edu

Follow this and additional works at: <https://digitalcommons.unl.edu/extensionhist>



Part of the [Agriculture Commons](#), and the [Curriculum and Instruction Commons](#)

Sumner, Susan S. and Albrecht, Julie A., "NF94-158 Escherichia coli 0157:H7" (1994). *Historical Materials from University of Nebraska-Lincoln Extension*. 495.

<https://digitalcommons.unl.edu/extensionhist/495>

This Article is brought to you for free and open access by the Extension at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Historical Materials from University of Nebraska-Lincoln Extension by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.



NebFact



Published by Cooperative Extension, Institute of Agriculture and Natural Resources,
University of Nebraska-Lincoln

Escherichia coli 0157:H7

By Susan S. Sumner, Extension Food Microbiologist
Julie A. Albrecht, Extension Food Specialist

- The Disease:** Symptoms of hemorrhagic colitis include severe abdominal cramps, followed by diarrhea (often bloody), nausea, vomiting, and occasionally a low-grade fever. A possible complication is hemolytic uremic syndrome (HUS), a urinary tract infection that is a leading cause of acute kidney failure in children. Symptoms generally begin three to four days after the food is eaten, last up to 10 days, and often require hospitalization.
- The Organism:** The specific serotype 0157:H7 produces a toxin that can cause hemorrhagic colitis. This bacteria grows slowly at refrigeration temperature. *E. coli* 0157:H7 has been shown to grow in acidic food products such as apple cider and mayonnaise.
- Sources:** Several strains of *Escherichia coli* bacteria, frequently associated with fecally contaminated water, have long been known to cause diarrhea in infants and travelers. Raw or rare ground beef, unpasteurized milk and unpasteurized apple cider have been involved in some illnesses.
- Control:** Cooking ground beef to 157°F, reheating, good sanitation, and refrigeration at 40°F or below are all important to control this microorganism.

*File NF157 under FOOD AND NUTRITION
F-13, Safety
Issued January 1994*

Issued in furtherance of Cooperative Extension work, Acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture. Elbert C. Dickey, Director of Cooperative Extension, University of Nebraska, Institute of Agriculture and Natural Resources.

University of Nebraska Cooperative Extension educational programs abide with the non-discrimination policies of the University of Nebraska-Lincoln and the United States Department of Agriculture.