

1998

Effect of Wet Corn Gluten Feed, Supplemental Protein, and Tallow on Steer Finishing Performance

C. J. Richards

University of Nebraska-Lincoln, Chris.Richards@okstate.edu

Rick Stock

University of Nebraska-Lincoln, rstock3@Unl.edu

Terry J. Klopfenstein

University of Nebraska-Lincoln, tklopfenstein1@unl.edu

D. H. Shain

University of Nebraska-Lincoln

Follow this and additional works at: <http://digitalcommons.unl.edu/animalscifacpub>

 Part of the [Animal Sciences Commons](#)

Richards, C. J.; Stock, Rick; Klopfenstein, Terry J.; and Shain, D. H., "Effect of Wet Corn Gluten Feed, Supplemental Protein, and Tallow on Steer Finishing Performance" (1998). *Faculty Papers and Publications in Animal Science*. 520.
<http://digitalcommons.unl.edu/animalscifacpub/520>

This Article is brought to you for free and open access by the Animal Science Department at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Papers and Publications in Animal Science by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Effect of Wet Corn Gluten Feed, Supplemental Protein, and Tallow on Steer Finishing Performance¹

C. J. Richards, R. A. Stock², T. J. Klopfenstein³, and D. H. Shain

Department of Animal Science, University of Nebraska, Lincoln 68583-0908

ABSTRACT: Two trials evaluated the effects of level of wet corn gluten feed (WCGF), type of supplemental protein, and supplemental tallow on steer finishing performance. In Trial 1, WCGF was fed at 0, 25 (two diets), or 50% of the dietary DM replacing dry-rolled corn (DRC), molasses, and a portion of the supplement. The DRC control diet and one 25% WCGF diet were supplemented with a combination of protein sources. The second 25% WCGF diet was supplemented with urea alone. The 50% WCGF diet contained no additional protein supplementation. No differences in DMI ($P > .10$) were observed. Calves fed 25% WCGF plus a combina-

tion of protein sources or 50% WCGF gained faster ($P < .10$) and more efficiently ($P < .10$) than calves fed the DRC control. Calves fed 25% WCGF plus urea gained faster ($P < .10$) and tended ($P = .14$) to be more efficient than calves fed the DRC control. In Trial 2, WCGF was fed to replace 0 or 50% of the DRC and the molasses-urea supplement (DM basis). Both diets were fed with or without 3% tallow. Steers fed WCGF gained faster ($P < .01$) and more efficiently ($P < .01$) than steers fed DRC. Inclusion of 3% tallow increased gain ($P < .05$) and improved efficiency ($P < .05$). Feed efficiency is improved by the addition of WCGF or tallow to DRC finishing diets.

Key Words: Protein, Gluten Feed, Tallow, Maize, Cattle

©1998 American Society of Animal Science. All rights reserved.

J. Anim. Sci. 1998. 76:421-428

Introduction

Wet corn gluten feed (**WCGF**) contains 90 to 100% of the relative energy value of dry-rolled corn (**DRC**) in finishing diets (Ham et al., 1995). Less starch and more highly digestible fiber is associated with WCGF than with corn grain (DeHaan, 1983) and may reduce problems associated with acidosis (Firkins et al., 1985). Ruminal escape protein concentration is lower in WCGF than in corn (McCoy, 1996), and this may create a deficiency in metabolizable protein, especially for young, rapidly gaining calves fed finishing diets.

Wet distillers grains improved feed efficiency of cattle fed dry-rolled corn finishing diets from 2 to 17% when fed at levels of 5.2 to 40% of the dietary DM (Larson et al., 1993). Wet corn distillers grains contain 30% CP (47% escape protein; Firkins et al., 1984) and approximately twice the lipid content found in corn or WCGF. Additional lipid content may explain a portion of the higher energy value (Larson et al.,

1993) of wet distillers by-products compared with the energy value of WCGF (Ham et al., 1995).

The objectives of these trials were to determine the effects of level of WCGF in the diet, the interaction of level of WCGF and type of supplemental protein (urea or a combination of protein sources), and the interaction of level of WCGF and supplemental tallow on steer finishing performance.

Materials and Methods

Trial 1. British crossbred steer calves ($n = 160$; 8 to 9 mo of age; mean BW 276 ± 19 kg) were blocked by weight and allotted randomly within block to one of four treatments (four pens per treatment). Wet corn gluten feed (Cargill Corn Milling, Eddyville, IA) was fed at 0, 25 (two treatments), or 50% of the dietary DM replacing dry-rolled corn (DRC) and molasses. The DRC control diet and one 25% WCGF diet were supplemented with a combination of urea (50% of supplemental protein), soybean meal (25% of supplemental protein), and an 80:20 combination (DM basis) of feather meal and blood meal (25% of supplemental protein). This combination of protein sources was formulated to supply N, amino acids, and peptides for microbial synthesis and escape protein in an attempt to meet metabolizable protein (**MP**)

¹Published with the approval of the director as paper no. 11833, journal ser., Nebraska Agric. Res. Div.

²Present address: Cargill, Inc., Blair, NE.

³To whom correspondence should be addressed.

Received February 26, 1997.

Accepted September 21, 1997.

needs. A second 25% WCGF treatment was supplemented with urea (100% of supplemental protein). The 50% WCGF diet received no additional supplemental protein. Final finishing diets (Table 1) were formulated (DM basis) to contain a minimum of 13% CP, .70% Ca, .30% P, .65% K, and included 28 mg/kg monensin (Elanco Animal Health, Indianapolis, IN), 11 mg/kg tylosin (Elanco Animal Health), and 53 mg/kg of thiamine (WCGF diets only). Nitrogen analysis of feed ingredients was performed using a combustion-type N analyzer (Perkin Elmer model PE 2410 Series II, Norwalk, CT). Actual dietary CP values (Table 1) were lower than calculated at the initiation of the trial due to lower than expected CP contents of DRC and WCGF.

Steers were implanted with Revalor (Hoeschst-Roussel Agri-Vet Company, Somerville, NJ) initially and on d 92. Initial weights were the average of two consecutive days' weights taken before feeding. To minimize gut fill differences, all steers were fed a common diet (45% alfalfa hay, 26% WCGF, 23% DRC,

3% molasses, and 3% supplement, DM basis) at an estimated 2% of their body weight (DM basis). All steers were adapted to their final diet in 21 d by feeding 45, 35, 25, and 15% forage diets for 4, 3, 7, and 7 d, respectively, assuming alfalfa hay contained 100% forage and corn silage contained 50% forage. In the WCGF treatments, WCGF was added to the maximum treatment level (25 or 50%, DM basis) before DRC increases began. The trial began November 30, 1993, and steers were fed for an average of 160 d in outside, dirt-mounded pens. At slaughter, hot carcass weights and liver scores were recorded. Liver scores were recorded in accordance with the system of Elanco Products Company (1974). Final weights were calculated from hot carcass weights assuming a constant 62% dressing percentage. After a 48-h chill, 12th rib fat thickness, quality grade, and yield grade were determined.

Net energy for gain for each diet was calculated according to procedures of Larson et al. (1993) with the exception that net energy required for daily gain

Table 1. Composition of finishing diets fed to calves in Trial 1

Item ^b	Treatment ^a			
	DRC	25% WCGF + urea	25% WCGF + protein	50% WCGF
Dry rolled corn	78.30	61.00	61.00	36.00
Molasses ^c	6.10	—	—	—
WCGF	—	25.00	25.00	50.00
Corn silage	5.00	5.00	5.00	5.00
Alfalfa hay	5.00	5.00	5.00	5.00
Dry supplement				
Finely ground corn	.09	1.89	1.54	1.83
Soybean meal	1.86	—	.28	—
Blood meal	.20	—	.03	—
Feather meal	.80	—	.12	—
Urea	.63	.18	.10	—
Limestone	1.46	1.47	1.47	1.71
Sodium chloride	.21	.30	.30	.30
Potassium chloride	.17	—	—	—
Tallow	.11	.08	.08	.08
Vitamin premix ^d	.01	.01	.01	.01
Trace mineral premix ^e	.03	.03	.03	.03
Rumensin premix ^f	.02	.02	.02	.02
Tylan premix ^g	.01	.01	.01	.01
Thiamin premix ^h	—	.01	.01	.01
Nutrient composition ⁱ , %				
Crude protein	11.62	11.25	11.27	13.22
Calcium	.70	.70	.70	.83
Phosphorus	.32	.49	.50	.69
Potassium	.65	.65	.66	.89

^aDRC = dry-rolled corn; WCGF = wet corn gluten feed; protein (% of supplemental protein) = 50% urea, 25% soybean meal, and 25% of an 80:20 combination of feather meal and blood meal.

^b% DM basis.

^cMolasses contained 1.5% NaCl.

^d15,000 IU of vitamin A, 3,000 IU of vitamin D, and 3.75 IU of vitamin E/g of premix.

^e10% Mg, 6% Zn, 4.5% Fe, 2% Mn, .5% Cu, .3% I, and .05% Co.

^f132 g of monensin/kg of premix.

^g88 g of tylosin/kg of premix.

^h88 g of thiamin/kg of premix.

ⁱCP based on measured values, other nutrients based on tabular values (NRC, 1984).

Table 2. Feed ingredient nutrient values^a used in NRC (1966) model calculations

Item	Trial 1		Trial 2		DIP ^c	Pretrial TDN ^d	eNDF ^{de}	NDF ^d
	Pretrial CP	Final CP ^b	Pretrial CP	Final CP ^b				
Dry-rolled corn	10.0	8.3	8.4	8.1	40	88	0	9.0
Wet corn gluten feed	20.0	18.3	18.3	19.9	80	88	0	36.2
Molasses	6.0	6.0	—	—	100	60	0	0
Molasses-urea supplement	—	—	50.6	50.6	100	60	0	0
Corn silage	8.0	7.3	8.0	5.0	75	75	71	41.0
Alfalfa hay	15.9	14.2	15.0	5.0	80	52	92	53.0
Tallow	—	—	0.0	0.0	0	0	0	0
Urea	291.0	291.0	291.0	291.0	100	0	0	0
Soybean meal	49.0	49.0	—	—	70	84	23	14.9
Blood meal	90.0	90.0	—	—	20	66	0	.9
Feather meal	91.3	91.3	—	—	40	55	23	39.0

^aPercentage (DM basis).

^bAverage crude protein content of dry-rolled corn, wet corn gluten feed, corn silage, and alfalfa hay samples determined in our laboratory.

^cDIP = degradable intake protein, assumed values for calculations.

^dValues listed in NRC (1996) model except for dry-rolled corn, which was changed to 0% eNDF and 9% NDF.

^eEffective NDF, percentage of NDF.

(**NE_gR**) was calculated using the equation $NE_{gR} = .0493 W^{.75} (ADG^{1.097})$, where NE_{gR} (Mcal/kg gain) is the net energy required for daily weight gain (ADG, kg/d) and W (kg) is the mean body weight (NRC, 1984). All other calculations were the same as reported by Larson et al. (1993).

Degradable intake protein (**DIP**) and MP supplies and requirements were calculated using the NRC (1996) model and values in Table 2. The animal evaluated was a 422-kg, 12-mo-old Simmental-Angus-Hereford growing/finishing steer that had a body condition score of 5 and a mature weight of 568 kg. Management settings included in the model were the use of an implant and ionophore. Environmental settings included the following values: 8 km/h wind speed, previous and current temperatures of 0°C, no night cooling, average hide thickness, some mud on the lower body, and no heat stress.

Assuming a DMI of 9.55 kg/d and an ADG of 1.64 kg/d before initiation of the trial, supplies of DIP and MP were calculated to exceed the animal's requirement (Table 3). Estimates of TDN and effective NDF (**eNDF**) on a DM basis and ruminal pH were also estimated.

After completion of the trial, degradable intake protein and MP supplies and requirements were estimated for each pen using the NRC (1996) model. Analyzed CP values for alfalfa, corn silage, corn, and WCGF were used for calculations. Actual pen DMI were used to calculate intakes of individual ingredients. Amounts of individual ingredients fed were entered into the NRC (1996) model. The NE_m and NE_g adjusters were changed equally until the proper ADG for the pen was achieved. Next, TDN percentages of the diets were calculated from NE_g using

Table 3. Pretrial protein, effective NDF, and pH estimates using the NRC (1996) model for Trial 1

Item ^b	Treatment ^a			
	DRC + protein	25% WCGF + urea	25% WCCF + protein	50% WCGF
DIP ^c intake, kg/d	.69	.78	.75	1.01
DIP requirement, kg/d	.65	.65	.65	.65
MP ^d supplied, kg/d	.86	.79	.80	.76
MP requirement kg/d	.77	.77	.77	.77
TDN, %	81	82	82	82
eNDF ^e , %	4.1	3.9	3.9	4.0
pH	5.60	5.60	5.60	5.60

^aDRC = dry-rolled corn; WCGF = wet corn gluten feed.

^bDM basis; 9.55 kg/d DMI; 1.64 kg ADG.

^cDIP = degradable intake protein.

^dMP = metabolizable protein.

^eeNDF = effective NDF.

formulas from NRC (1984). In the model, a dietary ingredient's TDN percentage was altered until the diet balanced for the TDN calculated from NE_g. Calculated TDN values were 82% for the DRC treatment and 84% (DM basis) for the remaining treatments. After obtaining the proper dietary TDN, ruminal pH was estimated. For diets containing WCGF, the eNDF value of WCGF in the model was adjusted to result in a ruminal pH increase of .09 and .18 units for the 25 and 50% WCGF diets, respectively. The adjustment of eNDF increasing ruminal pH in the model represented the average increase in ruminal pH observed by Ham et al. (1994) and Richards (1996) when feeding similar diets.

Statistical analysis was performed in accordance with procedures of Steel and Torrie (1980) for a randomized complete block design. Means were separated with Duncan's Multiple Range Test ($\alpha = .10$) using the GLM procedure of SAS (1989). Pen was the experimental unit and the model effects included treatment and block.

Trial 2. Yearling steers ($n = 160$; mean BW = 346 \pm 45) were limit-fed 6.8 kg (DM basis) of a 50% corn

silage and 50% alfalfa hay diet (DM basis) for 5 d. Initial weights were taken on two consecutive days before feeding. Steers were blocked by weight (four pens per treatment) and allotted randomly within block to one of four treatments (10 steers/pen, four pens/treatment) in a 2 \times 2 factorial arrangement of treatments. Wet corn gluten feed was fed to replace 0 or 50% of the DRC and molasses-urea supplement (DM basis). All diets were fed with or without the addition of 3% tallow (DM basis).

Final finishing diets (Table 4) were formulated (DM basis) to contain a minimum of 13% CP, .70% Ca, .30% P, .65% K, and included 33 mg/kg monensin and 11 mg/kg tylosin. Diets containing WCGF had 53 mg/kg thiamine added. Actual CP values were determined for feedstuffs as in Trial 1. The CP concentration of DRC was lower during the trial than formulated and resulted in lower CP concentrations than calculated in the DRC diets. Lipid content of WCGF and DRC were measured using the chloroform-methanol extraction procedures of Moore et al. (1986).

Table 4. Composition of finishing diets fed to steers in Trial 2

Item ^b	Treatment ^a			
	DRC	DRC + tallow	50% DRC: 50% WCGF	50% DRC: 50% WCGF + tallow
Dry-rolled corn	82.00	78.90	44.01	42.43
WCGF	—	—	44.00	42.43
Molasses-urea supplement ^c	5.00	5.00	—	—
Corn silage	5.00	5.00	5.00	5.00
Alfalfa hay	5.00	5.00	5.00	5.00
Tallow	—	3.00	—	3.00
Dry supplement				
Urea	.88	.97	.06	.20
Limestone	1.45	1.44	1.49	1.50
Sodium chloride	.30	.30	.30	.30
Potassium chloride	.25	.27	—	—
Tallow	.09	.09	.06	.06
Vitamin premix ^d	—	—	.01	.01
Trace mineral premix ^e	—	—	.03	.03
Rumensin premix ^f	.02	.02	.02	.02
Tylan premix ^g	.01	.01	.01	.01
Thiamin premix ^h	—	—	.01	.01
Nutrient composition ⁱ , %				
Crude protein	12.71	12.72	15.94	15.51
Calcium	.70	.70	.70	.70
Phosphorus	.30	.30	.52	.51
Potassium	.65	.65	.94	.91

^aDRC = dry-rolled corn; WCGF = wet corn gluten feed.

^b% DM basis.

^cSupplement was based on a molasses-urea liquor and contained (DM basis) 50.6% CP, 2.61% K, .95% Ca, .55% P, .41% S, .35% Mg, 78,830 IU of vitamin A/kg, 15,770 IU of vitamin D/kg, and 20 IU of vitamin E/kg.

^d15,000 IU of vitamin A, 3,000 IU of vitamin D, and 3.75 IU of vitamin E/g of premix.

^e10% Mg, 6% Zn, 4.5% Fe, 2% Mn, .5% Cu, .3% I, and .05% Co.

^f176 g of monensin/kg of premix.

^g88 g of tylosin/kg of premix.

^h88 g of thiamin/kg of premix.

ⁱCP based on measured values, other nutrients based on tabular values (NRC, 1984).

Table 5. Pretrial protein, effective NDF, and pH estimates using the NRC (1996) model for Trial 2

Item ^b	Treatment ^a			
	DRC	DRC + tallow	50% WCGF	50% WCGF + tallow
DIP ^c intake, kg/d	.95	.97	.98	.99
DIP requirement, kg/d	.75	.72	.77	.74
MP ^d supplied, kg/d	.86	.83	.85	.82
MP requirement kg/d	.77	.77	.77	.77
TDN, %	82	79	84	81
eNDF ^e , %	3.9	3.9	3.9	3.9
pH	5.59	5.59	5.59	5.59

^aDRC = dry rolled corn; WCGF = wet corn gluten feed.

^bDM basis; 10.91 kg/d DMI; 1.64 kg/d ADG.

^cDIP = degradable intake protein.

^dMP = metabolizable protein.

^eeNDF = effective NDF.

All steers were adapted to final diets in 21 d by feeding 45, 35, 25, and 15% forage diets for 2, 5, 7, and 7 d, respectively. Concentrate increases in the 50%DRC:50% WCGF treatment during the adaptation period were increased in equal portions on a DM basis. Cattle were implanted with Revalor on d 1 and the lightest replication received a second implant on d 78. The trial began December 21, 1994, and steer replicates were fed for 93, 107, 135, and 149 d.

Collection of hot carcass weights, liver scores, 12th rib fat thickness, quality grade, yield grade, and calculation of final weights were performed as in Trial 1. Net energy for gain for each diet was calculated as described in Trial 1 except NE_gR was calculated as $NE_gR = .0557 W^{.75} (ADG^{1.097})$.

Assuming a DMI of 10.91 kg/d and an ADG of 1.64 kg/d before initiation of the trial, protein supplies and

requirements were calculated as in Trial 1. These calculations estimated the supply of DIP and MP to exceed requirements (Table 5). Estimates of TDN and eNDF on a DM basis and ruminal pH were also made.

After completion of the trial, protein supplies and requirements were calculated using the NRC (1996) model as in the previous trial with protein levels of feedstuffs, dietary TDN, average age of the animal (15 mo), and average weight of the animal (445 kg) changed. Dietary TDN values calculated from NE_g were 77, 80, 84, and 88% (DM basis) for the DRC, DRC + tallow, 50% WCGF, and 50% WCGF + tallow, respectively.

Analysis of variance was according to procedures of Steel and Torrie (1980) for a randomized complete block with a 2 × 2 factorial arrangement of treatments. The GLM procedure of SAS (1989) was used

Table 6. Effect of wet corn gluten feed and supplemental protein on calf finishing performance in Trial 1

Item	Treatment ^a				SEM
	DRC + protein	25% WCGF + urea	25% WCCF + protein	50% WCGF	
DMI, kg/d	9.60	9.90	10.09	9.81	.19
Daily gain, kg	1.71 ^b	1.82 ^c	1.88 ^c	1.84 ^c	.03
Gain/feed	.179 ^b	.184 ^{bc}	.186 ^c	.187 ^c	.002
DIP intake ^d , kg/d	.64 ^a	.70 ^b	.70 ^b	.92 ^c	.16
DIP requirement, kg/d	.66 ^a	.77 ^b	.77 ^b	.79 ^b	.14
MP supplied ^e , kg/d	.83	.82	.86	.82	.15
Fat thickness, cm	.89 ^b	.97 ^b	1.09 ^c	1.09 ^c	.05
Quality grade ^f	18.6	18.3	18.4	18.3	.2
NE_g diet, Mcal/kg ^g	1.34 ^a	1.39 ^{ab}	1.41 ^b	1.41 ^b	.02

^aDRC = dry-rolled corn; WCGF = wet corn gluten feed; protein (% of supplemental protein) = 50% urea, 25% soybean meal, and 25% of a 80:20 combination of feather meal and blood meal.

^{b,c}Means within a row with unlike superscripts differ ($P < .10$).

^dDIP = degradable intake protein.

^eMP = metabolizable protein; requirement (NRC, 1996) for average WCGF daily gain (1.85 kg/d) = .83 kg/d.

^f18 = high Select, 19 = low Choice.

^gCalculated from animal performance.

Table 7. Effect of wet corn gluten feed and tallow on finishing steer performance in Trial 2

Item	Treatment ^a				SEM
	DRC	DRC + tallow	50% DRC: 50% WCGF	50% DRC: 50% WCGF + tallow	
DMI, kg/d	11.17	10.98	10.86	10.91	.12
Daily gain ^{bc} , kg	1.56	1.64	1.71	1.80	.03
Gain/feed ^{bc}	.142	.151	.159	.166	.008
DIP intake ^{bd} , kg/d	.96	.96	1.03	1.04	.10
DIP required ^{be} , kg/d	.72	.75	.89	.93	.10
MP supplied ^{bf} , kg/d	.85	.85	.94	.96	.09
Fat thickness, cm	.97	.94	.91	1.02	.05
Quality grade ^g	18.6	18.4	18.4	18.8	.3
NE _g diet, Mcal/kg ^{beh}	1.23	1.30	1.39	1.44	.02

^aDRC = dry-rolled corn; WCGF = wet corn gluten feed.

^bDRC vs 50% DRC:50% WCGF ($P < .01$).

^c0 vs 3% tallow ($P < .05$).

^dDIP = Degradable intake protein.

^e0 vs 3% tallow ($P < .01$).

^fMP = metabolizable protein supplied was calculated excluding energy from tallow; requirement (NRC, 1996) of 50% DRC:50% WCGF + tallow daily gain (1.80 kg/d) = .81 kg/d.

^g18.0 = high Select, 19.0 = low Choice.

^hCalculated from animal performance.

with model effects that included treatment and block. Preplanned contrasts of DRC vs 50% WCGF, 0 vs 3% tallow, and WCGF level \times tallow level were used to separate treatment means.

Results and Discussion

Trial 1. Steers fed 25 or 50% WCGF gained faster ($P < .10$; Table 6) than calves fed the DRC control, but no differences were observed ($P > .10$) in DMI. Calves fed the 25% WCGF diet plus a combination of protein sources or 50% WCGF diet were more efficient ($P < .10$) than calves fed the DRC control. Calves fed 25% WCGF plus urea tended ($P = .14$) to be more efficient than calves fed the DRC control. An increase in efficiency and gain may be attributed to a reduction of subacute acidosis (Stock et al., 1990). A reduction in subacute acidosis has been suggested by Farlin (1981) and Firkins et al. (1985) when corn by-products were fed. The faster and more efficient gains of the calves on the WCGF diets resulted in increased ($P < .10$) fat over the 12th rib, but quality grade, yield grade (data not shown), and liver scores (data not shown) were not affected.

Net energy for gain calculated from animal performance and NRC (1984) values for the DRC control diet were 1.34 vs 1.38 Mcal/kg, respectively. Assuming a NE_g of 1.55 Mcal/kg for DRC, energy values for WCGF were calculated to be 1.59, 1.66, and 1.62 Mcal/kg for the 25% plus urea, 25% plus protein, and 50% WCGF treatments, respectively. The average WCGF energy value calculated from this trial (1.62 Mcal/kg) is higher than the values of 1.40 and 1.55 Mcal/kg

found for WCGF diets in two finishing trials by Ham et al. (1995). In the two trials conducted by Ham et al. (1995), British crossbred yearling cattle were fed WCGF at 35 and 70% or 17.5, 35, 52.5, 70, and 87.5% of dietary DM. Two trials determining energy values for WCGF were conducted by Trenkle (1986a,b). When feeding heifers finishing diets containing 30 or 60% WCGF, the metabolizable energy value for WCGF was 22% lower than for DRC. In the second trial, an energy value of 9% less than DRC was calculated from steers fed finishing diets with 30, 50, and 70% of the DRC and corn silage replaced by WCGF.

Degradable intake protein was lowest for the DRC control, intermediate for 25% WCGF treatment, and highest for the 50% WCGF treatment ($P < .10$). The DIP intakes of cattle consuming the DRC and 25% WCGF diets were estimated to be deficient. However, the additional N needs of these treatments may have been met by recycled urea in the animal. The model uses a constant rate of N recycling. Ruminants may have a greater ability to recycle N when it is limiting than is allowed for by the NRC (1996) model. When calculating DIP requirements and adjusting pH level using effective NDF, the DIP required will be altered and a requirement should be calculated for each diet. Strobel and Russell (1986) conducted in situ experiments that showed that mixed ruminal bacteria produced 50% less protein at pH 5.7 than at pH 6.7. Therefore, when effective NDF is less than 20% of the dietary DM, the microbial yield predicted is reduced 2.5% for every 1% decrease in effective NDF (Russell et al., 1992). If DIP becomes limiting, microbial CP yield should decline. Metabolizable protein requirement is driven by ADG for a given animal. All animals

in a trial are assumed to be capable of equal rates of gain. Therefore, the requirement for MP should be based on the highest-gaining treatment or group of treatments. Because the WCGF treatment ADG were not different ($P > .10$), the average ADG from the WCGF treatments (1.85 kg) was used for determining the MP requirement for this trial. Metabolizable protein supplied was not different ($P < .10$) for any of the treatments, but the MP supplied was estimated to be slightly below the requirement for the 25% WCGF + urea and 50% WCGF treatments.

In a finishing trial, Krehbiel et al. (1995) fed varying levels of WCGF to yearling Hereford steers with or without addition of escape protein. The addition of escape protein did not result in an increase in performance. In contrast, Trenkle (1992) reported an increased MP requirement when feeding DRC-based diets to large-framed yearling steers implanted with Synovex-S and Finaplix-S.

Trial 2. Steers fed 50% WCGF diets gained faster ($P < .01$) and more efficiently ($P < .01$) than steers fed DRC diets; DMI did not differ among treatments ($P > .10$; Table 7). Daily gains of steers were higher ($P < .05$) and more efficient ($P < .05$) when 3% tallow was included in the diets. No treatment effects were observed ($P > .10$) for 12th rib fat thickness, quality grade, liver score (data not shown), or yield grade (data not shown). No significant WCGF level \times tallow level interaction ($P > .10$) was detected for any measure.

Net energy for gain estimated from dietary ingredients (1.38 Mcal/kg; NRC, 1984) was higher than the value calculated from animal performance (1.23 Mcal/kg) for the control diet. Greater precipitation over the feeding period may be responsible for reduced animal performance in this trial. Adding WCGF to the diet increased NE_g ($P < .01$), as did the addition of tallow ($P < .01$). The NE_g value of WCGF, when tallow was not fed, was calculated to be 1.91 Mcal/kg, which is 23% greater than the NE_g value of DRC (1.55 Mcal/kg; NRC, 1984). The NE_g value of tallow, when WCGF was not fed, was calculated to be 4.38 Mcal/kg, which is similar to NRC (1996).

Intake of DIP exceeded the requirement for all treatments and was greater ($P < .01$) for WCGF diets than for DRC diets. The requirement for DIP was increased ($P < .01$) by the inclusion of WCGF. Increases in DIP requirements due to additions of WCGF were an extension of the pH adjustment in the model during the calculation of the requirement. The ADG from the 50% DRC:50% WCGF + tallow treatment (1.80 kg/d) was used to calculate the MP requirement. Metabolizable protein supplied increased with the addition of WCGF ($P < .01$). The estimation of TDN from NE_g may have overestimated the energy available for microbial growth because the energy from tallow is included but is not available for microbial growth. Nocek and Russell (1988) reported that ruminal bacteria only grow on ruminally

degraded carbohydrates and do not use fats as an energy source. The inclusion of energy from tallow increases the DIP requirement and increases the estimate of MP due to a greater microbial growth.

The NE_g value of the WCGF and tallow combination was 2.00 Mcal/kg, which is 29% greater than DRC and approaches the energy value of distillers by-products, as reported by Larson et al. (1993) and Ham et al. (1994). Higher production responses reported for wet distillers grains than for WCGF may be attributed to highly digestible fiber, lipid, or escape protein. Both products are fibrous by-products, but wet distillers grains have a higher lipid content (15 vs 9%; Ham et al., 1995) and are higher in escape protein (47 vs 26%, DM basis; Firkins et al., 1984). Lipid content of the WCGF and DRC in our trial was 5.6 and 5.3%, respectively. The differences in lipid content and escape protein between WCGF and distillers by-products were evaluated by Lodge et al. (1997). In that trial, a composite of corn gluten feed, tallow, and corn gluten meal was fed. Wet corn gluten feed or the composite was fed as 40% (DM basis) of a DRC finishing diet. Steers consuming the composite were more efficient than steers consuming WCGF.

Implications

Wet corn gluten feed and distillers grains are fibrous by-products of grain processing. Wet corn gluten feed has been fed to replace dry-rolled corn in finishing cattle diets without causing a decrease in gains or efficiencies. A combination of tallow and wet corn gluten feed seems to further improve feed efficiency. Type of protein supplementation seems to be influenced by several animal and dietary factors that influence metabolizable protein and degradable intake protein supplies and requirements.

Literature Cited

- DeHaan, K. A. 1983. Improving the utilization of fiber and energy through the use of corn gluten feed and alkali compounds. Ph.D. Dissertation. Univ. of Nebraska, Lincoln.
- Elanco Products Company. 1974. Tylan premix for beef cattle. Technical B Manual. pp 4-5. Indianapolis, IN.
- Farlin, S. D. 1981. Wet distillers grains for finishing cattle. *Anim. Nutr. Health* 36 (No. 3):35.
- Firkins, J. L., L. L. Berger, and G. C. Fahey, Jr. 1985. Evaluation of wet and dry distillers grains and wet and dry corn gluten feeds for ruminants. *J. Anim. Sci.* 60:847-860.
- Firkins, J. L., L. L. Berger, G. C. Fahey, Jr., and N. R. Merchen. 1984. Ruminant nitrogen degradability and escape of wet and dry distillers grains and wet and dry corn gluten feeds. *J. Dairy Sci.* 67:1936-1944.
- Ham, G. A., R. A. Stock, T. J. Klopfenstein, and R. P. Huffman. 1995. Determining the net energy value of wet and dry corn gluten feed in beef growing and finishing diets. *J. Anim. Sci.* 73: 353-359.
- Ham, G. A., R. A. Stock, T. J. Klopfenstein, E. M. Larson, D. H. Shain, and R. P. Huffman. 1994. Wet corn distillers byproducts compared with dried corn distillers grains with solubles as a source of protein and energy for ruminants. *J. Anim. Sci.* 72: 3246-3257.

- Krehbiel, C. R., R. A. Stock, D. W. Herold, D. H. Shain, G. A. Ham, and J. E. Carulla. 1995. Feeding wet corn gluten feed to reduce subacute acidosis in cattle. *J. Anim. Sci.* 73:2931-2939.
- Larson, E. M., R. A. Stock, T. J. Klopfenstein, M. H. Sindt, and R. P. Huffman. 1993. Feeding value of wet distillers byproducts for finishing ruminants. *J. Anim. Sci.* 71:2228-2236.
- Lodge, S. L., R. A. Stock, T. J. Klopfenstein, D. H. Shain, and D. W. Herold. 1997. Evaluation of wet distillers composite for finishing ruminants. *J. Anim. Sci.* 75:44-50.
- McCoy, R. A. 1996. Receiving and finishing programs for calves. Ph.D. Dissertation. Univ. of Nebraska, Lincoln.
- Moore, J. A., R. S. Swingle, and W. H. Hale. 1986. Effects of whole cottonseed, cottonseed oil or animal fat on digestibility of wheat straw diets by steers. *J. Anim. Sci.* 63:1267-1273.
- Nocek, J. E., and J. B. Russell. 1988. Protein and energy as an integrated system. Relationship of ruminal protein and carbohydrate availability to microbial protein synthesis and milk production. *J. Dairy Sci.* 71:2070-2107.
- NRC. 1984. *Nutrient Requirements of Beef Cattle* (6th Ed.). National Academy Press, Washington, DC.
- NRC. 1996. *Nutrient Requirements of Beef Cattle* (7th Ed.). National Academy Press, Washington, DC.
- Richards, C. J. 1996. Evaluation of grains and grain byproducts for ruminants. M.S. Thesis. Univ. of Nebraska, Lincoln.
- Russell, J. B., J. D. O'Connor, D. G. Fox, P. J. Van Soest, and C. J. Sniffen. 1992. A net carbohydrate and protein system for evaluating cattle diets: I. Ruminal fermentation. *J. Anim. Sci.* 70:3551-3561.
- SAS. 1989. *SAS User's Guide Statistics* (6th Ed.). SAS Inst. Inc., Cary, NC.
- Steel, R.G.D., and J. H. Torrie. 1980. *Principles and Procedures of Statistics: A Biometrical Approach* (2nd Ed.). McGraw-Hill Publishing Co., New York.
- Stock, R. A., M. H. Sindt, J. C. Parrott, and F. K. Goedecken. 1990. Effects of grain type, roughage level and monensin level on finishing cattle performance. *J. Anim. Sci.* 68:3441-3455.
- Strobel, H. J., and J. B. Russell. 1986. Effect of pH and energy spilling on bacterial protein synthesis by carbohydrate limited cultures of mixed rumen bacteria. *J. Dairy Sci.* 69:2941-2947.
- Trenkle, A. 1986a. Comparison of wet and dry corn gluten feed when used to replace corn and corn silage in a ration for yearling steers. *Iowa State Univ. Beef Cattle Res. Rep.* RA402:23-28.
- Trenkle, A. 1986b. Feeding value of wet and dry corn gluten feed when used to replace corn and corn silage in a ration for yearling heifers. *Iowa State Univ. Beef Cattle Res. Rep.* RA401:19-22.
- Trenkle, A. 1992. Protein requirements of steers implanted with Synovex S and Finaplix S. *Iowa State Beef-Sheep Res. Rep.* AS-620:102-107.