University of Nebraska - Lincoln DigitalCommons@University of Nebraska - Lincoln

USGS Staff -- Published Research

US Geological Survey

1-1-2012

Effects of Mining-Derived Metals on Riffle-Dwelling Crayfish in Southwestern Missouri and Southeastern Kansas, USA

Ann L. Allert
United States Geological Survey, aallert@usgs.gov

Robert J. DiStefano Central Regional Office and Conservation Research Center

Christopher J. Schmitt *United States Geological Survey*, cjschmitt@usgs.gov

James F. Fairchild *United States Geological Survey*, jfairchild@usgs.gov

William G. Brumbaugh *United States Geological Survey*, bbrumbaugh@usgs.gov

Follow this and additional works at: http://digitalcommons.unl.edu/usgsstaffpub

Allert, Ann L.; DiStefano, Robert J.; Schmitt, Christopher J.; Fairchild, James F.; and Brumbaugh, William G., "Effects of Mining-Derived Metals on Riffle-Dwelling Crayfish in Southwestern Missouri and Southeastern Kansas, USA" (2012). USGS Staff -- Published Research. Paper 530.

http://digitalcommons.unl.edu/usgsstaffpub/530

This Article is brought to you for free and open access by the US Geological Survey at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in USGS Staff -- Published Research by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Effects of Mining-Derived Metals on Riffle-Dwelling Crayfish in Southwestern Missouri and Southeastern Kansas, USA

Ann L. Allert · Robert J. DiStefano · Christopher J. Schmitt · James F. Fairchild · William G. Brumbaugh

Received: 27 April 2012/Accepted: 31 July 2012/Published online: 8 September 2012 © Springer Science+Business Media, LLC (outside the USA) 2012

Abstract Riffle-dwelling crayfish populations were sampled at 16 sites in 4 tributaries of the Spring River located within the Tri-State Mining District in southwest Missouri. Crayfish density, physical habitat quality, and water quality were examined at each site to assess the ecological effects of mining-derived metals on crayfish. Metals (lead, zinc, and cadmium) were analyzed in samples of surface water, sediment, detritus, and whole crayfish. Sites were classified a posteriori into reference, mining, and downstream sites primarily based on metal concentrations in the materials analyzed. Three species of crayfish (Orconectes neglectus neglectus, O. macrus, and O. virilis) were collected during the study; however, only O. n. neglectus was collected at all sites. Mean crayfish densities were significantly lower at mining sites than at reference sites. Mean concentrations of metals were significantly correlated among the materials analyzed and were significantly greater at mining and downstream sites than at reference sites. Principal component analyses showed a separation of sites due to an inverse relationship among crayfish density, metals concentrations, and physical habitat quality variables. Sediment probable-

Electronic supplementary material The online version of this article (doi:10.1007/s00244-012-9797-9) contains supplementary material, which is available to authorized users.

A. L. Allert (⊠) · C. J. Schmitt · J. F. Fairchild · W. G. Brumbaugh Columbia Environmental Research Center, United States Geological Survey, 4200 New Haven Road, Columbia, MO 65201, USA e-mail: aallert@usgs.gov

R. J. DiStefano Missouri Department of Conservation, Central Regional Office and Conservation Research Center, 3500 East Gans Road, Columbia, MO 65201, USA effects quotients and surface-water toxic unit scores were significantly correlated; both indicated risk of toxicity to aquatic biota at several sites. Metals concentrations in whole crayfish at several sites exceeded concentrations known to be toxic to carnivorous wildlife. Mining-derived metals have the potential to impair ecosystem function through decreased organic matter processing and nutrient cycling in streams due to decreased crayfish densities.

The Tri-State Mining District (TSMD) occupies an area of approximately 647,500 ha in southwestern Missouri, southeastern Kansas, and northeastern Oklahoma. The TSMD was mined for zinc (Zn) and lead (Pb) for >150 years, beginning in the mid-1800s and ending in the late 1960s, with peak production occurring during World War II (Stewart 1986). Many sites in the TSMD are contaminated by wastes from historical mining, ore processing, and smelting (Barks 1977; Czarnezki 1985; Davis and Schumacher 1992), which has resulted in effects on aquatic organisms and potential risk to humans and wildlife (Angelo et al. 2007; Brumbaugh et al. 2005; MacDonald et al. 2010; Schmitt et al. 2006; Wang et al. 2010; Wildhaber et al. 1997, 2000). The United States Environmental Protection Agency (USEPA) has established four Superfund National Priority List (NPL) sites within the TSMD.

Crayfish are an important structural component of many aquatic systems, including Ozark streams, where they have been found to be the predominant macroinvertebrate (Rabeni 1985). Crayfish also play an integral role in stream function and the cycling of nutrients and energy through stream food webs by shredding organic matter (Creed 1994; Momot et al. 1978; Momot 1995; Rabeni et al. 1995) and as an important prey item for fish and other aquatic and terrestrial vertebrates (DiStefano 2005 and references



therein; Hobbs 1993; Rabeni et al. 1995). Recent research has showed that crayfish significantly affect aquatic microhabitats by way of ecosystem engineering (Zhang et al. 2004). Therefore, effects on crayfish may result in changes in the structure and function of Ozark stream ecosystems.

The objectives of this study were to determine crayfish species composition and densities in riffle habitats at selected stream sites in the TSMD; to evaluate crayfish densities relative to concentrations of the mining-derived metals, Pb, Zn, and cadmium (Cd), in surface water, sediment, detritus, and whole crayfish; to characterize physical habitat and water quality; and to evaluate the potential effects of metals in crayfish to carnivorous wildlife.

Materials and Methods

Crayfish were sampled in riffle habitats of tributaries of the Spring River of southwestern Missouri and southeastern Kansas. The 16 sample sites were located in Jenkins Creek, Center Creek, Turkey Creek, and Shoal Creek (Table 1; Fig. 1). Sites were selected based on stream order (Strahler 1952) and proximity to areas with mine waste. Sites were classified based on distances downstream from known

sources of mining-related contaminants and metal concentrations in the materials analyzed from each site. Sites considered upstream of contaminated areas where metals concentrations were low (e.g., reference sites) included J1, C1, T1, T2, S1, and S2; sites directly downstream of highly contaminated areas where metals concentrations were high (e.g., mining sites) included C2, C3, C4, T3, S3, and S6; and sites considered to be further downstream of mining or less contaminated areas where metals concentrations were intermediate (e.g., downstream sites) included C5, T4, S4, and S5.

Crayfish were collected by disturbing the substrate inside a 1 m² weighted polyvinyl chloride (PVC) quadrat frame placed on the stream bottom directly upstream of a kick seine (1.5 m length \times 1.5 m height) with a 3 mm diameter delta mesh on one date per site during the period of July 13 to 29, 2009 (15 sites) or on September 3, 2009 (Site S6). Each site consisted of a stream reach containing three riffles. Eight kick-seine subsamples were randomly located in each riffle (total n = 24/site). Crayfish were identified as to species (Pfleiger 1996), and carapace length (CL) was measured (to the nearest 0.1 mm) from the tip of rostrum to the posterior edge of the cephalothorax. All crayfish except those retained for metal analyses were released alive to the stream.


Table 1 Sampling locations in Jenkins, Center, Turkey, and Shoal creeks in southwestern Missouri and southeastern Kansas, USA

Stream/ site	Latitude	Longitude	Gradient (m/km)	Drainage area (ha)	Upstream tailings area (ha)	Tailings area/ drainage area	Distance downstream from tailings (km)	Stream order	Site group
Jenkins cı	reek								
J1	37°04′33.4″	94°15′40.1″	5.01	9,324	0.70	0.00008	5.1	3	R
Center cre	eek								
C1	37°06′46.9″	94°18′02.9″	4.36	29,526	27.8	0.00094	15	5	R
C2	37°10′46.6″	94°27′56.2″	2.52	66,045	588	0.00891	3.4	5	M
C3	37°10′04.6″	94°32′21.3″	2.52	67,340	785	0.01166	0.6	5	M
C4	37°10′45.8″	94°28′44.7″	2.46	73,297	815	0.01112	3.9	5	M
C5	37°09′05.4″	94°36′50.7″	2.72	77,182	908	0.01176	1.0	5	D
Turkey cr	reek								
T1	$37^{\circ}05'24.1''$	94°27′27.9″	8.98	4,144	81.7	0.01971	2.7	2	R
T2	37°06′38.4″	94°31′17.2″	3.66	6,734	143	0.02128	4.1	3	R
T3	37°06′51.6″	94°32′44.9″	3.63	9,324	225	0.02417	1.1	4	M
T4	37°07′46.9″	94°37′35.2″	4.79	11,396	318	0.02790	4.5	4	D
Shoal cree	ek								
S 1	36°56′34.7″	94°18′03.9″	3.36	69,671	96.0	0.00138	6.8	5	R
S2	37°01′25.7″	94°31′11.0″	2.84	84,952	120	0.00142	5.1	5	R
S3	37°02′07.7″	94°35′16.5″	1.52	113,441	181	0.00160	2.6	5	M
S4	37°02′24.0″	94°36′30.1″	2.32	114,736	209	0.00182	2.4	5	D
S5	37°02′31.2″	94°39′08.0″	3.08	116,549	213	0.00183	7.1	5	D
S 6	37°02′33.8″	94°39′23.6″	3.08	117,585	345	0.00294	0.5	5	M

R reference site, M mining site, D downstream site



Fig. 1 Study sites, mine-related waste (i.e., chat piles), city of Joplin boundary, and designated areas within USEPA NPL (Superfund) sites, as well as drainages, which are distinguished by gray shades



Water depth and current velocity were measured at each 1 m^2 kick-seine quadrat just before crayfish sampling using a Marsh-McBirney 2000 portable flow meter (Hach Inc., Loveland, Colorado, USA) at 6/10 of measured depth (Bain and Stevenson 1999). Substrate particle size was visually estimated at five random points within a 0.5 \times 0.5 m grid of each quadrat and assigned a value based on six coarseness categories (Bain et al. 1985).

A Hydrolab Quanta meter (Hach Inc., Loveland, Colorado, USA) was used to measure temperature, pH, specific conductance, dissolved oxygen, and turbidity in each riffle sampled for crayfish. A surface-water grab sample was collected from each riffle for analyses of metals (Pb, Zn, and Cd), alkalinity, hardness, and sulfate concentration. Alkalinity and hardness were measured by titration (American Public Health Association 2005). Sulfate concentrations were determined by colorimetric detection using a Hach 2100 spectrophotometer (Hach Inc., Loveland, Colorado, USA). A subsample of each surface-water sample was filtered on-site into a precleaned polyethylene bottle using a polypropylene syringe and filter cartridge (0.45 µm pore size), placed on ice, refrigerated, and subsequently acidified to 1 % (v/v) with nitric acid within 4 days of collection. Water samples were analyzed for Pb, Zn, and Cd using semiquantitative multielement inductively-coupled plasma-mass spectrometry (ICP-MS).

Crayfish were subsampled from those collected during kick-seine sampling for metals analyses. Generally, five crayfish from each riffle at each site were composited for analysis; however, samples from three sites (T4, S1, and S4) contained only three crayfish per riffle. Crayfish were rinsed with site water and placed in precleaned high-density polyethylene (HDPE) containers. Detritus (e.g., weathered leaves) was collected with a kick net and by hand on the day of crayfish sampling at the stream banks of each riffle sampled and placed in precleaned 125 mL HDPE bottles. Samples of crayfish and detritus were placed on ice, frozen (-20 °C) within 6 h of collection, and stored frozen until analyzed. Animal tissues and organic material were lyophilized (i.e., freeze-dried) and decreased to a coarse powder by mechanical crushing in a glass vial with a glass rod. Neither exoskeletons nor gut contents of the crayfish were removed before analyses. A dry mass of 0.25 g from each composite sample of crayfish or detritus sample was digested using concentrated nitric acid and microwave heating before being analyzed for Pb, Zn, and Cd using ICP-MS (Allert et al. 2008, 2009a; Besser et al. 2007). Surficial sediments (approximately the top 10 cm) were collected within the wetted stream channel of each riffle at each site using PVC scoops rinsed with site water before collection (Besser et al. 2009b). Sediment samples were composited and wet-sieved through a 2 mm stainlesssteel mesh sieve in the field to remove coarse particles using a minimal quantity of site water (Besser et al. 2009b; Brumbaugh et al. 2007); placed in precleaned 125 mL glass bottles on ice; then refrigerated (-4 °C) until analyzed. In the laboratory, a subsample of the <2 mm sediment sample was sieved using a 250 µm stainless-steel



mesh sieve. Total recoverable (TR) metals (Pb, Zn, and Cd) in the <2 and <250 μ m sediment fraction were analyzed by ICP-MS (Brumbaugh et al. 2007; May et al. 1997).

Site-mean TR metals concentrations in the <250 μ m sediment fraction were converted to probable-effects quotients (PEQs) by dividing site-mean TR metals concentrations by TSMD-specific probable-effects concentrations (PECs; MacDonald et al. 2000, 2010) for each metal. Individual PEQs for the three metals were summed (Σ PEQs) to estimate risks from the metal mixtures (Besser et al. 2009a; Ingersoll et al. 2001). In general, Σ PEQs greater than one are assumed to indicate a greater hazard of toxic effects (Besser et al. 2009a); however, MacDonald et al. (2010) calculated low-risk (Σ PEQ = 6.47) and high-risk (Σ PEQ = 10.04) toxicity thresholds for Pb, Zn, and Cd in TSMD sediments to better assess risks of toxicity in the TSMD.

The cumulative risk of toxic effects from metals in surface water was estimated using a toxic units (TU) approach (Wildhaber and Schmitt 1996). A TU is defined as the measured concentration of each dissolved metal in surface water divided by the chronic ambient water-quality criterion for the metal adjusted for hardness and the dissolved fraction of metal (USEPA 2005, 2006). TUs for metals were summed to produce a total toxicity estimate of the mixture (i.e., TU unit score = \sum TU) for site means; values >1.0 generally indicate potential toxicity to aquatic biota.

Increased concentrations of metals in fish and crayfish within the TSMD represent a risk to fish and carnivorous wildlife (Schmitt et al. 2006, 2008). The screening-level criteria developed by Schmitt et al. (2006) were used to assess the potential hazards of metals in crayfish collected during this study to carnivorous wildlife. Toxicity thresholds of metals in target species were determined through food-chain analysis using procedures developed for conducting ecological-risk assessments (USEPA 1992, 1993, 1997, 1999, 2007). This assessment used representative bird and mammal species based on body weight, such as the American robin (Turdus migratorius) and short-tailed shrew (Blarina brevicauda), which can be extrapolated to similar-sized species that consume crayfish. Hazard quotients (HQs) were calculated using site mean concentrations of Pb, Zn, and Cd in crayfish and no-effect hazard concentrations (NEHCs) to estimate daily contaminant intake rates; NEHCs are consensus-based no-adverse effect level-based toxic reference values normalized for estimated daily food ingestion rates (Schmitt et al. 2008). All assume a diet of 100 % crayfish. There is no analogous procedure to assess risk to predatory fish.

All statistical analyses were conducted using Statistical Analysis System (release 9.1; SAS, Chicago, IL) for Windows. Before analyses, data were tested for normality and homogeneity of variance using Shapiro-Wilk statistic. Data were not normally distributed; therefore, all analyses were conducted using rank-transformed data (Conover and Iman 1981). Ranks of site means for crayfish density, water-quality variables, physical habitat quality variables, and metal concentrations were used in the statistical analyses. Censored values [less than method detection limit (MDL); Supplemental Table S1] were replaced with 50 % of the MDL for statistical computations, figures, and tables; only metal concentrations in surface water samples from reference sites were censored.

Differences among groups of sites (i.e., reference, mining, downstream) were tested using nested analysis of variance (ANOVA; riffles within sites) with site considered a fixed effect. Differences in measured variables were tested as planned nonorthogonal contrasts using single df F-tests. The within-site mean squares for ranked variables were used in all tests, which were conducted using PROC GLM. Associations between site means metal concentrations were examined using Spearman's rank correlation analyses (Supplemental Table S2) and indicated a high degree of correlation between metals. Separate linear regression of O. n. neglectus and O. macrus densities against physical and chemical habitat variables was performed using PROC REG with variable selection based on Akaike's information criterion (AIC; Burnham and Anderson 2002). In these analyses, models were evaluated relative to each other based on corrected AIC values (AICc). The AICc values are adjusted upward for sample size relative to the number of independent variables, which protects against over-fitting models due to small sample size (Burnham and Anderson 2002). The best-fit equally fit model (variables n = 30) for O. n. neglectus density included five variables: current velocity; Pb and Cd in detritus; temperature and pH; $R^2 = 0.89$; and AICc = 55.65. The best two equally fit models for O. macrus density included two variables: alkalinity and current velocity ($R^2 = 0.99$; AICc = -6.68) or Pb in surface water and Zn in crayfish ($R^2 = 0.99$; AICc = -5.99). The relationship among O. n. neglectus and O. macrus densities and the variables identified in the best-fit models were further examined with principal component analyses (PCA). A significance level of P < 0.05 was used to judge all statistical tests. Supplemental material contains site mean summaries.

Results

Crayfish were collected at all sites (Supplemental Table S3). Only one crayfish species, *Orconectes (Procericambarus) neglectus neglectus* Faxon (ringed crayfish), was



Table 2 Mean densities of O. n. neglectus, water depth, and $\sum PEQ$ and $\sum TU$ for Pb, Zn, and Cd

Site group	O. n. neglectus density (n/m²)	∑PEQ	∑TU	Water depth (cm)
Reference ^a	16.9 (4.0)	9.2 (2.1)	0.23 (0.07)	21.8 (2.7)
Mining ^b	4.79 (1.33)	38.7 (6.3)	1.14 (0.14)	26.4 (2.4)
Downstream ^c	1.68 (0.41)	22.3 (4.0)	0.57 (0.15)	24.8 (1.9)
R versus M	$F_{(15,32)} = 15.0**$	$F_{(15,32)} = 352**$	$F_{(15,32)} = 2,191**$	$F_{(15,32)} = 6.41**$
R versus D	$F_{(15,32)} = 33.6**$	$F_{(15,32)} = 111**$	$F_{(15,32)} = 484**$	$F_{(15,32)} = 4.88*$
D versus M	$F_{(15,32)} = 5.44*$	$F_{(15,32)} = 39.3**$	$F_{(15,32)} = 395**$	$F_{(15,32)} = 0.00 \text{ ns}$

Data shown are arithmetic site groups means \pm SEs. Also shown are results of one-way ANOVA as F values and df for differences among site groups (** $p \le 0.01$; * $0.01 \le p \le 0.05$; ns ≥ 0.05). Within-site mean squares for ranked variables were used in all tests

R reference site, M mining site, D downstream site

collected at all sites. There were significant differences in mean densities of O. n. neglectus among site types; mean densities were significantly lower at mining and downstream sites than reference sites (Table 2). Orconectes (Procericambarus) macrus Williams (Neosho midget crayfish) was only collected at reference sites (J1, C1, S1, and S2). Mean densities of O. macrus were threefold greater than O. n. neglectus densities at C1; however, at the three other sites where both species were collected (J1, S1, and S2), mean densities of O. n. neglectus were 1.5- to 5-fold greater than O. macrus (Supplemental Table S3). A third species, Orconectes (Gremicambarus) virilis Hagen (virile crayfish), was collected only at J1 (virile 3) and C1 (virile 2).

Preliminary analyses showed there were no significant differences in mean CL between sexes for either $O.\ n.$ neglectus $(F_{(1,226)}=2.29;\ P=0.13)$ or $O.\ macrus$ $(F_{(1,54)}=3.37;\ P=0.06)$. However, there were significant differences in mean CL among sites $(F_{(1,226)}=2.29;\ P=0.13)$ and site groups for $O.\ n.\ neglectus$ collected for metal analyses: Crayfish from reference sites were significantly larger than crayfish collected at mining or downstream sites (reference vs. mining $F_{(3,15)}=159;\ P<0.001;$ reference vs. downstream $F_{(3,15)}=123;\ P<0.001;$ and mining vs. downstream $F_{(3,15)}=0;\ P=0.954)$. O. macrus were only collected at reference sites; however, there were no significant differences in CLs of $O.\ macrus$ collected for metals analyses among sites $(F_{(3,52)}=0.46;\ P=0.714)$.

Mean concentrations of Pb, Zn, and Cd in O. n. neglectus, surface water, <250 µm sediment fraction, and detritus were significantly correlated (Supplemental Table S2) and significantly greater at mining and downstream sites than at reference sites (Table 3; Supplemental Tables S4 through S7). Concentrations of metals in O. macrus were significantly greater than those in O. n. neglectus in

11 of the 12 comparisons (Supplemental Table S4). Mean concentrations of Pb, Zn, and Cd in the <250 μ m sediment fraction approached or exceeded TSMD-specific high-risk concentrations (PEC₂₀; concentrations associated with a 20 % decrease in a measured end point at all site groups (Table 3; Supplemental Table S6; MacDonald et al. 2010). Mean concentrations of Pb in surface water did not exceed the chronic USEPA water-quality criterion at any site group; however, mean concentration of Zn and Cd did exceed these criteria at mining sites (Table 3).

Sediment \sum PEQs indicated risk of toxicity at mining and downstream sites (Table 2; Supplemental Table S8); \sum PEQs were greater than both the TSMD-specific low-(\sum PEQs = 6.47) and high-risk (\sum PEQs = 10.04) toxicity thresholds at mining and downstream sites (MacDonald et al. 2010). *O. n. neglectus* densities were decreased relative to reference sites where TSMD-specific \sum PEQs were exceeded (Fig. 2). Surface water \sum TUs were significantly greater at mining and downstream sites than at reference sites (Table 2; Supplemental Table S9). Surface water \sum TUs at all mining sites were <1, indicating potential risk to aquatic biota.

Criteria used to evaluate risks of Pb, Zn, and Cd concentrations in crayfish to wildlife indicated that metals concentrations at mining and downstream sites are potentially hazardous to carnivorous wildlife (Fig. 3). HQs were greater for birds than mammals in their respective size category (Supplemental Tables S10 and S11). HQs for mean concentrations of Pb and Zn in *O. n. neglectus* were >1.0 for robin-size birds at mining and downstream sites. The Cd hazard quotient was <1.0 for robin-sized birds and approached 1.0 for shrew-size mammals at mining sites (Fig. 3).

Mean water depth and current velocities were significantly greater at mining and downstream sites than at reference sites (Table 2; Supplemental Tables S12 and S13).



^a Reference sites: J1, C1, T1, T2, S1, and S2

b Mining sites: C2, C3, C4, T3, S3, and S6

^c Downstream sites: T4, C5, S4, and S5

Table 3 Metal concentrations in O. n. neglectus <250 µm fraction and surface water

Table 3 Metal concentrations in <i>O. n. neglectus</i> , detritus,	Site group/criteria	Pb	Zn	Cd		
<250 µm fraction of sediment,		O. n. neglectus (μg/g dry weight)				
and surface water	Reference ^a	3.65 (0.7)	159 (17)	1.22 (0.24)		
	Mining ^b	10.1 (1.7)	339 (26)	5.76 (0.77)		
	Downstream ^c	10.5 (2.6)	272 (35)	3.81 (0.94)		
	R versus M	$F_{(15,32)} = 47.3**$	$F_{(15,32)} = 120**$	$F_{(15,32)} = 531**$		
	R versus D	$F_{(15,32)} = 22.3**$	$F_{(15,32)} = 32.4**$	$F_{(15,32)} = 155**$		
	D versus M	$F_{(15,32)} = 2.05 \text{ ns}$	$F_{(15,32)} = 17.0**$	$F_{(15,32)} = 66.6*$		
Data shown are arithmetic site		Detritus (µg/g dry weig	dry weight)			
groups means ± SEs. Also	R	156 (36)	3,281 (663)	34 (11)		
shown are results of one-way ANOVA as <i>F</i> values and <i>df</i> for	M	643 (70)	10,772 (1,438)	138 (20)		
differences among site group	D	543 (124)	8,500 (1,261)	92 (20)		
$(**p \le 0.001; \text{ ns} \ge 0.05).$	R versus M	$F_{(15,32)} = 167**$	$F_{(15,32)} = 312**$	$F_{(15,32)} = 129**$		
Within-site mean squares for ranked variables were used in	R versus D	$F_{(15,32)} = 71.9**$	$F_{(15,32)} = 132**$	$F_{(15,32)} = 41.6**$		
all tests	D versus M	$F_{(15,32)} = 9.59**$	$F_{(15,32)} = 18.8**$	$F_{(15,32)} = 13.6**$		
R reference site, M mining site,		<250 μm sediment fraction (μg/g dry weight)				
D downstream site	R	235 (73)	2,093 (409)	14.0 (3.7)		
$*0.01 \le p \le 0.05$	M	947 (231)	8,991 (1,435)	58.6 (8.7)		
 a Reference sites: J1, C1, T1, T2, S1, and S2 b Mining sites: C2, C3, C4, T3, S3, and S6 	D	534 (128)	5,393 (1,011)	31.9 (4.2)		
	R versus M	$F_{(15,32)} = 155**$	$F_{(15,32)} = 235**$	$F_{(15,32)} = 341**$		
	R versus D	$F_{(15,32)} = 51.2**$	$F_{(15,32)} = 54.5**$	$F_{(15,32)} = 93.4**$		
^c Downstream sites: T4, C5, S4,	D versus M	$F_{(15,32)} = 15.9**$	$F_{(15,32)} = 40.1**$	$F_{(15,32)} = 47.1**$		
and S5	PEC ₂₀ ^d	179	2,409	14.1		
^d PEC ₂₀ = TSMD probable-		Surface water (µg/L)				
effects concentration high-risk	R	0.17 (0.04)	43.2 (13.4)	0.14 (0.17)		
threshold associated with a	M	0.56 (0.13)	235 (33)	0.91 (0.16)		
20 % decrease in a measured end point (MacDonald et al.	D	0.40 (0.09)	93.9 (25.1)	0.43 (0.16)		
2010)	R versus M	$F_{(15,32)} = 17.4**$	$F_{(15,32)} = 8.68**$	$F_{(15,32)} = 7.41**$		
^e WQC = federal water-quality	R versus D	$F_{(15,32)} = 180**$	$F_{(15,32)} = 373**$	$F_{(15,32)} = 439**$		
criteria adjusted for hardness	D versus M	$F_{(15,32)} = 93.7**$	$F_{(15,32)} = 482**$	$F_{(15,32)} = 343**$		
and the dissolved fraction of the metal (USEPA 2006)	WQC ^e	5–9	93–145	0.4–0.6		

Substrate coarseness was not significantly difference among site groups (Supplemental Tables S12 and S13). In situ water-quality variables generally did not exceed state or federal criteria (Supplemental Tables S14 and S15). However, water-quality constituents associated with metalmining, such as specific conductance and concentrations of hardness and sulfate, were significantly greater at mining than at reference sites (Supplemental Tables S16 and S17).

A series of interpretable ordinations were obtained by PCA for each species (Fig. 4). Each ordination resulted in the first two PCA axes explaining more of the variation (69-90 %) than expected by chance alone. In the ordination for the O. n. neglectus, variables with strong positive associations with PC1 included Pb (0.67) and Cd (0.64) in detritus. Current velocity (0.58) had a strong positive association, but density (-0.65) had a strong negative association with PC2. Reference sites were separated from all other sites with negative values for PC1, whereas mining sites had positive values. In the ordination for O. macrus, density (-0.45) had a strong negative association with PC1, whereas, temperature (0.47) and alkalinity (0.41) had a strong positive association with PC1. Variables that had strong positive associations with PC2 included Pb in surface water (0.66) and Zn in the <250 µm fraction of sediment (0.47). Current velocity had a strong negative association (-0.46). These ordinations suggest that habitat variables and metals are important in explaining densities of O. n. neglectus and O. macrus among sites.

Discussion

Our results indicate that environmental metals concentrations remain increased in the TSMD as indicated by high metal concentrations in surface water, sediment, detritus, and crayfish in the streams investigated. We found






Fig. 2 Relationship between sediment ∑PEQs (MacDonald et al. 2010) and mean densities of *O. n. neglectus. Lines* represent TSMD low-risk (*dashed line*) and high-risk (*solid line*) toxicity thresholds (MacDonald et al. 2010)

decreased densities of O. n. neglectus and the absence of O. macrus at mining-affected sites as well as a negative association of crayfish densities with metal concentrations. Our results are consistent with previous investigations that have documented increased metal concentrations in macroinvertebrates, fish, detritus, and sediment in the TSMD (Angelo et al. 2007; Brumbaugh et al. 2005; MacDonald et al. 2010; Schmitt et al. 1993, 2006; Wildhaber et al. 1997, 2000) and that documented the absence or decreased densities of cravfish below mining-affected sites in the Viburnum Trend Mining District of southeast Missouri (Allert et al. 2008, 2009a). Allert et al. (2009a) also documented decreased survival of caged crayfish at sites directly downstream (0.4–3.7 km) of mining sites. Survival and biomass of caged crayfish were significantly lower at mining sites than reference or downstream sites, and survival was negatively correlated with metal concentrations in surface water, detritus, macroinvertebrates, stonerollers, and whole crayfish. In situ testing of crayfish was an important tool for showing that the absence of crayfish populations below mining sites was the result of metal exposure as opposed to habitat loss due to physical impairment by mine waste (e.g., sedimentation by mine tailings).

Although the diversity of crayfish in the Missouri Ozarks is high (>25 species; Pfleiger 1996), many taxa are restricted to a single drainage and are further restricted within drainages due to macrohabitat partitioning between species (DiStefano et al. 2003; Flinders and Magoulick 2005; Rabeni 1985). O. macrus is endemic to the Ozark Highlands and is primarily found in the westward-flowing rivers north of the Illinois River in Oklahoma, including the Spring River. Within the Spring River drainage, O. macrus does not occur in the North Fork Spring River or in





Fig. 3 HQs for dietary metal exposure for representative species for a Zn, b Cd, and c Pb in O. n. neglectus. Line represents risk threshold

portions of Shoal Creek (Dillman et al. 2010; Pfleiger 1996). O. macrus typically inhabits rocky substrates in swift, shallow water and is a relatively sedentary species that spends most of its time in cavities beneath rocks or in excavated tunnels in gravelly substrate (Pfleiger 1996). It is unclear why O. macrus is not found at sites in the lower reaches of Center Creek and Shoal Creek despite being present in the lower Neosho River and Spring River (Dillman et al. 2010). O. macrus may be more sensitive to metals than O. n. neglectus, or its sedentary nature within excavated tunnels may expose O. macrus to greater metal concentrations typical of porewater and sediment (Brumbaugh et al. 2007). Either or both mechanisms could have eliminated O. macrus from Center Creek and Shoal









Fig. 4 PC ordination for selected physical habitat and water-quality variables and **a** mean densities of *O. n. neglectus* and **b** mean densities of *O. macrus*. Variables strongly associated with the axes are shown: the lengths of the *lines* leading from the centroid toward the labels are proportional to the strength of the correlation of that variable with the axes

Creek where metals concentrations are high (Allert et al. 1997, 2008, 2009a; Knowlton et al. 1983; Wigginton and Birge 2007).

Previous studies differed in their characterization of habitat selection by *O. neglectus*. Gore and Bryant (1990) found that *O. neglectus* partition their habitats, with young typically found in moderate-velocity (25–45 cm/s) cobble habitats and adults in low-velocity (0 cm/s) macrophyte beds or high-velocity (>65 cm/s) cobbled habitats. However, other studies (Flinders and Magoulick 2005; Magoulick and DiStefano 2007) reported no significant difference in habitat selection by different size classes of *O. neglectus* or *O. n. chaenodactylus* (Williams 1952). The range of water depth, current velocities, and substrate in riffles measured in our study were comparable with previously reported values (Flinders and Magoulick 2005; Magoulick and DiStefano 2007; Rabalais and Magoulick



Previous studies have used crayfish to assess the bioavailability of metals through the analysis of whole-body crayfish (Allert et al. 2008, 2009a, 2010; Besser et al. 2007; Schmitt et al. 2006, 2008). Crayfish, freshwater mussels, and benthic fishes have been identified as possible sentinel organisms in the investigation of heavy-metal pollution in the environment because of their important functional role in streams and limited home ranges (Allert et al. 2008, 2009a, 2009b, 2010; Angelo et al. 2007; Maret and MacCoy 2002). Dietary pathways have been identified as important routes of metal exposure (Besser et al. 2005; Farag et al. 1999). Therefore, the likelihood of food-chain transfer of metals from crayfish and other environmental media to wildlife in the TSMD is high and potentially hazardous.

Conclusion

Crayfish densities were significantly decreased downstream of mine waste relative to upstream reference sites. Decreased crayfish densities imply a loss of ecosystem function because crayfish are a key structural and functional component of Ozark streams and their surrounding ecosystems. This study provides additional support for the use of crayfish in the assessment of the bioavailability and ecological effects of metals in aquatic ecosystems. However, risk assessments are more effective when contaminant (e.g., metals) concentrations are linked with the status of aquatic organisms (e.g., crayfish density) and toxicological information (e.g., mortality or other sublethal measurements). In situ and laboratory toxicity tests would provide additional information regarding metal exposure,



the bioaccumulation of metals, and the potential toxicity to known age classes of crayfish because it is impossible to observe temporal changes and bioaccumulation in the environment. Genomic research may provide information on the association of distribution and metal concentrations among crayfish in the TSMD.

Acknowledgments We thank the private landowners who allowed us to access the studied streams. Personnel who assisted in the study included staff from the Missouri Department of Conservation and from the United States Geological Survey (USGS) Columbia Environmental Research Center. We thank J. Besser, J. Hinck, N. Wang, and two anonymous reviewers for providing insightful comments that greatly improved the quality of this manuscript. This study was jointly funded by the USGS, the United States Department of the Interior Natural Resource Damage Assessment and Restoration Program, the Missouri Department of Conservation, and the Missouri Department of Natural Resources. This manuscript has been reviewed in accordance with USGS policy. All procedures performed conformed with USGS guidelines for the humane treatment of the test organisms during culture and experimentation. The use of trade names does not constitute USGS or United States Government endorsement.

References

- Allert AL, Wildhaber ML, Schmitt CJ, Chapman D, Callahan E (1997) Toxicity of sediments and pore-waters and their potential impact on Neosho madtom, *Noturus placidus*, in the Spring River system affected by historic zinc-lead mining and related activities in Jasper and Newton Counties, Missouri; and Cherokee County, Kansas. Final report to the United States Fish and Wildlife Service-Region 3, Columbia Missouri Ecological Services Field Office. United States Geological Survey, Columbia Environmental Research Center, Columbia
- Allert AL, Fairchild JF, DiStefano RJ, Schmitt CJ, Besser JM, Brumbaugh WG et al (2008) Effects of lead-zinc mining on crayfish (Orconectes hylas) in the Black River watershed, Missouri. Freshw Crayfish 16:97–111
- Allert AL, Fairchild JF, DiStefano RJ, Schmitt CJ, Brumbaugh WG, Besser JM (2009a) Ecological effects of lead mining on Ozark streams—in-situ toxicity to woodland crayfish (*Orconectes hylas*). Ecotoxicol Environ Saf 72:1207–1219
- Allert AL, Fairchild JF, Schmitt CJ, Besser JM, Brumbaugh WG, Olson SJ (2009b) Effects of mining-derived metals on riffledwelling benthic fishes in Southeast Missouri, USA. Ecotoxicol Environ Saf 72(6):1642–1651
- Allert AL, DiStefano RJ, Fairchild JF, Schmitt CJ, Brumbaugh WG (2010) Effects of mining-derived metals on riffle-dwelling crayfish and in-situ toxicity to juvenile *Orconectes hylas* and *Orconectes luteus* in the Big River of southeast Missouri, USA. United States Geological Survey Administrative report, submitted to the United States Fish and Wildlife Service, Columbia
- American Public Health Association, American Water Works Association, and Water Environment Federation (2005) Standard methods for the examination of water and wastewater. American Public Health Association, Washington DC
- Angelo RT, Cringan MS, Chamberlain DL, Stahl AJ, Haslouer SG, Goodrich CA (2007) Residual effects of lead and zinc mining on freshwater mussels in the Spring River Basin (Kansas, Missouri, and Oklahoma, USA). Sci Total Environ 384:467–496
- Bain MB, Stevenson NJ (1999) Aquatic habitat assessment—common methods. American Fisheries Society, Bethesda

- Bain MA, Finn JT, Booke HE (1985) Quantifying stream substrate for habitat analysis studies. N Am J Fish Manag 5:499–506
- Barks JH (1977) Effects of abandoned lead and zinc mines and tailings piles on water quality in the Joplin area, Missouri. United States Geological Survey Water Resources Investigations Report 77–75. United States Geological Survey, Rolla
- Besser JM, Brumbaugh WG, Brunson EL, Ingersoll CG (2005) Acute and chronic toxicity of lead in water and diet to the amphipod *Hyalella azteca*. Environ Toxicol Chem 24(7):1807–1815
- Besser JM, Brumbaugh WG, May TW, Schmitt CJ (2007) Biomonitoring of lead, zinc, and cadmium in streams draining leadmining and non-mining areas, southeast Missouri, USA. Environ Monit Assess 129:227–241
- Besser JM, Brumbaugh WG, Allert AL, Poulton BC, Schmitt CJ, Ingersoll CG (2009a) Ecological impacts of lead mining on Ozark streams—toxicity of sediment and pore water. Ecotoxicol Environ Saf 72:516–526
- Besser JM, Brumbaugh WG, Hardesty DK, Hughes JP, Ingersoll CG (2009b) Assessment of metal-contaminated sediments from the Southeast Missouri (SEMO) mining district using sediment toxicity tests with amphipods and freshwater mussels. United States Geological Survey Administrative report submitted to the United States Fish and Wildlife Service, Ecological Services Office, Region 3, Environmental Contaminants Division, Columbia Missouri. United States Geological Survey, Columbia
- Brumbaugh WG, Schmitt CJ, May TW (2005) Concentrations of cadmium, lead, and zinc in fish from mining-influenced waters of northeastern Oklahoma—sampling of blood, carcass, and liver for aquatic biomonitoring. Arch Environ Contam Toxicol 49:76–88
- Brumbaugh WG, May TW, Besser JM, Allert AL, Schmitt CJ (2007)
 Assessment of elemental concentrations in streams of the New
 Lead Belt in southeastern Missouri, 2002–05. United States
 Geological Survey Scientific Investigations Report 2007–5057.
 http://pubs.usgs.gov/sir/2007/5057/. Accessed 14 May 2009
- Burkey JL, Simon TP (2010) Reach- and watershed-scale associations of crayfish within an area of varying agricultural impact in West-Central Indiana. Southeast Nat 9(sp3):199–216
- Burnham KP, Anderson DR (2002) Model selection and multimodel inference: a practical information theoretic approach, 2nd edn. Springer-Verlag, New York, p 488
- Clark JM (2009) Abiotic and biotic factors affecting size-dependent crayfish (Orconectes obscures) distribution, density, and survival. Doctoral dissertation, Kent State University, Kent
- Conover WJ, Iman RL (1981) Rank transformations as a bridge between parametric and nonparametric statistics. Am Stat 35:124–129
- Creed RP Jr (1994) Direct and indirect effects of crayfish grazing in a stream community. Ecology 75:2091–2103
- Czarnezki JM (1985) Accumulation of lead in fish from Missouri streams impacted by lead mining. Bull Environ Contam Toxicol 34:736–745
- Davis JV, Schumacher JG (1992) Water quality characterization of the Spring River basin, southwestern Missouri and southeastern Kansas. United States Geological Survey Water Resources Investigations Report 90–4176. United States Geological Survey, Rolla
- Dillman CB, Wagner BK, Wood RM (2010) Phylogenetic estimation of species limits in dwarf crayfishes from the Ozarks, *Orconectes macrus* and *Orconectes nana* (Decapoda: Cambaridae). Southeast Nat 9(sp3):185–198
- DiStefano RJ (2005) Trophic interactions between Missouri Ozarks stream crayfish communities and sport fish predators—increased abundance and size structure of predators cause little change in crayfish community densities. Project F-1-R-054, Study S-41, Job 4, Final report. Missouri Department of Conservation, Columbia



- DiStefano RJ, Decoske JJ, Vangilder TM, Barnes LS (2003) Macrohabitat partitioning among three crayfish species in two Missouri streams, USA. Crustaceana 76:343–362
- DiStefano RJ, Herleth-King SS, Imhoff EI (2008) Distribution of the imperiled Meek's crayfish (Orconectes meeki meeki, Faxon) in the White River drainage of Missouri, USA—associations with multi-scale environmental variables. Freshw Crayfish 16:27–36
- Farag AM, Woodward DF, Brumbaugh WG, Goldstein JG, MacConnell E, Hogstrand C et al (1999) Dietary effects of metals-contaminated invertebrates from the Coeur d'Alene River, Idaho, on cutthroat trout. Trans Am Fish Soc 128:578–592
- Flinders CA, Magoulick DD (2003) Effects of stream permanence on crayfish community structure. Am Midl Nat 149(1):134–147
- Flinders CA, Magoulick DD (2005) Distribution, habitat use, and life history of stream-dwelling crayfish in the Spring River drainage of Arkansas and Missouri with a focus on the Mammoth Spring crayfish (*Orconectes marchandi*). Am Midl Nat 154:358–374
- Gore JA, Bryant RM Jr (1990) Temporal shifts in physical habitat of the crayfish, *Orconectes neglectus* (Faxon). Hydrobiologia 199:131–142
- Hill AM, Lodge DM (1995) Multi-trophic-level impact of sublethal interactions between bass and omnivorous crayfish. J N Am Benthol Soc 14:306–314
- Hobbs HH Jr (1993) Trophic relationships of North American freshwater crayfishes and shrimps. Milwaukee Public Museum, Milwaukee
- Ingersoll CG, MacDonald DD, Wang N, Crane JL, Field LJ, Haverland PS et al (2001) Predictions of sediment toxicity using consensus-based freshwater sediment quality guidelines. Arch Environ Contam Toxicol 41(1):8–21
- Knowlton MF, Boyle TP, Jones JR (1983) Uptake of lead from aquatic sediment by submersed macrophytes and crayfish. Arch Environ Contam Toxicol 12(5):535–541
- Lodge DM, Hill AM (1994) Factors governing species composition, population size, and productivity of cool-water crayfish. Nord J Freshw Res 69:111–136
- MacDonald DD, Ingersoll CG, Berger TA (2000) Development and evaluation of consensus-based sediment quality guidelines for freshwater ecosystems. Arch Environ Contam Toxicol 39: 20–31
- MacDonald DD, Ingersoll CG, Crawford M, Prencipe H, Besser JM, Brumbaugh WG et al (2010) Advanced screening-level ecological risk assessment (SLERA) for aquatic habitats within the Tri-State Mining District, Oklahoma, Kansas, and Missouri. Draft Final Technical Report submitted to the United States Environmental Protection Agency, Region 6, Dallas, Texas, Region 7, Kansas City, Kansas, and the United States Fish and Wildlife Service, Columbia
- Magoulick DD, DiStefano RJ (2007) Invasive crayfish *Orconectes neglectus* threatens native crayfishes in the Spring River drainage of Arkansas and Missouri. Southeast Nat 6(1):141–150
- Maret TR, MacCoy DE (2002) Fish assemblages and environmental variables associated with hardrock mining in the Coeur d'Alene river basin, Idaho. Trans Amer Fish Soc 131:865–884
- May TW, Wiedmeyer RH, Brumbaugh WG, Schmitt CJ (1997) The determination of metals in sediment pore waters and in 1 N HClextracted sediments by ICP-MS. At Spectrosc 18:133–139
- Mitchell DJ, Smock LA (1991) Distribution, life history and production of crayfish in the James River, Virginia. Am Midl Nat 126(2):353–363
- Momot WT (1995) Redefining the role of crayfish in aquatic ecosystems. Rev Fish Sci 3:33–63
- Momot WT, Gowing H, Jones PD (1978) The dynamics of crayfish and their role in ecosystems. Am Midl Nat 99(1):10–35
- Pfleiger WL (1996) The crayfishes of Missouri. Missouri Department of Conservation, Jefferson

- Rabalais MR, Magoulick DD (2006) Influence of an invasive crayfish species on diurnal habitat use and selection by a native crayfish species in an Ozark stream. Am Midl Nat 155:295–306
- Rabeni CF (1985) Resource partitioning by stream-dwelling crayfish—the influence of body size. Am Midl Nat 113:20–29
- Rabeni CF, Gossett M, McClendon DD (1995) Contribution of crayfish to benthic invertebrate production and trophic ecology of an Ozark stream. Freshw Crayfish 10:163–173
- Schmitt CJ, Wildhaber ML, Hunn JB, Nash T, Tieger MN, Steadman BL (1993) Biomonitoring of lead-contaminated Missouri streams with an assay for erythrocyte δ-aminolevulinic acid dehydratase activity in fish blood. Arch Environ Contam Toxicol 25:464–475
- Schmitt CJ, Brumbaugh WG, Linder GL, Hinck JE (2006) A screening-level assessment of lead, cadmium, and zinc in fish and crayfish from northeastern Oklahoma, USA. Environ Geochem Health 38:445–471
- Schmitt CJ, Brumbaugh WG, Besser JM, Hinck JE, Bowles DE, Morrison LW et al (2008) Protocol for monitoring metals in Ozark National Scenic Riverways, Missouri, Version 1.0. United States Geological Survey Open-File Report 2008–1269. http://pubs.usgs.gov/of/2008/1269/. Accessed 14 May 2009
- Stein RA, Magnuson JJ (1976) Behavioral response of crayfish to a fish predator. Ecology 57(4):751–761
- Stewart DR (1986) A brief description of the historical, ore production, mine pumping, and prospecting of the Tri-State Zinc-Lead District of Missouri, Kansas, and Oklahoma. In: Hagni RD (ed) Guidebook to the geology and environmental concerns in the Tri-States Lead-Zinc District, Missouri, Kansas, Oklahoma. Division of Geology and Land Survey-Department of Natural Resources, Department of Geology and Geophysics, University of Missouri-Rolla, Rolla, pp 16–29
- Strahler AN (1952) Dynamic basis of geomorphology. Geol Soc Am Bull 63:305–923
- United States Environmental Protection Agency (1992) Framework for ecological risk assessment. EPA-630-R-92-001. USEPA, Washington DC
- United States Environmental Protection Agency (1993) Wildlife exposure factors handbook. EPA-600-R-93-187. USEPA, Washington DC
- United States Environmental Protection Agency (1997) Ecological risk assessment guidance for Superfund—process for designing and conducting ecological risk assessments. EPA-540-R-97-006. USEPA, Washington DC
- United States Environmental Protection Agency (1999) Screening level ecological risk assessment protocol for hazardous waste combustion facilities. Peer draft review, v. 1. EPA-530-D-99-001A. USEPA, Washington DC
- United States Environmental Protection Agency (2005) Procedures for the derivation of equilibrium partitioning sediment benchmarks (ESBs) for the protection of benthic organisms: Metal mixtures (cadmium copper, lead nickel, silver, and zinc). EPA-600-R-Q2-011, USEPA, Office of Research and Development, Washington DC
- United States Environmental Protection Agency (2006) National recommended water quality criteria, updated 2009. http://www. epa.gov/waterscience/criteria/wqctable/index.html. Accessed 15 May 2009
- United States Environmental Protection Agency, Office of Solid Waste and Emergency Response (2007) Ecological soil screening levels for nickel. Interim final OSWER directive 9285.7–76. http://www.epa.gov/ecotox/ecossl/pdf/eco-ssl_nickel.pdf. Acces sed 15 May 2009
- Vannote RL, Minshall GW, Cummins KW, Sedell JR, Cushing CE (1980) The river continuum concept. Can J Fish Aquat Sci 37:130–137



- Wang N, Ingersoll CG, Ivey CD, Hardesty DK, May TW, Augspurger T et al (2010) Sensitivity of early life stages of freshwater mussels (Unionidae) to acute and chronic toxicity of lead, cadmium, and zinc in water. Environ Toxicol Chem 29:2053–2063
- Westhoff JT, Guyot JA, DiStefano RJ (2006) Distribution of the imperiled Williams' crayfish (*Orconectes williamsi*) in the White River drainage of Missouri—associations with multi-scale environmental variables. Am Midl Nat 156(2):273–288
- Wigginton AJ, Birge WJ (2007) Toxicity of cadmium to six species in two genera of crayfish and the effect of cadmium on molting success. Environ Toxicol Chem 26:548–554
- Wildhaber ML, Schmitt CJ (1996) Estimating aquatic toxicity as determined through laboratory tests of Great Lakes sediments containing complex mixtures of environmental contaminants. Environ Monit Assess 41:255–289
- Wildhaber ML, Schmitt CJ, Allert AL (1997) Elemental concentrations in benthic invertebrates from the Neosho, Cottonwood, and Spring rivers. Final report to the United States Fish and Wildlife Service, Region 6, Manhattan, Kansas. United States Geological Survey, Columbia
- Wildhaber ML, Allert AL, Schmitt CJ, Tabor VM, Mulhern D, Powell KL et al (2000) Natural and anthropogenic influences on the distribution of the threatened Neosho madtom in a midwestern warmwater stream. Trans Am Fish Soc 129:243–261
- Williams AB (1952) Six new crayfishes of the genus *Orconectes* (Decapoda: Astacidae) from Arkansas, Missouri and Oklahoma. Trans Kans Acad Sci 55(3):330–351
- Zhang Y, Richardson JS, Negishi JN (2004) Detritus processing, ecosystem engineering and benthic diversity—a test of predatoromnivore interference. J Anim Ecol 73:756–766

