## University of Nebraska - Lincoln Digital Commons @ University of Nebraska - Lincoln

USGS Staff -- Published Research

**US** Geological Survey

1-1-2009

# Ecological effects of lead mining on Ozark streams: In-situ toxicity to woodland crayfish (*Orconectes hylas*)

Ann L. Allert
United States Geological Survey, aallert@usgs.gov

James F. Fairchild *United States Geological Survey*, jfairchild@usgs.gov

Robert J. DiStefano Central Regional Office and Conservation Research Center

C.J. Schmitt *United States Geological Survey*, cjschmitt@usgs.gov

William G. Brumbaugh
United States Geological Survey, bbrumbaugh@usgs.gov

See next page for additional authors

Follow this and additional works at: http://digitalcommons.unl.edu/usgsstaffpub

Allert, Ann L.; Fairchild, James F.; DiStefano, Robert J.; Schmitt, C.J.; Brumbaugh, William G.; and Besser, J.M., "Ecological effects of lead mining on Ozark streams: In-situ toxicity to woodland crayfish (*Orconectes hylas*)" (2009). *USGS Staff -- Published Research*. Paper 531.

http://digitalcommons.unl.edu/usgsstaffpub/531

This Article is brought to you for free and open access by the US Geological Survey at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in USGS Staff -- Published Research by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors Ann L. Allert, James F. Fairchild, Robert J. DiStefano, C.J. Schmitt, William G. Brumbaugh, and J.M. Besser

ELSEVIER

Contents lists available at ScienceDirect

### Ecotoxicology and Environmental Safety

journal homepage: www.elsevier.com/locate/ecoenv



## Ecological effects of lead mining on Ozark streams: In-situ toxicity to woodland crayfish (*Orconectes hylas*)

A.L. Allert a,\*, J.F. Fairchild A, R.J. DiStefano b, C.J. Schmitt A, W.G. Brumbaugh A, J.M. Besser

#### ARTICLE INFO

Article history:
Received 22 April 2008
Received in revised form
28 July 2008
Accepted 28 August 2008
Available online 20 December 2008

Keywords:
Crayfish
Orconectes hylas
Lead-zinc mining
Lead
Zinc
Cadmium
Nickel
Cobalt
In-situ toxicity

#### ABSTRACT

The Viburnum Trend mining district in southeast Missouri, USA is one of the largest producers of lead–zinc ore in the world. Previous stream surveys found evidence of increased metal exposure and reduced population densities of crayfish immediately downstream of mining sites. We conducted an insitu 28-d exposure to assess toxicity of mining-derived metals to the woodland crayfish (*Orconectes hylas*). Crayfish survival and biomass were significantly lower at mining sites than at reference and downstream sites. Metal concentrations in water, detritus, macroinvertebrates, fish, and crayfish were regulits support previous field and laboratory studies that showed mining-derived metals negatively affect *O. hylas* populations in streams draining the Viburnum Trend, and that in-situ toxicity testing was a valuable tool for assessing the impacts of mining on crayfish populations.

Published by Elsevier Inc.

#### 1. Introduction

Extensive deposits of lead (Pb) ore have been mined in Missouri for more than three centuries. Mining of the Viburnum Trend in southeast Missouri, USA, which contains economically significant Pb-zinc (Zn) ores, began in the 1950s. By 1970, the Viburnum Trend was the largest Pb-producing region in the world (Ryck and Whitley, 1974). Deposits within the Viburnum Trend also contain considerable quantities of copper-(Cu), cobalt-(Co), and nickel-(Ni) bearing minerals (Jessey, 1981). Although the mining district remains a major producer of Pb, and a minor producer of Zn and Cu (Missouri Department of Natural Resources, 2004), there is no commercial recovery of Co or Ni (Kuck, 2004; Shedd, 2004). We conducted no analyses of copper because previous studies have indicated that it is a minor component of metal contamination from mining in the Viburnum Trend (Besser et al., 2008; Brumbaugh et al., 2007).

Although modern mining practices incorporate efficient extraction technologies and operate within environmental regulations, early studies (Duchrow, 1983; Ryck and Whitley, 1974;

Wixson, 1977) showed that metal exposures to aquatic biota had occurred within the Viburnum Trend. Subsequent investigations have documented elevated metal concentrations in water, stream sediments, and aquatic food chains (Besser et al., 2006; Brumbaugh et al., 2007; Schmitt et al., 2007a,b), and the loss of biota including crayfish (Allert et al., 2008) and other macroinvertebrates (unpublished data; B. Poulton, USGS, Columbia, MO). In addition, sportfish including smallmouth bass (*Micropterus dolomieu*), longear sunfish (*Lepomis megalotis*), and suckers (Catostomidae) are known to contain Pb at concentrations exceeding recommended human food consumption levels (Missouri Department of Health and Senior Services, 2005).

Crayfish hold an intermediate trophic position and facilitate the flow of nutrients and energy in aquatic ecosystems (Lodge et al., 1995). Crayfish process a significant portion of organic matter in streams (Usio, 2000), breaking allocthonous and autochthonous material into smaller particles that are ultimately consumed by aquatic insects, snails, and microbial fauna (Huryn and Wallace, 1987; Parkyn et al., 2001). Crayfish are the dominant invertebrate biomass in some Ozark streams (Rabeni et al., 1995) and are the primary food source for several centrarchid fishes (Whitledge and Rabeni, 1997). Crayfish are also prey for more than 200 species of insects, arachnids, amphibians, fish, reptiles, birds, and mammals in and around North American water bodies (DiStefano, 2005).

<sup>&</sup>lt;sup>a</sup> US Geological Survey, Columbia Environmental Research Center (USGS/CERC), 4200 New Haven Road, Columbia, MO 65201, USA

<sup>&</sup>lt;sup>b</sup> Missouri Department of Conservation, 1110 South College Avenue, Columbia, MO 65201, USA

<sup>\*</sup> Corresponding author. Fax: +1573 876 1896. E-mail address: aallert@usgs.gov (A.L. Allert).

The southern-most mines within the Viburnum Trend are located in the headwaters of the Black River watershed (Fig. 1). The woodland crayfish (*Orconectes hylas*) is endemic to the Black River watershed (Pflieger, 1996) and occurs at high densities (25–37/m²) in riffle habitats of Ozark streams (DiStefano et al., 2002). Allert et al. (2008) reported reduced densities of *O. hylas* (0–2/m²) at sites immediately downstream of mining activities compared with densities (15–20/m²) at sites upstream of mining or at sites greater than 10 km downstream of mining areas in the Black River watershed. These findings suggest that mining activities are adversely affecting crayfish populations. We therefore conducted chronic in-situ exposures of *O. hylas* in two tributaries of the Black River watershed with the following objectives: (1) to evaluate crayfish responses relative to metal

exposure through aqueous and dietary pathways, and (2) to evaluate the effects of mining-derived metals on survival and growth of *O. hylas*.

#### 2. Methods

#### 2.1. Study area

Crayfish cages were deployed from 30 June to 28 July 2005 at seven sites in three tributaries of the Black River (Table 1). Previously collected physical, chemical and biological data (Allert et al., 2008; Besser et al., 2006; Brumbaugh et al., 2007) were used to identify reference sites (sites upstream of known mining activities); mining sites (sites impacted by mining activity); and downstream sites, where possible biological recovery may occur. Two sites were designated as




Fig. 1. Map of study sites in the Black River watershed of Missouri, USA.

Table 1
Sampling sites and cage locations in three tributaries of the Black River.

Site	Stream	Latitude, longitude <sup>a</sup>	Prominent mining feature upstream	Stream order	Stream distance from tailings or mine (km)	Type of site
SC2	Strother Creek	37°36′07.2″, 91°01′40.8″	Effluent pond; Buick tailings	2	3.7	M
WF1	West Fork	37°30′39.6″, 91°09′43.2″	None	2	NA	R
WF3	West Fork	37°29′49.2″, 91°05′13.2″	Brushy Creek tailings; West Fork tailings	2	2.2	M
WF4	West Fork	37°29′13.2″, 91°03′03.6″	Brushy Creek tailings; West Fork tailings	2	10.1	D
BF1	Bee Fork	37°26′43.4″, 91°06′29.4″	None	1	NA	R
BF3	Bee Fork	37°26′28.7″, 91°05′38.0″	Fletcher clarification dam and tailings	1	0.4	M
BF5	Bee Fork	37°27′36.0″, 91°01′30.0″	Fletcher clarification dam and tailings	2	7.4	D

Latitude, longitude as determined by global positioning system (GPS) receiver. R = reference site; M = mining site; D = downstream site; NA = not applicable.

a (GPS; ±10 m) based on the WGS84 geodetic datum.




Fig. 2. Diagram of cage. Cage frame consists of low-density polyethylene (LDPE) strips (0.635-cm width) heat welded to stainless-steel-woven-wire mesh (2.7-mm diagonal mesh opening). Cage includes a hinged door with 6-mm nylon shock cord used to secure door by fastening around stainless steel machine screws installed on cage frame and cage door. The feeding/stocking port consists of a twist-off polyvinyl chloride (PVC) cap. Polyethylene mesh packs (1.27-cm diagonal mesh opening) filled with rolled up scour pads (for crayfish refuge), organic material, and cobble were inserted into cage prior to stocking with crayfish and secured to bottom of cage with stainless-steel wire.

reference sites (WF1, BF1); three as mining sites (SC2, WF3, BF3); and two as downstream sites (WF4, BF5). Sites classified as mining sites were 0.4–3.7 km downstream of mining activity, whereas those classified as downstream sites were at least 7 km downstream of mining-related activity.

#### 2.2. Crayfish collection and rearing

Ovigerous *O. hylas* females were collected from the reference site on the West Fork (WF1) in early April in 2005 and returned to the US Geological Survey's Columbia Environmental Research Center (CERC) located in Columbia, MO, USA. Sixteen females were held in individual 2-L flow-through aquaria with CERC well water (temperature 18 °C, pH 7.7, alkalinity 254 mg/L as CaCO<sub>3</sub>, hardness 286 mg/L as CaCO<sub>3</sub>) and fed frozen brine shrimp (San Francisco Bay Brand, Inc., San Francisco, CA, USA) *ad libitum* daily. Upon hatching and detaching from the adult females, juvenile crayfish were placed in a fiberglass tank filled with CERC well water and fed flake food (Ziegler Brothers, Inc., Gardner, PA, USA) *ad libitum* daily until their body width was greater than 2 mm. Prior to stocking crayfish into cages, a subset of crayfish available for stocking into cages (n=88) was measured for mean carapace length (CL, from tip of rostrum to posterior edge of the cephalothorax, to the nearest 0.1 mm) and weighed (to the nearest 0.01 g).

#### 2.3. Toxicity test

A 28-d in-situ toxicity test was conducted with juvenile O. hylas, at seven sites in the West Fork, Bee Fork, and Strother Creek of the Black River watershed (Fig. 1). Crayfish were exposed in hemicylindrical (0.28-m<sup>2</sup>) cages constructed of stainlesssteel wire mesh (2.7-mm diagonal opening) and polyethylene (LDPE) reinforcing strips (Fig. 2). We collected cobble substrates (2.5-7.5 cm) and approximately 10-g organic material (henceforth detritus) from each site to provide both food and shelter for caged crayfish. Cobble, detritus, and three polyethylene scour pads were placed in three polyethylene-mesh packs (15-cm length  $\times$  30-cm width; 1.27-cm diagonal opening), which were closed using plastic cable ties and secured to the bottom of each cage using stainless-steel wire. Prior to placing organic material into the mesh packs, all predatory insects (i.e., Odonata and Plecoptera larvae) were removed. The bottom of the cages were buried 2-4-cm into the stream sediment to expose cravfish to sub-surface water and to anchor the cages. Minced fish (largescale stonerollers, Campostoma oligolepis; henceforth stonerollers) from each site were added to each cage weekly in increasing increments to maintain dietary rations proportional to anticipated crayfish biomass (0.1, 0.1, 0.2, 0.4 g minced fish in weeks 1, 2, 3, and 4, respectively). Ten juvenile crayfish were placed in each of the six cages at each site except at SC2, where seven cages were deployed. Three cages (n = 30 crayfish) were sampled on both day 14 and 28,

except at SC2, where four cages were sampled on day 28. Crayfish were measured (CL and wet weight) and frozen for metal analyses. Test endpoints included survival and growth. Biomass (i.e., standing crop) at day 28 was estimated by multiplying the number of survivors by the mean wet weight of survivors at each site.

#### 2.4. Wild crayfish collections

Wild crayfish ( $O.\ hylas$ ) were collected on days 0 and 28 of the toxicity test at each site to compare CL, wet weight, and metal concentrations of wild crayfish to those stocked in cages. Crayfish were collected by disturbing substrate directly upstream of a small kick seine (1-m length  $\times$  1.5-m height) with 3-mm delta mesh (Flinders and Magoulick, 2005). Wild crayfish were collected within 200 m of cage placement. Collections were made upstream or downstream of cage placement, depending on the nearest available riffle, and number of kick seines required to collect 30 individuals. All crayfish collected were identified to species, measured (CL and wet weight) and frozen for metal analyses.

#### 2.5. Water samples

Surface water quality parameters (temperature, pH, conductivity, dissolved oxygen, and turbidity) were measured in-situ on days 0, 7, 14, 21, and 28 using a Hydrolab<sup>®</sup>. Quanta meter (Loveland, CO, USA). A sub-surface grab sample was collected in a pre-cleaned 4-L carboy at each site on days 0, 14, and 28 for additional water quality and nutrient analyses at CERC. Alkalinity and hardness were measured by titration (APHA, 2005). Sulfate was measured by colorimetric detection with a Hach<sup>®</sup> 2100 spectrophotometer (Loveland, CO, USA). Recovery of reference standards used as laboratory control samples for surface water quality parameters (pH, conductivity, dissolved oxygen, turbidity, alkalinity, hardness, and sulfate) ranged from 90% to 118%. Overall, detection and recoveries of water quality parameters were within acceptable criteria, thus none of the sample results were corrected.

Samples for dissolved nutrient analyses were filtered through 0.4-µm polycarbonate filters under vacuum pressure within 4 days of collection. Nutrients were measured in surface water samples with a Technicon® Autoanalyzer (Tarrytown, NY, USA) using colorimetric detection (APHA, 2005). Total ammonia (NH<sub>3</sub>) was analyzed using a salicylate/nitroprusside colorimetric reaction. Dissolved nitrite/nitrate (NO<sub>2</sub>/NO<sub>3</sub>) was measured following cadmium reduction and measured using colorimetric reaction (APHA, 2005). Soluble reactive phosphorus (SRP) was determined using the automated ascorbic acid method (APHA, 2005). Samples for total phosphorous (TP) and total nitrogen (TN) were digested in sodium hydroxide and potassium persulfate then analyzed using the automated ascorbic acid and the automated cadmium reduction methods, respectively (APHA, 2005). Dissolved organic carbon (DOC) was analyzed using a persulfate/UV digestion followed by colorimetric analysis of CO<sub>2</sub>. Method limits of detection limits (MLDs) for nutrients are listed in Table S-1. Recovery of reference standards used as laboratory control samples for nutrients ranged from 100% to 128%, except for one standard for NH<sub>3</sub> (1 mg N/L) that was 145%. Overall, detection and recoveries of nutrients were within acceptable criteria and all measured concentrations exceeded MLDs, thus none of the sample results were corrected for recovery.

#### 2.6. Metal concentrations

Water samples were filtered for metal analyses using a polypropylene syringe and filter cartridge (0.45- $\mu$ m pore size) into a pre-cleaned polyethylene bottle on-site, and placed on ice. Water samples were subsequently acidified to 1% (v/v) with nitric acid (J.T. Baker Inc., Phillipsburg, NJ, USA) within 4 days of collection.

Surface water samples were analyzed for Pb, Zn, Cd, Ni, and Co by inductively-coupled plasma-mass spectrometry (ICP-MS) (Brumbaugh et al., 2007; May et al., 1997). Calibration verification, method limits of detection, and recoveries of metals in reference solutions, duplicates, and spikes were all within acceptable criteria and all measured concentrations exceeded MLDs (Brumbaugh et al., 2007).

Detritus, macroinvertebrates (i.e., Ephemeroptera, Odonata, Plecoptera, Megaloptera, Trichoptera, Diptera, and Chirononmidae), stonerollers, and whole crayfish from each site were analyzed for Pb, Zn, Cd, Ni, and Co by ICP-MS (Besser et al., 2006; Brumbaugh et al., 2005). Tissues were lyophilized and reduced to a coarse powder by mechanical crushing in a glass vial with a glass rod. Neither exoskeletons nor gut contents of any of the biota were removed before analysis. A dry mass of 0.25 g from each composited sample was digested using concentrated nitric acid and microwave heating. Quality control measures incorporated at the digestion stage included digestion blanks, certified reference materials, replicates, and spikes. A calibration blank and an independent calibration verification standard were analyzed with every 10 samples to confirm the calibration status of the ICP-MS during instrumental analyses of digestates. The MLDs for detritus,

macroinvertebrates, stonerollers, and crayfish are listed in Table S-2. All measured concentrations exceeded the MLDs.

Recoveries of the elements from reference materials (fish, mussel, oyster, plant, and plankton) ranged from 87% to 115%. Relative percent differences (RPDs) for replicate analyses were < 26% for all elements except for Pb analyses of one sample of detritus (50%) and one sample of macroinvertebrates (85%). Instrumental precision, estimated by determining the RPDs from the duplicate analysis of detritus and biota digestates, was < 4%. Recoveries of method spikes for all five metals in 14 separate spiked samples of all the sample types analyzed averaged 97%. Post-digestion or analysis spike recoveries ranged from 83% to 104%. As a check for potential interferences, dilution percent differences (DPDs) based on 5X dilutions of detritus and biota sample digestates were determined; DPDs were < 10% for all metals. Blank-equivalent concentrations (BECs) for digestion blanks were less than corresponding MLDs; therefore sample results were not corrected for BECs. Overall, quality assurance results indicated that the methods used provided acceptable accuracy and precision, thus none of the sample results were corrected.

#### 2.7. Statistical analysis

Statistical analyses were conducted using Statistical Analysis System (SAS) for Windows (Release 9.1; SAS Institute, Cary, NC, USA). Censored values (< MLD) were replaced with 50% of the MLD for statistical computations and graphing. Survival and biomass data of caged crayfish on day 28 of the toxicity test and the overall means (e.g., data from days 0, 7, 14, 21, and 28) for water quality, nutrients, and metal concentrations were used in the statistical analyses. No wild crayfish were collected within 200 m of our cages at SC2, so our search was expanded until one adult wild crayfish was collected at a distance greater than 500 m away from the cages; therefore it was not used in the statistical analyses of CL and wet weight of wild crayfish. However, data from this individual is present in tables and figures for comparative purposes. All data were tested for normality and homogeneity of variance using the PROC UNIVARIATE module in SAS. Data were not normally distributed, therefore rank transformation were used in statistical analyses. Differences in caged crayfish survival and biomass among sites and groups of sites were tested using nested analysis-of-variance (ANOVA; cages nested within site), with site considered a fixed effect. Differences in caged crayfish survival among groups of sites were tested as planned non-orthogonal contrasts using single degree-of-freedom F-tests. The mean square for cage survival within site was used in all tests, which were conducted using the PROC GLM module in SAS. Differences in crayfish survival and biomass among individual sites were also evaluated with Duncan's multiple range test. Differences in wild crayfish growth, water quality, nutrient, and metal concentrations among groups of sites were also tested using the same procedures. Differences in mortality, size of crayfish, water quality, nutrient, and metal concentrations were tested on a stream-by-stream basis using the Kruskal-Wallis test. Results were similar to those that included all seven sites; therefore we list only exceptional results in Table S-3. Finally, associations among caged crayfish survival and biomass on day 28, wild crayfish (excluding SC2) CL and wet weight, water quality, nutrient, and metal concentrations were examined with Spearman's correlation analysis. A significance level of P < 0.05 was used to judge all statistical tests.

#### 3. Results

#### 3.1. Toxicity test

Mean percent survival of caged crayfish on day 28 (henceforth survival) was significantly greater at reference sites (90%) than at mining sites (39%; Table 2; Fig. 3), and decreased sharply with increasing metal concentrations in surface water (Fig. 4). When analyzed on a stream-by-stream basis, survival at the mining site (BF3) in the Bee Fork was significantly lower; however, survival was not significantly lower at the mining site (WF3) in the West Fork (Table S-3).

Mean CL and wet weight of caged crayfish at all sites on day 28 were significantly greater than those on day 0 (n=88; CL =  $6.7\pm0.11$ ; wet wt. =  $0.06\pm0.003$ ), except mean wet weight of crayfish at SC2 (Table 2). Mean CL and wet weight of crayfish at day 28 were significantly greater at downstream sites than at reference or mining sites; however, CL and weight were not significantly different among mining and reference sites (Table 2). Mean biomass of crayfish at reference and downstream sites on day 28 was significantly greater than at mining sites (Table 2; Fig. 3).

**Table 2**Number (n) of living cravfish, percent (%) survival, carapace length (CL), and wet weight (means with standard error in parenthesis) of caged *Orconectes hylos*.

Day/site	п	% Survival	CL (mm)	Wet wt. (g)	Biomass (g/m²)
Day 14 Reference sites WF1 <sup>a</sup> BF1 Group mean	24 28	83 (3) b 93 (7) a 88 (4) A	9.3 (0.19) b 8.5 (0.22) c 8.9 (0.16) C	0.17 (0.01) c 0.13 (0.01) d 0.15 (0.01) B	4.3 (0.7) ab 5.3 (0.8) ab 4.7 (0.5) A
Mining sites SC2 WF3 <sup>b</sup> BF3 Group mean	23 27 17	84 (9) b 90 (6) a 57 (23) b 77 (27) A	8.4 (0.24) c 9.7 (0.21) b 9.8 (0.31) b 9.3 (0.16) A	0.12 (0.01) d 0.20 (0.01) bc 0.18 (0.02) c 0.17 (0.01) B	3.3 (0.4) b 5.3 (0.6) ab 4.0 (1.8) ab 4.3 (0.6) B
Downstream sites WF4 BF5 Group mean	28 29	84 (3) b 97 (3) a 92 (8) A	11.2 (0.17) a 9.8 (0.23) b 10.5 (0.01) B	0.32 (0.02) a 0.23 (0.02) b 0.27 (0.01) A	8.3 (1.8) a 6.0 (1.8) ab 7.3 (1.3) A
<b>Day 28</b> Reference sites WF1 BF1 Group mean	27 26	90 (6) a 90 (0) a 90 (3) A	10.5 (0.33) bc 10.8 (0.37) b 10.7 (0.15) B	0.40 (0.02) cd 0.29 (0.03) de 0.33 (0.14) B	1.2 (3.3) c 14.7 (2.3) bc 13.3 (1.9) B
Mining sites SC2 <sup>c</sup> WF3 BF3 Group mean	3 20 16	7 (3) d 67 (9) bc 53 (3) c 39 (9) B	9.2 (0.03) c 11.3 (0.47) b 13.2 (0.46) a 11.9 (0.36) B	0.17 (0.03) e 0.37 (0.05) cd 0.58 (0.05) ab 0.44 (0.04) B	0.3 (0.2) d 7.7 (2.5) cd 7.7 (3.5) cd 4.7 (1.6) C
Downstream sites WF4 BF5 Group mean	26 25	87 (3) ab 77 (13) ab 82 (16) A	14.0 (0.31) a 12.9 (0.3) a 13.4 (1.6) A	0.71 (0.05) a 0.53 (0.04) bc 0.61 (0.03) A	32.0 (4.9) a 22.7 (5.2) ab 27.3 (3.8) A

Sites with the same lower case letter and groups of sites with the same capital letter are not significantly different for each test day (P < 0.05). Initial stocking number = 30, unless otherwise noted.

#### 3.2. Wild crayfish collections

Very few wild crayfish were collected at any of the mining sites on either day 0 or 28, despite significant sampling effort (i.e., >15 kick seines). Mean wet weight of wild crayfish on days 0 and 28 were significantly greater at reference and downstream sites than mining sites; however, there was no significant difference in CL among groups of sites on day 28 (Table 3).

#### 3.3. Water quality

Mean conductivity, hardness, and sulfate concentrations in surface water were significantly higher at mining and downstream sites than reference sites (Table 4); whereas alkalinity was significantly lower at downstream sites than at reference and mining sites (Table 4). Reference sites had significantly higher turbidity (Table 4); however, values were still very low. Dissolved oxygen and pH at reference sites were significantly lower than mining and downstream sites (Table 4); however, at all the sites, the range in dissolved oxygen and pH readings were narrow.

Ammonia,  $NO_2/NO_3$ , and TN concentrations in surface water from mining sites were significantly higher than reference and downstream sites (Table 5). Soluble reactive phosphorous concentrations in surface waters from mining and downstream sites were significantly lower than reference sites (Table 5); however,

there were no significant differences in TP concentrations among reference, mining and downstream sites (Table 5). Dissolved organic carbon was significantly lower at downstream sites than reference or mining sites (Table 5).

Analyses on a stream-by-stream basis produced slightly different results for several parameters. There were no significant differences in dissolved oxygen or DOC at sites in the West Fork; however, there were at sites in the Bee Fork (Table S-3). There were significant differences in alkalinity at sites in the West Fork; however, alkalinity was not significantly different at sites in the Bee Fork. There was no significant difference in SRP or NH<sub>3</sub> in either the Bee Fork or West Fork, despite an overall significant difference when SC2 was included in the analyses.

#### 3.4. Metal concentrations


Concentrations of metals in surface water were all below surface water chronic criteria (WQC; USEPA, 2002; Table 6). Cobalt concentrations in surface water were also all below a proposed Canadian guideline for chronic exposure of  $4\,\mu g\, Co/L$  (Nagpal, 2004). However, concentrations of Pb, Zn, Ni, and Co in surface waters were significantly higher at mining sites compared with reference or downstream sites (Table 6).

Concentrations of all metals in detritus, macroinvertebrates, stonerollers, and caged crayfish were significantly higher at

<sup>&</sup>lt;sup>a</sup> Stocking number = 29.

<sup>&</sup>lt;sup>b</sup> Stocking number = 31.


<sup>&</sup>lt;sup>c</sup> Stocking number = 40.



**Fig. 3.** (a) Mean percent (%) survival and (b) mean biomass of caged *Orconectes hylas* within groups of sites on days 14 and 28. Groups of sites with the same letter are not significantly different (P<0.05).

mining sites than at reference or downstream sites, with the exception of Cd concentrations in macroinvertebrates, and stonerollers (Table 7; Fig. 5). Mean concentration of Pb in wild crayfish was significantly higher at mining sites than at reference or downstream sites (Table 7). Mean concentrations of Zn and Co in all sample types were highest at SC2. Lead and Ni concentrations were highest at SC2 and/or BF3. Metal concentrations in samples at all mining sites were generally two to ten-fold higher in all sample types compared with those at downstream or reference sites. Concentrations of Pb, Zn, and Ni were highest in detritus and were generally higher in macroinvertebrates compared with stonerollers or crayfish. Metal concentrations in the West Fork were generally lower than those in the Bee Fork. Concentrations of Pb and Cd in fish; Zn in invertebrates, and Zn in surface water were not significantly different among the three groups of sites in the West Fork (Table S-3).

Mean concentrations of metals in crayfish at day 0 were: Pb  $(6.0\pm0.4\,\mu\text{g/g}\ \text{dry}\ \text{weight})$ , Zn  $(81\pm0.3\,\mu\text{g/g}\ \text{dry}\ \text{weight})$ , Cd  $(0.16\pm0.2\,\mu\text{g/g}\ \text{dry}\ \text{weight})$ , Ni  $(4.0\pm0.7\,\mu\text{g/g}\ \text{dry}\ \text{weight})$ , and Co  $(2.0\pm0.4\,\mu\text{g/g}\ \text{dry}\ \text{weight})$ . Caged crayfish rapidly accumulated all metals at all mining sites (Fig. 6). Metal concentrations in caged crayfish at SC2 continued to increase throughout the exposure; however, they did not at WF3 or BF3. Metal concentrations in caged crayfish on day 28 at reference and downstream sites were generally lower than concentrations in crayfish stocked into cages on day 0. Concentrations of metals in caged crayfish at day 28 were comparable to those in wild crayfish at all sites except at SC2, where metal concentrations in caged crayfish were greater than the single large wild crayfish collected at SC2.



**Fig. 4.** Relationship between mean percent (%) survival of caged *Orconectes hylas* on day 28 and mean metal concentrations in surface water at sampling sites: (a) lead; (b) zinc; (c) cadmium; (d) nickel; and (e) cobalt. Dashed lines are mean chronic water quality criteria for the sample sites.

#### 3.5. Correlation analyses

Carapace length and wet weight of caged crayfish were negatively correlated with Zn concentrations in stonerollers (Table S-4). Carapace length and wet weight of caged crayfish

**Table 3**Number (*n*), carapace length (CL), and wet weight (means with standard error in parenthesis) of wild *Orconectes hylas* collected at sampling sites on days 0 and 28 of the in-situ exposure.

Day/site	n	CL (mm)	Wet wt. (g)
Day 0 Reference sites WF1 BF1 Group mean	30 35	10.1 (0.3) a 9.5 (0.2) a 9.8 (0.3) A	0.26 (0.03) a 0.21 (0.02) a 0.23 (0.02) A
Mining sites SC2 WF3 BF3 Group mean	0 1 32	- 5.4 (-) a 9.2 (0.3) a 7.3 (1.9) B	- 0.03 (-) a 0.18 (0.01) a 0.11 (0.08) B
Downstream sites WF4 BF5	30 43	9.4 (0.3) a 7.1 (0.2) a 8.2 (0.1) A	0.22 (0.03) a 0.08 (0.01) a 0.15 (0.07) A
Day 28 Reference sites WF1 BF1 Group mean	30 37	12.9 (0.3) a 12.0 (0.4) a 12.5 (0.4) A	0.58 (0.04) a 0.47 (0.04) a 0.53 (0.05) A
Mining sites SC2 WF3 BF3 Group mean	1 12 2	35.0 (-) 12.2 (0.4) a 12.3 (0.2) b 12.2 (0.02) A	12.1 (-) 0.48 (0.05) a 0.34 (0.05) a 0.41 (0.07) B
Downstream sites WF4 BF5 Group mean	18 30	13.2 (0.3) a 12.3 (0.3) a 12.7 (0.5) A	0.56 (0.04) a 0.53 (0.05) a 0.55 (0.02) A

Sites with the same lower case letter and groups of sites with the same capital letter are not significantly different for each test day (P<0.05).

were negatively correlated with SRP, and CL was negatively correlated with TP (Table S-5).

Mean percent survival of caged crayfish was negatively correlated with most metal concentrations in surface water, detritus, macroinvertebrates, stonerollers, and caged crayfish (Table 8). Biomass of caged crayfish was negatively correlated with Zn concentrations in stonerollers and Co concentrations in caged crayfish. Concentrations of Pb, Zn, Ni, and Co in caged crayfish and all environmental matrixes were highly inter-correlated (Table 8). Lead concentrations in caged crayfish were positively correlated with Pb, Zn, and Ni concentrations in wild crayfish (Table 8). Zinc concentrations in caged crayfish were positively correlated with Co concentrations in wild crayfish. Survival of caged crayfish was negatively correlated with NH<sub>3</sub>, NO<sub>2</sub>/NO<sub>3</sub>, TN, conductivity, hardness, and sulfate (Table S-5). Biomass of caged crayfish was negatively correlated with DOC (Table S-5).

Wet weight of wild crayfish was correlated with Co concentrations in wild crayfish (Table S-4); however, CL and wet weight of wild crayfish were not significantly correlated with any water quality parameter or nutrient concentration (Table S-5). There were fewer significant correlations between metal concentrations in wild crayfish and the other samples collected at the sites; however, concentrations of Pb in wild crayfish were significantly correlated with most metals in all sample types analyzed (Table 8). Zinc concentrations in wild crayfish were significantly correlated with Pb, Zn, and Cd concentrations in detritus, and with Zn concentrations in macroinvertebrates. Nickel concentrations in wild crayfish were correlated with Zn concentrations in macroinvertebrates, and Pb, Zn, and Cd concentrations in detritus.

Ammonia, NO<sub>2</sub>/NO<sub>3</sub>, TN, conductivity, hardness, sulfate, and Ni concentrations in surface water were all significantly correlated (Table S-6). Soluble reactive phosphorous (SRP) was negatively correlated with dissolved oxygen; dissolved organic carbon was negatively correlated with pH; and temperature was negatively correlated with Co concentrations in surface water. Hardness was also significantly correlated with Pb, Ni, and Co concentrations in surface water were significantly correlated with Co concentrations in surface water, as were Ni and Co concentrations in surface water.

#### 4. Discussion

We documented decrease survival of caged crayfish at sites directly downstream (0.4–3.7 km) of mining sites. Survival and

**Table 4**Water quality (means with standard error in parenthesis) of surface water at sampling sites.

Site	Temp (°C)	PH (SU)	Cond (µs/cm)	DO (mg/L)	Alk (mg/L as CaCO <sub>3</sub> )	Hard (mg/L as CaCO <sub>3</sub> )	Turb (NTU)	Sulfate (mg/L)
Reference sites WF1 BF1	23.6 (0.6) a 23.1 (1.4) b	8.04 (0.03) ab 7.84 (0.09) c	344 (5) e 282 (6) f	7.8 (0.5) a 6.3 (0.4) b	182 (3) a 143 (3) c	183 (3) d 145 (2) e	0.60 (0.04) a 0.50 (0.04) a	0.3 (0.2) e 1 (0.3) e
Group mean	23.4 (0.7) A	7.94 (0.06) B	313 (11) C	7.1 (0.4) B	162 (8) A	164 (7) C	0.55 (0.03) B	0.6 (0.2) C
Mining sites SC2 WF3 BF3 Group mean	25.8 (0.5) a 24.6 (0.9) a 24.2 (0.9) a 24.9 (0.5) A	7.94 (0.04) bc 8.09 (0.03) a 8.01 (0.05) ab 8.01 (0.03) A	858 (31) a 418 (7) cd 592 (15) b 623 (42) A	8.0 (0.2) a 8.4 (0.3) a 8.4 (0.1) a 8.3 (0.1) A	117 (3) d 170 (3) b 143 (2) c 143 (7) A	409 (11) a 212 (3) c 247 (5) b 289 (26) A	0.50 (0.04) a 0.40 (0.05) a 0.40 (0.06) a 0.45 (0.03) A	304 (10) a 53 (3) d 129 (6) b 162 (32) A
Downstream sit WF4 BF5 Group mean	24.3 (0.4) a 22.9 (0.8) a 23.6 (0.5) A	8.10 (0.04) a 8.05 (0.04) ab 8.08 (0.03) A	395 (2) d 446 (4) c 420 (9) B	8.3 (0.1) a 8.2 (0.1) a 8.3 (0.1) A	165 (2) b 141 (1) c 152 (5) B	198 (1) c 199 (1) c 199 (1) B	0.50 (0.12) a 0.40 (0.02) a 0.44 (0.05) A	42 (1) d 80 (1) c 64 (8) B

Temp = temperature; Cond = conductivity; DO = dissolved oxygen; Alk = alkalinity; Hard = hardness; Turb = turbidity. Sites with the same lower case letter and groups of sites with the same capital letter are not significantly different (P<0.05).

**Table 5**Nutrient concentrations (means with standard error in parenthesis) of surface waters at sampling sites.

Site	NH <sub>3</sub> (mg N/L)	SRP (μg P/L)	NO <sub>2</sub> /NO <sub>3</sub> (mg N/L)	DOC (mg C/L)	TN (mg N/L)	TP (μg P/L)
Reference sites WF1 BF1	0.01 (0.002) b 0.01 (0.002) b	0.5 (0.3) a 0.8 (0.4) a	0.04 (0.01) d 0.07 (0.01) d	0.59 (0.04) bc 0.90 (0.07) a	0.10 (0.01) c 0.20 (0.01) c	3.7 (0.6) ab 2.4 (0.5) ab
Group mean	0.01 (0.002) B	0.6 (0.3) A	0.06 (0.01) C	0.75 (0.06) A	0.12 (0.01) B	2.9 (0.4) A
Mining sites SC2 WF3 BF3 Group mean	0.06 (0.01) a 0.01 (0.002) b 0.01 (0.003) b 0.03 (0.01) A	0.3 (0.2) a 0.2 (0.1) a 0 c 0.1 (0.1) B	1.2 (0.09) a 0.13 (0.01) cd 0.51 (0.02) b 0.61 (0.10) A	0.91 (0.06) a 0.60 (0.02) bc 0.69 (0.06) b 0.73 (0.04) A	1.30 (0.08) a 0.20 (0.02) c 0.50 (0.03) b 0.68 (0.10) A	4.4 (0.4) a 2.8 (0.8) b 2.0 (0.6) a 3.1 (0.4) A
Downstream sites WF4 BF5 Group mean	0.01 (0.003) b 0.01 (0.003) b 0.01 (0.001) B	0 a 0.2 (0.2) a 0.1 (0.1) B	0.08 (0.01) d 0.19 (0.01) c 0.14 (0.02) B	0.57 (0.03) c 0.50 (0.04) c 0.53 (0.03) B	0.10 (0.01) c 0.20 (0.01) c 0.17 (0.02) B	2.0 (0.2) b 2.6 (1.2) ab 3.1 (0.4) A

 $NH_3$  = ammonia; SRP = soluble reactive phosphorus;  $NO_2/NO_3$  = nitrite/nitrate; DOC = dissolved organic carbon; TN = total nitrogen; TP = total phosphorous. Sites with the same lower case letter and groups of sites with the same capital letter are not significantly different (P<0.05).

Table 6 Concentrations ( $\mu$ g/L; means with standard error in parenthesis) of metals measured in surface waters (WC) at sampling sites, and site hardness-adjusted chronic water quality criteria (WQC).

Site	Lead		Zinc	Zinc		Cadmium		Nickel	
	WC	WQC	WC	WQC	WC	WQC	WC	WQC	WC
Reference sites									
WF1	0.03 (0) c	4.9	4.1 (2) b	197	0.03 (0) b	0.37	0.13 (0) c	46	0.09 (0.01) de
BF1	0.03 (0) c	3.8	2.5 (0) b	162	0.03 (0) b	0.32	0.13 (0) c	38	0.14 (0.01) d
Group mean	0.03 (0) B	4.4	3.3 (1) B	180	0.03 (0) A	0.35	0.13 (0) B	42	0.12 (0.01) B
Mining sites									
SC2	0.71 (0.02) b	11.3	56 (5) a	390	0.06 (0.02) a	0.65	26 (2) a	90	0.64 (0.05) a
WF3	0.12 (0.01) c	5.7	2.5 (0) b	223	0.03 (0) b	0.41	1.9 (0.01) c	52	0.39 (0.02) b
BF3	1.59 (0.06) a	6.8	6.6 (0.2) b	254	0.03 (0) b	0.46	6.0 (0.01) b	59	0.28 (0.01) c
Group mean	0.86 (0.18) A	7.9	20 (7) A	289	0.03 (0.01) A	0.51	11 (3) A	67	0.43 (0.05) A
Downstream sites									
WF4	0.04 (0.01) c	5.3	2.5 (0) b	211	0.03 (0) b	0.40	0.40 (0) c	49	0.15 (0.01) d
BF5	0.03 (0) c	5.3	3.9 (1) b	211	0.03 (0) b	0.40	0.50 (0) c	49	0.12 (0.01) d
Group mean	0.04 (0.01) B	5.3	3.3 (1) B	211	0.03 (0) A	0.40	0.46 (0.02) B	49	0.13 (0.01) B

Sites with the same lower case letter and groups of sites with the same capital letter are not significantly different (P<0.05).

biomass of caged crayfish were significantly lower at mining sites than reference or downstream sites, and survival was negatively correlated with metal concentrations in surface water, detritus, macroinvertebrates, stonerollers, and whole crayfish. Survival was also negatively correlated with several water quality parameters. Our study supports previous results which found reduced densities of O. hylas populations in riffle habitats, and elevated metals concentrations (Pb, Zn, Cd, Ni, and Co) in O. hylas at sites directly downstream of mining in the Black River watershed (Allert et al., 2008). Metal concentrations in detritus and biota at mining sites in this study were comparable to those of Besser et al. (2006), and decreased with distance from mining sites. Alikhan et al. (1990) also reported an inverse relationship between distance from a contaminant source and metal concentrations in crayfish. Conductivity, hardness, sulfate, and nitrogen ions often are elevated below mining sites (Gray, 1998; Tiwary, 2001), and may be useful in indicating the extent of mining impacts. Finally, in-situ testing of crayfish was an important tool for demonstrating that absence of crayfish populations below mining sites was the result of metal exposure as opposed to habitat loss due to physical impairment by mine waste (e.g., sedimentation by mine tailings).

Although no toxicity studies have been conducted with juvenile O. hylas in single-metal exposures, previous studies have shown several species of crayfish to be relatively sensitive to metals, and that juvenile crayfish are more sensitive than adult crayfish. Wigginton and Birge (2007) calculated 96-h median lethal concentrations ( $LC_{50}$ ) of Cd to six species of crayfish (Cambaridae) with a mean  $LC_{50}$  of  $1510\,\mu g\,Cd/L$  for adult crayfish and of  $111\,\mu g\,Cd/L$  for juvenile crayfish at a water hardness of approximately  $45\,m g/L$  as  $CaCO_3$ . Other studies (Lindhjem and Bennet-Chambers, 2002; Mirenda, 1986a, b; Naqvi and Howell, 1993) have also reported lethal concentrations that are greater than the metal concentrations measured in surface waters from our study streams; however, these laboratory studies were short in duration and conducted with different species and sizes of crayfish. Thorp et al. (1979) reported that long-term exposure to 5

 $\begin{tabular}{ll} \textbf{Table 7} \\ \textbf{Metal concentrations ($\mu g/g$ dry weight; means with standard error in parenthesis) in detritus and biota at sampling sites. \\ \end{tabular}$ 

Metal and site(s)	Detritus <sup>a</sup>	Macro-invertebrates <sup>a</sup>	Stonerollers <sup>b</sup>	Wild crayfish <sup>c</sup>	Caged crayfish <sup>d</sup>
<b>Lead</b> Reference sites WF1 BF1 Group mean	11 (2) b 7.2 (1) b 8.9 (1) B	2.0 (1) c 1.0 (0.2) c 1.5 (0.5) B	2.6 (0.1) d 4.3 (1) cd 3.4 (1) B	0.53 (-) 0.42 (-) 0.48 (0.1) B	1.2 (0.2) b 0.99 (0.2) b 1.1 (0.2) B
Mining sites SC2 WF3 BF3 Group mean	650 (82) a 143 (30) b 567 (63) a 453 (74) A	81 (12) b 11 (2) c 114 (15) a 69 (14) A	42 (2) a 8.9° (-) c 33 (3) b 32 (6) A	14 (-) 1.7 (-) 42 (-) 15 (9) A	37 <sup>e</sup> (7) a 6.4 (1) b 37 (4) a 26 (4) A
Downstream sites WF4 BF5 Group mean	54 (8) b 32 (2) b 41 (5) B	5.3 (1) c 8.1 (1) c 6.9 (1) B	4.6 (1) cd 6.2 (1) cd 5.6 (1) B	1.5 (-) 2.0 (-) 1.7 (0.2) B	2.8 (0.3) b 2.6 (0.3) b 2.7 (0.2) B
<b>Zinc</b> Reference sites WF1 BF1 Group mean	26 (3) d 25 (1) d 25 (1) B	100 (9) c 90 (9) c 95 (6) B	112 (4) b 129 (3) b 120 (5) B	62 (-) 57 (-) 60 (3) A	64 (1) d 68 (1) cd 66 (1) B
Mining sites SC2 WF3 BF3 Group mean	2288 (309) a 529 (27) bc 693 (15) b 1170 (257) A	392 (35) a 162 (41) bc 196 (28) b 250 (36) A	317 (31) a 148 <sup>c</sup> (-) b 108 (15) b 200 (50) A	73 64 83 71 (5) A	236° (10) a 105 (5) b 97 (3) b 146 (16) A
Downstream sites WF4 BF5 Group mean	234 (11) cd 129 (5) d 174 (22) B	151 (44) bc 146 (14) bc 148 (18) B	93 (7) b 102 (13) b 98 (8) B	79 (-) 65 (-) 72 (7) A	68 (1) cd 68 (3) cd 68 (1) B
<b>Cadmium</b> Reference sites WF1 BF1 Group mean	0.32 (0.02) b 0.27 (0.01) b 0.30 (0.02) B	0.81 (0.3) bc 0.72 (0.1) cd 0.77 (0.5) A	0.1 (0.02) de 0.3 (0.01) a 0.2 (0.04) A	0.75 (-) 0.77 (-) 0.76 (0.01) A	0.60 (0.04) bc 0.56 (0.03) bc 0.57 (0.02) B
Cadmium (continued) Mining sites SC2 WF3 BF3 Group mean	3.6 (1) a 0.91 (0.1) b 1.3 (0.04) b 1.9 (0.4) A	1.2 (0.1) ab 0.32 (0.1) d 1.4 (0.3) a 0.97 (0.2) A	0.2 (0.01) ab 0.1° (-) de 0.2 (0.01) cd 0.2 (0.02) A	0.78 (-) 1.4 (-) 0.41 (-) 1.0 (0.25) A	2.1° (0.2) a 0.32 (0.02) cd 0.53 (0.1) bc 0.98 (0.2) A
Downstream sites WF4 BF5 Group mean	0.56 (0.1) b 0.53 (0.03) b 0.54 (0.03) B	0.41 (0.1) cd 1.2 (0.03) ab 0.90 (0.2) A	0.1 (0.01) e 0.2 (0.01) bc 0.2 (0.02) A	0.54 (-) 0.70 (-) 0.62 (0.08) A	0.25 (0.03) d 0.76 (0.1) b 0.50 (0.1) B
<b>Nickel</b> Reference sites WF1 BF1 Group mean	8.5 (1) d 12 (0.4) d 10 (1) B	2.3 (1) c 2.3 (0.4) c 2.3 (0.4) B	2.0 (0.02) de 1.2 (0.1) e 1.6 (0.24) B	1.1 (-) 1.1 (-) 1.1 (0.01) A	1.2 (0.04) c 1.5 (0.1) c 1.3 (0.1) B
Mining sites SC2 WF3 BF3 Group mean	487 (62) a 124 (7) c 298 (8) b 303 (49) A	57 (9) a 17 (4) bc 28 (6) b 34 (6) A	19 (0.2) a 9.5° (-) b 5 (0.3) c 12 (3) A	4.2 (-) 1.2 (-) 16 (-) 5.6 (3) A	17° (2) a 7.5 (0.4) b 8.3 (1) b 11 (1) A
Downstream sites WF4 BF5	34 (2) d 41 (2) d	4.0 (0.2) c 6.3 (1) c	2.1 (2) de 2.7 (0.5) d	2.1 (-) 2.0 (-)	2.1 (0.2) c 1.9 (0.1) c

Table 7 (continued)						
Metal and site(s)	Detritus <sup>a</sup>	Macro-invertebrates <sup>a</sup>	Stonerollers <sup>b</sup>	Wild crayfish <sup>c</sup>	Caged crayfish <sup>d</sup>	
Group mean	38 (2) B	5.3 (0.6) B	2.5 (0.3) B	2.1 (0.04) A	2.0 (0.1) B	
Cobalt Reference sites WF1 BF1 Group mean	8.7 (1) c 10 (1) c 9.4 (1) B	4.8 (2) d 6.4 (2) d 5.6 (1) B	1.4 (0.01) d 1.5 (0.1) d 1.4 (0.05) B	0.94 (-) 1.6 (-) 1.3 (0.3) A	0.87 (0.03) c 1.8 (0.1) bc 1.3 (0.1) B	
Mining sites SC2 WF3 BF3 Group mean	328 (40) a 86 (8) b 66 (9) b 160 (38) A	111 (9) a 24 (6) bc 38 (9) b 58 (12) A	26 (0.3) a 9.5° (-) b 3.7 (0.2) c 14 (5) A	11 (-) 1.4 (-) 8.2 (-) 5.5 (2) A	57 <sup>e</sup> (5) a 6.9 (1) b 5.2 (1) bc 23 (6) A	
Downstream sites WF4 BF5 Group mean	18 (0.3) c 8.9 (1) c 13 (2) B	9.9 (5) cd 4.9 (1) d 7.0 (2) B	1.6 (0.2) d 1.1 (0.2) d 1.3 (0.2) B	1.6 (-) 0.96 (-) 1.3 (0.3) A	1.5 (0.1) bc 0.79 (0.1) c 1.1 (0.1) B	

Sites with the same lower case letter and groups of sites with the same capital letter are not significantly different (P<0.05).

and  $10 \,\mu g \, Cd/L$  completely eliminated a crayfish population in an artificial stream. In addition, toxicity of metals to crayfish has been shown to be higher just after molting due to the high permeability of the soft carapace (Knowlton et al., 1983; Wigginton and Birge, 2007). Therefore, tests that include this sensitive life stage are likely to demonstrate greater sensitivity (e.g., Kunz et al., 2005).

None of the metals in our study exceeded chronic WQC at any of the study sites; however, reduced survival of O. hylas may be most associated with Ni concentrations, due to the bioavailability of Ni in surface and pore waters (Besser et al., 2008; Brumbaugh et al., 2007) or reflect the cumulative action of a mixture of metals. Results from our study indicate that O. hylas are highly sensitive to chronic exposure to metals associated with Pb mining. These results are consistent with findings of laboratory studies reported by Kunz et al. (2005), who compared the sensitivity of four aquatic taxa to chronic toxicity of a five-metal mixture (Pb, Zn, Cd, Ni, and Co), which was based on concentrations measured in sediment pore waters from our study streams. Survival of juvenile O. hylas (approximately 30-d old) was reduced at metal concentrations an order of magnitude lower than those affecting rainbow trout (Onchorhynchus mykiss) and mottled sculpin (Cottus bairdi), fishes that are known to be sensitive to metal toxicity (Besser et al., 2007). The relative sensitivity of crustaceans to metal toxicity was further supported by a laboratory study conducted by Besser et al. (2008) that found toxic effects on survival, growth, and reproduction of *H. azteca* to be significantly correlated with metal concentrations in both sediment and sediment pore water collected from mining sites in the Viburnum Trend.

Previous field studies (Mirenda, 1986a, b; Vijayram and Geraldine, 1996) have shown a significant correlation between metal concentrations in water and those in crayfish. In our study, metal concentrations in caged crayfish were significantly higher at mining sites than at reference sites, and were significantly correlated with surface water metal concentrations; however, there were fewer correlations between metal concentrations in surface water and metal concentration in wild crayfish. Wild crayfish on day 0 were significantly larger than crayfish stocked into cages at all sites except WF3, which may have resulted in

fewer molts in wild crayfish during the 28-d exposure. On day 28, only wild crayfish at WF1, BF1, and the single crayfish collected at SC2 were significantly larger than caged crayfish, which indicated caged crayfish grew faster than wild crayfish, and thus molted more frequently than wild crayfish. More frequent molting may have resulted in higher metal concentrations in caged crayfish due to greater permeability in the carapace after molting (Knowlton et al., 1983; Wigginton and Birge, 2007).

Diet may also influence bioaccumulation of metals in crayfish (Besser et al., 2006; Giesy et al., 1980). Besser et al. (2006) reported strong correlations between Pb, Zn, and Cd concentrations in field-collected crayfish, plant material, and benthic organisms in streams within the Viburnum Trend. We also found significant correlations between concentrations of Pb, Zn, Ni, and Co in caged crayfish and metal concentrations in detritus, macroinvertebrates, and stonerollers. Although there were fewer significant correlations between metal concentrations in wild crayfish, there were significant correlations among Pb, Zn, Cd, and Ni concentrations in wild crayfish and metal concentrations in detritus, macroinvertebrates, and stonerollers. Caged and wild crayfish may have fed on different food items due to availability, or differences may reflect dietary preference for macroinvertebrates and fish by juvenile crayfish (Parkyn et al., 2001; Whitledge and Rabeni, 1997).

Biomass of caged crayfish was significantly higher at downstream sites than at reference sites and mining sites, and may reflect higher nutrient concentrations. Nutrient concentrations were significantly higher at mining sites, and probably contributed to the development of large algal mats found at some of our mining sites (Gale et al., 1973). These algal mats may provide wild crayfish with additional food resources; however, algae contain high concentrations of metals at sites downstream of mining (Besser et al., 2006), which may increase metal effects.

#### 5. Conclusions

These findings suggest that despite the use of efficient extraction technologies and operations that meet environmental


 $<sup>^{</sup>a}$  n = 4.

 $<sup>^{\</sup>rm b}$  n=2; Campostoma oligolepis.


 $<sup>^{</sup>c}$  n = 1.

<sup>&</sup>lt;sup>d</sup> n = 6.

 $<sup>^{</sup>e}$  n = 7.



**Fig. 5.** Metal concentrations in food-web components placed in cages at sampling sites: (a) lead; (b) zinc; (c) cadmium; (d) nickel; and (e) cobalt. Sample types: Detr = detritus; Invert = macroinvertebrates; Fish = stoneroller (*Campostoma oligolepis*); Wild = wild *Orconectes hylas*; and Caged = caged *O. hylas*.



**Fig. 6.** Mean metal concentrations in caged *Orconectes hylas* (days 0, 14, and 28), and wild *O. hylas* (day 28) at sampling sites: (a) lead; (b) zinc; (c) cadmium; (d) nickel; and (e) cobalt.

regulations, Pb–Zn mining in the Viburnum Trend is impacting water quality, metal accumulation in food webs, and *O. hylas* populations of the Black River watershed. Our results indicate that

metals were the main factor causing the observed toxic effects on *O. hylas* in streams of the Black River watershed, including elevated concentrations of metals in crayfish. Survival and

**Table 8**Spearman correlation coefficients for correlations among percent (%) survival and biomass on day 28, and metal concentrations in caged crayfish (*Orconectes hylas*) and metal concentrations in surface water, detritus, macroinvertebrates, stonerollers (*Campostoma oligolepis*), and wild crayfish (*O. hylas*) on day 28.

Metal		Caged crayfish						Wild crayfish					
		Lead	Zinc	Cadmium	Nickel	Cobalt	Biomass	% Survival	Lead	Zinc	Cadmium	Nickel	Cobalt
Lead	Surface water Detritus Macroinvertebrates Stonerollers Caged crayfish	0.93 1.0 0.93 0.93	0.85 0.79 0.71 0.86 0.79	-0.19 0.04 0.07 0.18 0.04	0.85 0.93 0.82 0.93 0.93	0.78 0.71 0.57 0.71 0.71	-0.44 -0.43 -0.36 -0.54 -0.43	-0.86 -0.95 -0.95 -0.99 -0.95	0.64 <b>0.83</b> <b>0.94</b> <b>0.89</b> <b>0.83</b>	0.70 <b>0.83</b> 0.77 0.71 <b>0.83</b>	-0.34 -0.43 -0.37 -0.31 -0.43	0.70 <b>0.83</b> 0.77 0.71 <b>0.83</b>	0.58 0.37 0.26 0.49 0.37
Zinc	Surface water Detritus Macroinvertebrates Stonerollers Caged crayfish	0.56 1.0 1.0 0.29 0.79	0.37 <b>0.79</b> <b>0.79</b> 0.57	0.67 0.04 0.04 0.39 0.04	0.30 <b>0.93</b> <b>0.93</b> 0.43 <b>0.86</b>	0.26 0.71 0.71 0.75 <b>0.93</b>	-0.30 -0.43 -0.43 - <b>0.96</b> -0.75	-0.58 - <b>0.95</b> - <b>0.95</b> -0.34 - <b>0.81</b>	0.52 <b>0.83</b> <b>0.83</b> -0.31 0.43	0.39 <b>0.83</b> <b>0.83</b> -0.66 0.37	-0.58 -0.43 -0.43 0.83 -0.09	0.39 <b>0.83</b> <b>0.83</b> -0.66 0.37	-0.03 0.37 0.37 0.03 <b>0.83</b>
Cadmium	Surface Water Detritus Macroinvertebrates Stonerollers Caged crayfish	0.61 <b>1.0</b> 0.39 -0.14 0.04	0.61 <b>0.79</b> 0.32 0.29 0.04	0.61 0.04 0.61 0.71	0.61 <b>0.93</b> 0.14 -0.07 -0.04	0.61 0.71 0.07 0.14 -0.04	-0.61 -0.43 -0.11 -0.46 -0.32	-0.62 - <b>0.95</b> -0.49 -0.07 -0.20	- 0.83 0.49 -0.09 -0.02	- <b>0.83</b> 0.37 -0.37 -0.37	- -0.43 -0.66 0.09 0.09	- <b>0.83</b> 0.37 -0.37 -0.37	- 0.37 0.14 0.20 -0.48
Nickel	Surface Water Detritus Macroinvertebrates Stonerollers Caged crayfish	0.95 0.93 0.96 0.93 0.93	<b>0.81 0.86</b> 0.75 0.71 <b>0.86</b>	0.20 0.18 0.21 0.18 -0.04	0.91 0.93 0.89 0.93	<b>0.68</b> 0.71 0.64 <b>0.68 0.82</b>	-0.50 -0.54 -0.46 -0.54 -0.57	-1.0 -0.99 -0.99 -0.95 -0.91	0.93 0.89 0.94 0.83 0.66	0.75 0.71 0.77 0.60 0.60	-0.35 -0.31 -0.37 -0.09 -0.09	0.75 0.71 0.77 0.60 0.60	0.38 0.49 0.26 0.09 0.43
Cobalt	Surface Water Detritus Macroinvertebrates Stonerollers Caged crayfish	0.86 0.86 0.89 0.82 0.71	0.93 0.93 0.96 0.89 0.93	-0.11 -0.11 -0.07 -0.14 -0.04	0.96 0.96 0.93 0.89 0.82	0.93 0.93 0.89 0.96	-0.68 -0.68 -0.61 -0.71 - <b>0.86</b>	- <b>0.83</b> - <b>0.82</b> - <b>0.86</b> -0.74 -0.69	0.43 0.43 0.54 0.26 0.14	0.43 0.43 0.60 0.31 0.09	0.03 0.03 -0.26 0.09 0.26	0.43 0.43 0.60 0.31 0.09	0.60 0.60 0.77 0.54 0.60

For wild crayfish, no data from SC2 is included. Correlations in boldface are significant (P<0.05).

biomass data from the in-situ toxicity test suggest that recovery of crayfish populations may occur approximately 10 km downstream of mining sites. The absence of crayfish may have negative effects on organic (e.g., leaves, woody debris) processing, nutrient cycling, and energy transfer in Ozark streams. Crayfish are the predominant prey item of smallmouth bass, an important species sought by recreational anglers in the Ozarks (Mayers, 2003; Weithman, 1991), and other centrarchid fishes in Ozark streams (DiStefano, 2005; Probst et al., 1984). High metal concentrations in centrarchids are likely related to high metal concentrations in crayfish, which could potentially represent a hazard to both wildlife and humans (Schmitt et al., 2006).

#### Acknowledgments

Funding for this research was provided by the US Government through the US Geological Survey. We thank the private landowners and the Doe Run Corporation who allowed access to these streams. CERC personnel who assisted included: J. Arms, E. Brunson, J.M. Fairchild (CERC volunteer), K. Grabner, L. Johnson, S. Koppi, T. May, S. Olson, B. Poulton, L. Sappington, D. Stoppler, M. Struckhoff, C. Vishy, M. Walther, D. Whites, and C. Witte. Missouri Department of Conservation personnel who assisted were: T. Boersig, L. Cole, I. Dasho (an AFS Hutton Junior Biologist), S. Herleth-King, J. Lane, and J. Westhoff. We thank K. Buhl, L. Cleveland, and two anonymous reviewers, for providing insightful comments that greatly improved the quality of this manuscript.

#### Disclaimer

Crayfish were collected in accordance with Missouri Department of Conservation Wildlife Collectors Permit regulations. To ensure humane treatment of the test organisms during culture and experimentation, this research was conducted according to the CERC Animal Welfare plan, in compliance with requirements of the US Laboratory Animal Welfare Act and the Intergency Research Animal Committee. Use of trade names does not constitute USGS or US Government endorsement.

#### Appendix A. Supplementary Materials

Supplementary data associated with this article can be found in the online version at doi:10.1016/j.ecoenv.2008.08.005.

#### References

Alikhan, M.A., Bagatto, G., Zia, S., 1990. Crayfish as a "biological indicator" of aquatic contamination by heavy metals. Water Resour. 24, 1069–1076.

Allert, A.L., Fairchild, J.F., DiStefano, R.J., Schmitt, C.J., Besser, J.M., Brumbaugh, W.G., Poulton, B.C., 2008. Effects of lead-zinc mining on crayfish (Orconectes hylas) in the Black River watershed, Missouri, USA. Freshwater Crayfish 16, 99-113.

APHA (American Public Health Association), American Water Works Association, Water Environment Federation, 2005. Standard Methods for the Examination of Water and Wastewater, 21st ed. American Public Health Association, Alexandria, VA.

Besser, J.M., Brumbaugh, W.G., May, T.W., Schmitt, C.J., 2006. Biomonitoring of lead, zinc, and cadmium in streams draining lead-mining and non-mining areas, southeast Missouri, USA. Environ. Monit. Assess. 129, 227–241.

- Besser, J.M., Mebane, C.A., Mount, D.R., Ivey, C.D., Kunz, J.L., Greer, I.E., May, T.W., Ingersoll, C.G., 2007. Sensitivity of mottled scuplins (*Cottus bairdi*) and rainbow trout (*Onchorhynchus mykiss*) to acute and chronic toxicity of cadmium, copper, and zinc, Environ. Tox. Chem. 26, 1657–1665.
- Besser, J.M., et al., 2008. Ecological impacts of lead mining on Ozark streams: toxicity of sediment and pore water. Ecotoxicol. Environ. Saf. 72, 516–526.
- Brumbaugh, W.G., Schmitt, C.J., May, T.W., 2005. Concentrations of cadmium, lead, and zinc in fish from mining-influenced waters of Northeastern Oklahoma: sampling of blood, carcass, and liver for aquatic biomonitoring. Arch. Environ. Contam. Toxicol. 49, 76–88.
- Brumbaugh, W.G., May, T.W., Besser, J.M., Allert, A.L., Schmitt, C.J., 2007.
  Assessment of elemental concentrations in streams of Missouri's New Lead
  Belt: 2002–2005. US Geological Survey Report 2007-5057.
- DiStefano, R.J., 2005. Trophic interactions between Missouri Ozarks stream crayfish communities and sport fish predators: increased abundance and size structure of predators cause little change in crayfish community densities. Missouri Department of Conservation, Dingell-Johnson Project F-1-R-054, Study S-41, Job 4, Final Report. Columbia, MO.
- DiStefano, R.J., Young, J., Noltie, D.B., 2002. A study of the life history of *Orconectes hylas* with comparisons to *Orconectes peruncus* and *Orconectes quadruncus* in Ozark streams. Missouri. USA. Freshwater Crayfish 13. 439–456.
- Duchrow, R.M., 1983. Effects of lead tailings on benthos and water quality in three Ozark streams. Trans. Miss. Acad. Sci. 17. 5–17.
- Gale, N.L., Wixson, B.G., Hardie, M.G., Jennett, J.C., 1973. Aquatic organisms and heavy metals in Missouri's new lead belt. Water Resour. Bull. 9, 673–688.
- Giesy, J.P., Bowling, J.W., Kania, H.J., 1980. Cadmium and zinc accumulation and elimination by freshwater crayfish. Arch. Environ. Contam. Toxicol. 9, 683–697.
- Gray, N.F., 1998. Acid mine drainage composition and the implications for its impact on lotic systems. Water Resour. 32, 2122–2134.
- Flinders, C.A., Magoulick, D.D., 2005. Distribution, habitat use and life history of stream-dwelling crayfish in the Spring River drainage of Arkansas and Missouri with a focus on the Mammoth Spring crayfish (*Orconectes marchandi*). Am. Mid. Nat. 154. 358–374.
- Huryn, A.D., Wallace, J.B., 1987. Production and litter processing by crayfish in an Appalachian mountain stream. Freshwater Biol. 18, 277–286.
- Jessey, D.R., 1981. An investigation of the nickel-cobalt occurrence in the southeast Missouri mining district: Rolla, MO. Ph.D. Dissertation, University of Missouri-Rolla.
- Knowlton, M.F., Boyle, T.P., Jones, J.R., 1983. Uptake of lead from aquatic sediment by submersed macrophytes and crayfish. Arch. Environ. Contam. Toxicol. 12, 535–541.
- Kuck, P.H., 2004. Nickel. In Minerals Yearbook, Volume I, Metals and Minerals: US Geological Survey, accessed January 2007 at URL <a href="http://minerals.usgs.gov/minerals/pubs/myb.html">http://minerals.usgs.gov/minerals/pubs/myb.html</a>.
- Kunz, J.L., Besser, J.M., May, T.W., Ingersoll, C.W., 2005. Toxicity of a metal mixture to stream-dwelling sculpins and crayfish. Poster TP109. SETAC North American 26th Annual Meeting, Baltimore, MD.
- Lindhjem, P.A., Bennet-Chambers, M.G., 2002. Bioaccumulation and acute toxicity of zinc in marron, *Cherax tenuimanus* (Smith) [Decapoda: Parastacidae]. Freshwater Crayfish 13, 424–430.
- Lodge, D.M., Kershner, J.W., Aloi, J.E., 1995. Effects of an omnivorous crayfish (*Oronectes rusticus*) on a freshwater littoral food web. Ecology 75, 165–1281.
- May, T.W., Wiedmeyer, R.H., Brumbaugh, W.G., Schmitt, C.J., 1997. The determination of metals in sediment pore waters and in 1N HCl-extracted sediments by ICP-MS. Atom. Spectrosc. 18, 133–139.
- Mayers, D.A., 2003. 2001 Mail Survey Results of Lower Current River Anglers. Missouri Department of Conservation, Jefferson City, MO.
- Mirenda, R.J., 1986a. Acute toxicity and accumulation of zinc in the crayfish, Orconectes virilis (Hagen). Bull. Environ. Contam. Toxicol. 37, 387–394.
- Mirenda, R.J., 1986b. Toxicity and accumulation of cadmium in the crayfish, Orconectes virilis (Hagen). Arch. Environ. Contam. Toxicol. 15, 401–407.
- Missouri Department of Natural Resources, Geological Survey and Resources Assessment Division, 2004. The mineral industry of Missouri. In: Minerals

- Yearbook, Volume I: US Geological Survey, accessed January 2007, at URL <a href="http://minerals.usgs.gov/minerals/pubs/state/2004/mostmyb04.pdf">http://minerals.usgs.gov/minerals/pubs/state/2004/mostmyb04.pdf</a>.
- Missouri Department of Health and Senior Services, 2005. 2005 Fish Advisory, accessed February 2008, at URL <a href="http://www.dhss.mo.gov/NewsAndPublicNotices/05FishAdvisory.pdf">http://www.dhss.mo.gov/NewsAndPublicNotices/05FishAdvisory.pdf</a>.
- Nagpal, N.K., 2004. Water quality guidelines for cobalt (electronic resource). Technical Report, accessed March 2008, at URL <a href="http://www.env.gov.bc.ca/wat/wq/BCguidelines/cobalt/cobalt\_tech.pdf">http://www.env.gov.bc.ca/wat/wq/BCguidelines/cobalt/cobalt\_tech.pdf</a>>.
- Naqvi, S.M., Howell, R.D., 1993. Toxicity of cadmium and lead to juvenile red swamp crayfish, *Procambarus clarkia*, and the effects on fecundity of adults. Bull. Environ. Contam. Toxicol. 51, 303–308.
- Parkyn, S.M., Collier, K.J., Hicks, B.J., 2001. New Zealand stream crayfish: functional omnivores but trophic predators. Freshwater Biol. 46, 641–652.
- Pflieger, W.L., 1996. The Crayfishes of Missouri. Missouri Department of Conservation, Jefferson City, MO.
- Probst, W.E., Rabeni, C.F., Covington, W.G., Marteney, R.E., 1984. Resource use by stream-dwelling rock bass and smallmouth bass. Trans. Am. Fish. Soc. 113, 283–294.
- Rabeni, C.F., Gossett, M., McClendon, D.D., 1995. Contribution of crayfish to benthic invertebrate production and trophic ecology of an Ozark stream. Freshwater Crayfish 10, 163–173.
- Ryck, F.M., Whitley, J.R., 1974. Pollution abatement in the lead mining district of Missouri. In: Proceedings of Purdue Industrial Waste Conference, vol. 29, pp. 857–863
- Schmitt, C.J., Brumbaugh, W.G., Linder, G.L., Hinck, J.E., 2006. A screening-level assessment of lead, cadmium, and zinc in fish and crayfish from northeastern Oklahoma. USA. Environ. Geochem. Health 38, 445–471.
- Schmitt, C.J., Brumbaugh, W.G., May, T.W., 2007a. Accumulation of metals in fish from lead-zinc mining areas of southeastern Missouri, USA. Ecotoxicol. Environ. Saf. 67, 14–30.
- Schmitt, C.J., Whyte, J.J., Roberts, A.P., Annis, M.L., Tillitt, D.E., 2007b. Biomarkers of metals exposure in fish from lead–zinc mining in southeastern Missouri, USA. Ecotoxicol. Environ. Saf. 67, 31–47.
- Shedd, K.B., 2004. Cobalt. In: Minerals Yearbook, Volume I, Metals and Minerals: US Geological Survey, accessed January 2007 at URL <a href="http://minerals.usgs.gov/minerals/pubs/myb.html">http://minerals.usgs.gov/minerals/pubs/myb.html</a>.
- Thorp, J.H., Giesy, J.P., Wineriter, S.A., 1979. Effects of chronic cadmium exposure on crayfish survival, growth, and tolerance to elevated temperatures. Arch. Environ. Contam. Toxicol. 8, 449–456.
- Tiwary, R.K., 2001. Environmental impacts of coal mining on water regime and its management. Water Air Soil Pollut. 132, 185–199.
- USEPA, 2002. National Recommended Water Quality Criteria: 2002. Accessed January 2007, at <a href="http://www.epa.gov/waterscience/criteria/aqlife.html">http://www.epa.gov/waterscience/criteria/aqlife.html</a>.
- Usio, N., 2000. Effects of crayfish on leaf processing and invertebrate colonization of leaves in a headwater stream: decoupling of a trophic cascade. Oecologia 124. 608–614.
- Vijayram, K., Geraldine, P., 1996. Regulation of essential heavy metals (Cu, Cr, and Zn) by the freshwater prawn *Macrobrachium malcolmsonii* (Milne Edwards). Bull. Environ. Contamin. Toxicol. 56, 335–342.
- Weithman, A.S., 1991. Recreational use and economic value of Missouri fisheries. Missouri Department of Conservation. Federal Aid in Sport Fish Restoration Project F-1-R-40, Study SI-1. Jefferson City, MO.
- Whitledge, G.W., Rabeni, C.F., 1997. Energy sources and ecological role of crayfishes in an Ozark stream: insights from stable isotopes and gut analysis. Can. J. Fish. Aquat. Sci. 54, 2555–2563.
- Wigginton, A.J., Birge, W.J., 2007. Toxicity of cadmium to six species in two genera of crayfish and the effect of cadmium on molting success. Environ. Toxicol. Chem. 26, 548–554.
- Wixson, B.G., 1977. The Missouri lead study, vol. 1: an interdisciplinary investigation of environmental pollution by lead and other heavy metals from industrial development in the New Lead Belt of southeastern Missouri. Final report to the National Science Foundation. University of Missouri, Columbia and Rolla, MO.