# **Emergence and Survival of Legumes Seeded into Pastures Varying** in Landscape Position

John A. Guretzky,\* Kenneth J. Moore, Allen D. Knapp, and E. Charles Brummer

#### **ABSTRACT**

Landscape position affects legume establishment in pastures. We conducted this study to (i) determine the role of emergence and survival on establishment of legumes on summit and backslope positions and (ii) examine how competition, as influenced by sward-cutting height and N fertilization, affects legume emergence and survival across these positions. We no-till drilled a mixture composed of red clover (Trifolium pratense L.), alfalfa (Medicago sativa L.), and birdsfoot trefoil (Lotus corniculatus L.) into an existing cool-season grass pasture at Rhodes, IA, in 1998 and 1999. Treatments consisted of landscape positions (summit and backslope), N fertilizer rates (0, 22, 44, and 89 kg ha<sup>-1</sup>), and sward heights (uncut, 5, and 13 cm). Landscape position did not affect legume emergence in swards cut at 5 and 13 cm. At those heights, emergence averaged 1041 and 831 seedlings m<sup>-2</sup>. In swards cut at 5 cm, 62 plants m<sup>-2</sup> survived on summits vs. 183 plants m<sup>-2</sup> on backslopes. Legume emergence and survival generally decreased as sward height and N fertilization increased but interactions with landscape position occurred. Addition of 89 kg ha<sup>-1</sup> N was required to reduce survival on backslopes to 39 plants m<sup>-2</sup>, a density similar to that on summits. Our results show that seedling survival limits legume establishment on summit positions in pastures. Legume establishment is successful on backslopes because of less competition from grass. We recommend that legumes be seeded on backslope positions in pastures, N fertilizer not be applied, and grass competition be reduced before seeding legumes in pastures.

Legumes improve the production and nutritional value of forage in cool-season grass pastures. Pastures composed of grasses and legumes produce more dry matter than grass-only pastures (Taylor and Allinson, 1983). Through symbiotic N<sub>2</sub> fixation with rhizobial bacteria, legumes increase soil mineral N (Heichel, 1985). The transfer of N, from the decomposition of roots and leaf residues of legumes, improves the nutritional value and growth of the grass (Heichel and Henjum, 1991). Legumes also are higher in crude protein than are grasses (Van Soest, 1982), and their establishment with cool-season grasses can improve the seasonal distribution of forage growth and quality in pastures (Sleugh et al., 2000).

Harmoney et al. (2001) found establishment of legumes to vary according to landscape position. Legumes contributed more to forage production on backslope than on summit or toeslope positions, and the proportion of dry matter comprised of legumes was related linearly to diversity of leguminous species in pastures. They could not conclude why legume establishment differed at summit and backslope landscape positions, al-

Dep. of Agronomy, Iowa State Univ., Ames, IA 50011-1010. Received 1 July 2002. \*Corresponding author (john.a.guretzky@erdc.usace.army.mil).

Published in Crop Sci. 44:227–233 (2004). © Crop Science Society of America 677 S. Segoe Rd., Madison, WI 53711 USA though differential competitive pressures from established grasses may have been the cause.

Competition from grass often limits the establishment of legumes in pastures. Legume interseeding is most successful when competition from grasses is reduced through herbicide application (Cuomo et al., 2001; Groya and Sheaffer, 1981) or by cutting swards to low heights before seeding legumes (Taylor and Allinson, 1983). Soil N availability also influences grass-legume competition. Grasses have a greater relative growth rate than legumes in areas with high soil N availability. In areas with low soil N availability, legumes are more competitive because they can supplement mineral N uptake with N<sub>2</sub> fixation (Schwinning and Parsons, 1996).

How competition, as influenced by sward height and N fertilization, affects legume establishment in pastures varying in landscape position is unknown. Our objectives were to (i) determine the role of emergence and survival during the seeding year on establishment of several legumes sown as a mixture on summit and backslope positions in pastures and (ii) examine how sward height and N fertilization affect emergence and survival of legumes on these landscape positions.

### **MATERIALS AND METHODS**

This experiment was conducted at the Iowa State University Rhodes Research Farm (41°52′ N, 93°10′ W). A 1737 pure live seed (PLS) m<sup>-2</sup> mixture consisting of red clover (*T. pratense* 'Marathon'), alfalfa (*M. sativa* 'Alfagraze'), and birdsfoot trefoil (*L. corniculatus* 'Norcen') was no-till drilled using a cone seeder (Almaco, Nevada, IA) into a cool-season grass pasture composed of three cool-season grasses: smooth brome (*Bromus inermis* Leyss.), Kentucky bluegrass (*Poa pratensis* L.), and reed canarygrass (*Phalaris arundinacea* L.). The legumes were sown at 582 PLS m<sup>-2</sup> of red clover, 587 PLS m<sup>-2</sup> of alfalfa, and 568 PLS m<sup>-2</sup> of birdsfoot trefoil. On a PLS basis, these rates corresponded to 11.3 kg ha<sup>-1</sup> red clover, 12.6 kg ha<sup>-1</sup> alfalfa, and 8.6 kg ha<sup>-1</sup> birdsfoot trefoil. The legumes were inoculated with a humus-based rhizobium specific to each legume.

We conducted the experiment in small plots, 1.4 m² in 1998 and 2.8 m² in 1999, to facilitate counts of legume emergence and survival under several treatments. Treatments consisted of two landscape positions (summit and backslope), four N fertility rates (0, 22, 44, and 89 kg ha⁻¹) applied as NH₄NO₃, and three sward heights (uncut, 5, and 13 cm above the soil). The backslope had a west-facing aspect and 10% slope. The summit had zero aspect and slope. Tillage and herbicide burndown are common procedures to establish legumes in pastures. However, we attempted to follow procedures of Harmoney et al. (2001), reducing competition from existing grasses by a mechanical rather than a chemical means. Also, because aspect does not affect legume establishment or persistence in

these pastures (Harmoney et al., 2001), it was not incorporated into this experiment.

For several years before 1998, the pasture was fertilized annually in the spring with N, P, and K at 77, 26, and 34 kg ha<sup>-1</sup>, respectively, cut once for hay, and grazed by cattle July through August. In 1998, ground adjacent to the experimental plots was cut and raked with a tractor but was not fertilized or grazed. Plots in 1999 were arranged  $\approx$ 3 m away from the 1998 plots. Before seeding, the abundance of litter appeared to be similar among 1998 and 1999 plots.

Plots were seeded on 28 Apr. 1998 and 20 May 1999. Nitrogen treatments were applied on 27 Apr. 1998 and 26 Apr. 1999. Wet-field conditions delayed planting following N application in 1999. Cutting dates in 1998 were 27 April, 7 May, 19 May, and 19 June for the 5-cm height treatment and 7 May, 19 May, and 19 June for the 13-cm height treatment. In 1999, the cutting dates were 26 Apr., 21 May, and 7 June for the 5-cm height and 21 May and 7 June for the 13-cm height. Uncut swards were not cut until August of each year. In August, all plots were cut at 13 cm. The 5 and 13-cm height treatments consisted of clipping the grass with a sickle bar push mower and lightly hand-raking the plots. Mowing did not result in a chopping of the grass, only clipping at the respective height. Raking removed the fresh, clipped grass lying on top of the canopy but did not disturb existing litter. Clipping was terminated before legume seedlings reached 5 cm in height.

Our goal with the sward height and N fertilizer rate treatments was to establish several levels of competition. Because sward height treatments were used only to reduce competition from existing grasses, sward heights or yields of grass were not measured at the time of clipping. We clipped the 5-cm height treatment frequently to inhibit grass competition. The 13-cm height treatment was cut less frequently and the uncut sward not at all to establish the different levels of competition.

Because of wet field conditions, the 5 and 13-cm height treatments were cut one less time in 1999 than in 1998.

We measured legume emergence using a destructive sampling technique. The technique involved counting, identifying, and then removing each legume seedling that emerged within a 0.1-m<sup>2</sup> subplot. Emergence was measured on 15 May, 4 June, and 30 June 1998 and 9 and 28 June 1999. As a percentage of overall emergence, 88, 10, and 2% of legumes emerged in 1998 and 80 and 20% of overall legumes emerged in 1999 by those dates, respectively. The data we present represents the sum of emergence after all sampling dates each year. Seedling survival was measured within a different 0.1-m<sup>2</sup> subplot on 15 Aug. 1998 and 1999, 109 and 87 d after planting in 1998 and 1999, respectively. We used a LAI-2000 Plant Canopy Analyzer (LI-COR, Lincoln, NE) to measure leaf area index (LAI) of the grass swards. Leaf area index was measured on 28 May and 22 June 1998 and 22 June and 16 July 1999, before the legumes produced appreciable leaf area.

We conducted the experiment in a randomized complete block. Four replicated blocks were established on each landscape position, and separate blocks were established for each seeding year. Within each block, N fertilizer rate and swardheight treatments were arranged in a split plot. Nitrogen fertilizer rate treatments were randomly arranged as whole plots. Sward height treatments were randomly arranged within each N fertilizer rate as subplots. Every block consisted of 12 plots. We considered the effects of landscape position, N, and sward height fixed and replications random. We conducted an analysis of variance to test these effects by using F ratios for splitplot experiments arranged in a randomized complete block design (McIntosh, 1983). Because interactions of treatments and years occurred, the analysis of variance was conducted by year (Table 1). Significance was defined at  $P \le 0.05$ . However, if treatment effects were significant at  $P \le 0.10$  in one

Table 1. Analysis of variance (ANOVA) for emergence and survival of a legume mixture interseeded into a cool-season grass pasture at Rhodes, IA, in 1998 and 1999. Sources of variation included two landscape positions (L), four N fertilizer rates (N), three sward heights (H), and four replications (R). Mean squares (MS) are presented for overall legume, red clover, alfalfa, and birdsfoot trefoil emergence and survival by year.

			Overa	ll legume	Red	Red clover		Alfalfa		Birdsfoot trefoil	
Source	F test	df	1998 MS	1999 MS	1998 MS	1999 MS	1998 MS	1999 MS	1998 MS	1999 MS	
			Emergence								
L	MS(L)/MS[R(L)]	1	12 558	349 451	33 153	76 784	1 240	33 153	66 993	17 281	
R(L)	- ( ) E ( ) J		342 513	168 992	51 142	32 288	51 369	24 841	31 961	10 162	
N N	$MS(N)/MS[R \times N(L)]$	3	70 243	1 921 573***	19 460	338 338***	7 327	202 810***	24 648	128 961***	
$L \times N$	$MS(L \times N)/MS[R \times N(L)]$	3	46 980	251 941	11 246	54 633	16 379	23 298	9 960	18 608	
$\mathbf{R} \times \mathbf{N}(\mathbf{L})$	. , , , , , , , , , , , , , , , , , , ,	18	133 481	170 476	23 891	32 362	20 545	20 435	11 726	15 793	
H	$MS(H)/MS[R \times N \times H(L)]$		164 054	3 774 791***	10 578	784 986***	4 026	287 179***	74 123**	278 910***	
$L \times H$	$MS(L \times H)/MS[R \times N \times H(L)]$	2	104 999	139 936	17 186	35 253	17 078	7 707	16 028	10 166	
$N \times H$	$MS(N \times H)/MS[R \times N \times H(L)]$	6	196 746*	119 832	18 517	23 271	33 996**	9 208	20 323	12 468	
$L \times N \times H$	$MS(L \times N \times H)/MS[R \times N \times H(L)]$	6	243 468**	237 022†	44 423**	62 937*	25 312†	21 462	21 741†	15 384	
$\mathbf{R} \times \mathbf{N} \times \mathbf{H}(\mathbf{L})$		36	69 957	113 257	14 490	24 133	11 299	14 577	11 359	32 989	
						Survi	val				
L	MS(L)/MS[R(L)]	1	61 560**	173 740***	45 067*	66 993***	13	1 935†	1 528†	13 090**	
R(L)		6	4 111	1 193	2 101	686	108	425	284	493	
N N	$MS(N)/MS[R \times N(L)]$	3	5 552	25 370***	2 967	5 387*	82	1 015*	570	3 684**	
$L \times N$	$MS(L \times N)/MS[R \times N(L)]$	3	4 911	9 435*	2 859	2 257	13	387	573	1 359	
$\mathbf{R} \times \mathbf{N}(\mathbf{L})$		18	3 998	2 839	2 652	1 075	56	249	303	615	
H	$MS(H)/MS[R \times N \times H(L)]$	2	51 145***	161 044***	28 574***	46 014***	855***	7 770***	1 423***	10 076***	
$L \times H$	$MS(L \times H)/MS[R \times N \times H(L)]$	2	3 911†	50 809***	3 609*	18 095***	41	922†	129	3 771***	
$N \times H$	$MS(N \times H)/MS[R \times N \times H(L)]$	6	2 215	11 677*	1 495	3 604*	70	467	170	1 729***	
$L \times N \times H$	$MS(L \times N \times H)/MS[R \times N \times H(L)]$	6	1 415	7 784	700	2 578	18	217	290	1 104**	
$\mathbf{R} \times \mathbf{N} \times \mathbf{H}(\mathbf{L})$		36	1 568	4 262	828	1 583	56	374	233	369	

<sup>\*</sup> Denotes significance at the 0.05 probability level.

<sup>\*\*</sup> Denotes significance at the 0.01 probability level.

<sup>\*\*\*</sup> Denotes significance at the 0.001 probability level.

<sup>†</sup> Denotes significance at the 0.100 probability level.

year and  $P \le 0.05$  the other year, both were considered significant

# **RESULTS Emergence**

#### **Landscape Position Effects**

In 1998 and 1999, overall legume emergence was similar among summits and backslopes at all N rates in swards cut at 5 and 13 cm (Table 2). At those heights, overall legume emergence averaged 1041 and 831 seedlings m<sup>-2</sup> across years. Overall legume emergence was greater on backslopes than on summits by 686 plants m<sup>-2</sup> in uncut swards fertilized at 22 kg ha<sup>-1</sup> N in 1998 and 1003 plants m<sup>-2</sup> in uncut swards at 0 kg ha<sup>-1</sup> N in 1999. Emergence was greater for red clover than alfalfa and birdsfoot trefoil in 1998 and 1999. Landscape position effects on emergence of each species, however, were similar to their effects on overall legume emergence (Table 2).

# **Nitrogen Fertilization Effects**

Nitrogen fertilization did not affect overall legume emergence in swards cut at 5 cm in 1998 or 1999 nor in swards cut at 13 cm in 1998 (Table 2). Addition of N from 0 to 89 kg ha<sup>-1</sup> reduced overall legume emergence by 68% in swards cut at 13 cm in 1999. In uncut swards, addition of N from 0 to 89 kg ha<sup>-1</sup> reduced overall

legume emergence on backslopes by 37% in 1998 and 92% in 1999. Nitrogen fertilization had negligible effects on overall legume emergence on summits. Effects of N fertilization on red clover, alfalfa, and birdsfoot trefoil emergence were similar to those on overall legume emergence in 1998 and 1999.

#### **Sward Height Effects**

Overall legume emergence was similar among sward heights at most N fertilization rates on summits and backslopes in 1998. In 1999, increased sward height reduced overall legume emergence at all N rates on summits and at 44 and 89 kg ha<sup>-1</sup> N on backslopes by 61 to 98% (Table 2). Sward height affected the emergence of red clover, alfalfa, and birdsfoot trefoil similarly as overall emergence in 1998 and 1999.

#### **Survival**

# **Landscape Position Effects**

Landscape position and sward height interactions affected overall legume survival in 1998 and 1999 (Table 1). In swards cut at 5 cm, overall legume survival was greater on backslopes than on summits by 69 plants m<sup>-2</sup> in 1998 and 173 plants m<sup>-2</sup> in 1999 (Table 3). In swards cut at 13 cm and in uncut swards, overall legume survival was, on average across years, greater on backslopes than on summits by 60 and 22 plants m<sup>-2</sup>, respec-

Table 2. Landscape position, sward height, and N fertilizer rate effects on emergence of legumes interseeded into a cool-season grass pasture at Rhodes, IA. Legumes were interseeded as a mixture across summit (S) and backslope (B) landscape positions on 28 Apr. 1998 and 20 May 1999. Values represent sum of overall legume, red clover, alfalfa, and birdsfoot trefoil emergence after measuring on 15 May, 4 June, and 30 June 1998 and 9 and 28 June 1999 (n = 4).

			Emergence of legumes								
		r Sward height	Nitrogen, kg ha <sup>-1</sup>								
Legume species			0		22		44		89		
	Year		S	В	S	В	S	В	S	В	SE
		cm		seedlings m <sup>-2</sup>							
Overall	1998	5	1047	853	893	842	1025	1028	1103	1023	233.0†
		13	675	1052	893	648	807	950	971	710	213.5‡
		Uncut	1068	1068	678	1364	960	686	500	673	187.0§
	1999	5	1217	1480	1386	1015	958	969	818	993	257.1
		13	1074	1066	928	1168	<b>791</b>	880	361	329	257.2
		Uncut	345	1348	412	608	374	162	14	108	238.0
Red clover	1998	5	441	288	385	315	498	374	412	406	119.6
		13	315	398	388	266	355	409	422	302	114.8
		Uncut	481	398	291	508	525	323	245	323	117.4
	1999	5	506	684	654	420	490	466	371	493	115.9
		13	544	474	428	525	369	428	180	202	115.9
		Uncut	159	614	197	288	178	102	3	62	109.8
Alfalfa	1998	5	371	248	280	315	334	328	390	307	113.4
		13	213	363	312	258	296	377	355	261	107.6
		Uncut	361	339	288	516	261	229	202	207	110.5
	1999	5	415	436	415	342	256	304	240	280	91.9
		13	266	345	261	350	243	245	108	65	90.9
		Uncut	135	401	129	237	170	49	11	41	85.3
Birdsfoot trefoil	1998	5	234	318	229	213	194	326	301	309	81.1
		13	148	291	194	124	156	164	194	148	75.8
		Uncut	226	331	100	339	175	135	54	143	75.4
	1999	5	296	361	318	253	213	199	207	221	75.0
		13	264	248	240	294	181	207	73	62	76.5
		Uncut	51	334	86	83	27	11	0	6	69.6

<sup>†</sup> SE, by year and legume species, for comparison of two landscape position means at the same or different N rate and sward height.

<sup>‡</sup> SE, by year and legume species, for comparison of two N rate means at the same landscape position and sward height.

<sup>§</sup> SE, by year and legume species, for comparison of two sward height means at the same N rate and landscape position.

Table 3. Landscape position and sward height effects on survival of legumes interseeded as a mixture into a cool-season grass pasture at Rhodes, IA. Values represent overall legume, red clover, alfalfa, and birdsfoot trefoil survival on 15 Aug. 1998 and 1999 (n = 16).

			Survival of legumes						
Legume species	Year	Landscape position	5	13		Uncut	SE		
					plants m <sup>-2</sup>				
Overall	1998	Summit	63	20	F	5	14.0†		
3,61	2570	Backslope	132	77		31	2.00		
		SE	102	,,	<b>17.4</b> ‡	<b>71</b>			
	1999	Summit	61	11	17007	3	23.1		
	1,,,,	Backslope	234	75		21	2011		
		SE	254	75	20.1	21			
Red clover	1998	Summit	42	18	20.1	2	10.2		
Rea cloves	1770	Backslope	101	69		22	10.2		
		SE SE	101	0)	12.5				
	1999	Summit	28	7	12.0	1	14.1		
	1777	Backslope	133	50		12	14.1		
		SE SE	133	30	12.7	12			
Alfalfa	1998	Summit	12	1	12.7	1	2.6		
Allalla	1770	Backslope	9	2		0	2.0		
		SE	,	2	3.0	U			
	1999	Summit	18	3	3.0	1	6.8		
	1777	Backslope	42	9		3	0.0		
		SE	72	,	7.0	3			
Birdsfoot trefoil	1998	Summit	10	1	7.0	2	5.4		
Bitusioot treion	1770	Backslope	22	5		2 9	3.4		
		SE	22	3	5.6	,			
	1999	Summit	14	3	3.0	2	6.8		
	1777	Backslope	62	20		2 7	0.0		
		SE	02	20	7.2	1			
		SE			1.4				

<sup>†</sup> SE by year and legume species for comparison of two sward height means at the same landscape position.

tively (Table 3). Red clover had the greatest survival of the legume species and was affected similarly by landscape position as overall legume survival in 1998 and 1999.

Survival of alfalfa and birdsfoot trefoil was similar among summits and backslopes in 1998 (Table 3). In 1999, alfalfa survival was 24 plants m<sup>-2</sup> greater on backslopes than on summits in plots cut at 5 cm but similar among landscape positions in swards cut at 13 cm and in uncut swards. Birdsfoot trefoil survival was also greater on backslopes than on summits in plots cut at 5 cm (Table 3) and in unfertilized swards cut at 13 cm in 1999 (data not shown).

## **Nitrogen Fertilization Effects**

Overall legume survival was also affected by a land-scape position and N fertilization interaction in 1999. Neither N main effects nor the landscape position and N interaction on overall legume survival were statistically significant in 1998 (Table 1). In both years, addition of N from 0 to 44 kg ha<sup>-1</sup> reduced overall legume survival similarly on summits and backslopes (Fig. 1). Across these rates, overall legume survival was greater on backslopes than on summits by 64 plants m<sup>-2</sup> in 1998 and 105 plants m<sup>-2</sup> in 1999. When N fertilization increased to 89 kg ha<sup>-1</sup>, overall legume survival declined on backslopes to 39 plants m<sup>-2</sup> across years, a density similar to that on summits (Fig. 1).

The effect of N fertility on overall legume survival also varied with sward height in 1999. In swards cut at 5 cm, overall legume survival increased from 155 plants m<sup>-2</sup> at 0 kg ha<sup>-1</sup> N to 217 plants m<sup>-2</sup> at 22 kg ha<sup>-1</sup> N (Table 4). In 13-cm and uncut swards, addition of N

fertilizer decreased overall legume survival. Similar results occurred for red clover and birdsfoot trefoil as overall legume survival in 1999. Survival of alfalfa decreased with increased N fertilization at all sward heights in 1999. Nitrogen fertilization did not affect red clover, alfalfa, or birdsfoot trefoil survival in 1998 (Table 1).

#### **Sward Height Effects**

Overall legume survival was greatest in swards cut at 5 cm (Table 3). Across years, overall legume survival on summits was 46 and 58 plants m<sup>-2</sup> greater in swards cut at 5 cm than in swards cut at 13 cm and in uncut swards, respectively. On backslopes in 1998 and 1999, respectively, overall legume survival was greater in swards cut at 5 cm than swards cut at 13 cm by 55 and 159 plants m<sup>-2</sup> and greater in swards cut at 5 cm than uncut swards by 101 and 213 plants m<sup>-2</sup>. Individually, survival of each legume species also decreased with increased sward height in 1998 and 1999 (Table 3).

# **Sward Leaf Area Index**

Leaf area index of the grass sward increased at a greater rate on summits than on backslopes. From 1 to 22 June 1998, LAI increased from 3.0 to 4.7 on summits and from 2.3 to 2.9 on backslopes (Table 5). Similar results occurred between 22 June and 16 July 1999. Leaf area index also increased with addition of N fertilizer (Table 5). Landscape position and N fertilizer rate interactions did not occur, however. Cutting at a greater height also increased LAI of the grass sward in 1998 and 1999 (Table 5).

<sup>‡</sup> SE by year and legume species for comparison of two landscape position means at the same or different sward height.

Table 4. Sward height and N fertilizer rate effects on survival of legumes interseeded as a mixture into a cool-season grass pasture at Rhodes, IA. Values represent overall legume, red clover, alfalfa, and birdsfoot trefoil survival on 15 Aug. 1998 and 1999 (n = 8).

		Sward height	Survival of legumes						
Legume species	Year		0	22	44	89	SE		
		cm		pla	nts m <sup>-2</sup>				
Overall	1998	5	126	100	79	85	24.3†		
		13	58	44	71	22			
		Uncut	30	40	4	0			
		SE			19.8‡	*			
	1999	5	155	217	152	66	30.8		
		13	107	26	38				
		Uncut	33	10	1	2 3			
		SE			32.6	_			
Red clover	1998	5	90	67	65	65	18.9		
		13	53	36	68	19			
		Uncut	22	23	3	0			
		SE	14.3						
	1999	5	70	115	93	45	18.8		
	2,,,,	13	61		30		1010		
		Uncut	18	20 5	1	2 3			
		SE	10		19.9				
Alfalfa	1998	5	11	16	4	10	3.7		
1	2570	13	2	1	1	1			
		Uncut	2 3	0	0	0			
		SE		v	3.7	v			
	1999	5	44	33	32	8	9.1		
	1,,,,	13	9	1	3	0	7.1		
		Uncut	3	i	Õ	Ŏ			
		SE		•	9.7	v			
Birdsfoot trefoil	1998	5	25	17	11	11	8.0		
Ditusioot ticion	1570	13	3	7	1	1	0.0		
		Uncut	5	17	i	0			
		SE		1,	7.6	v			
	1999	5	42	69	28	14	10.6		
	1///	13	38	6	4	0	10.0		
		Uncut	12	5	i	ĭ			
		SE	12	3	9.6	1			

 $<sup>\</sup>dagger$  SE by year and legume species for comparison of two N fertilizer rate means at the same or different sward height.

<sup>‡</sup> SE by year and legume species for comparison of two sward height means at the same N fertilizer rate.




Fig. 1. Landscape position and N fertilizer rate effects on survival of a legume mixture interseeded into a cool-season grass pasture at Rhodes, IA. Values represent overall survival of red clover, alfalfa, and birdsfoot trefoil on 15 Aug. 1998 and 1999. Standard errors for comparison of landscape position means at the same or different N rate were 25.9 in 1998 and 20.1 in 1999. Standard errors for comparison of N rate means at the same landscape position were 25.8 in 1998 and 21.8 in 1999.

Table 5. Analysis of variance and means for leaf area index of pasture swards measured at Rhodes, IA, in 1998 and 1999. Sources of variation included two landscape positions (L), four N fertilizer rates (N), three sward heights (H), and four replications (R). Leaf area index was measured in grass swards at two sampling dates each year before accumulation of appreciable leaf area by legume seedlings.

			Leaf area index						
Source		df	1	998	1999				
	F test		28 May	22 June	22 June	16 July			
L	MS(L)/MS[R(L)]	1	11.9	75.7***	2.1	12.2**			
R(L)	~ ( )- ~ L ( )J	6	3.4	1.8	1.5	0.5			
N	$MS(N)/MS[R \times N(L)]$	3	5.4***	2.1†	3.9**	9.2***			
$L \times N$	$MS(L \times N)/MS[R \times N(L)]$	3	0.1	0.2	1.4	0.7			
$\mathbf{R} \times \mathbf{N}(\mathbf{L})$	. , , , , , , , , , , , , , , , , , , ,	18	0.6	0.7	0.6	0.3			
H	$MS(H)/MS[R \times N \times H(L)]$	2	43.8***	114.1***	114.9***	91.2***			
$L \times H$	$MS(L \times H)/MS[R \times N \times H(L)]$	2	0.5	2.1†	1.8	1.3			
$N \times H$	$MS(N \times H)/MS[R \times N \times H(L)]$	6	0.7	0.5	0.8	0.8			
$L \times N \times H$	$MS(L \times N \times H)/MS[R \times N \times H(L)]$	6	0.6	0.2	1.8†	0.7			
$\mathbf{R} \times \mathbf{N} \times \mathbf{H}(\mathbf{L})$	7 2	36	0.4	0.7	0.8	0.9			
Landscape position	Summit		3.0	4.7	2.8	4.0			
	Backslope		2.3	2.9	2.5	3.3			
	SE‡		0.38	0.27	0.25	0.14			
Sward height, cm	5		1.6	2.0	0.9	2.2			
9 /	13		2.5	3.5	2.4	3.3			
	Uncut		3.9	5.8	4.6	5.5			
	SE§		0.16	0.21	0.22	0.24			
Nitrogen, kg ha-1	0		2.2	3.4	2.2	2.9			
5 , 8	22		2.4	3.7	2.4	3.4			
	44		3.0	4.0	2.9	4.0			
	89		3.2	4.0	3.0	4.3			
	SE¶		0.22	0.34	0.22	0.16			

<sup>\*\*</sup> Denotes significance at the 0.01 probability level.

#### **DISCUSSION**

Seedling survival rather than emergence limits establishment of legumes on summit positions in pastures. Repeated defoliation of the swards to 5 cm improved legume survival on summits; however, relative to backslopes, survival on summits remained low. In swards cut at 5 cm, overall legume survival on summits was less than on backslopes by 69 plants m<sup>-2</sup> in 1998 and 173 plants m<sup>-2</sup> in 1999 (Table 3). Overall legume survival was greater on backslopes than on summits in swards cut at 5 and 13 cm despite similar legume emergence among these positions at these sward heights. Only when swards remained uncut was emergence limited on summits relative to backslopes. Emergence on summits was less than on backslopes by 686 plants m<sup>-2</sup> in uncut swards fertilized at 22 kg ha<sup>-1</sup> N in 1998 and 1003 plants m<sup>-2</sup> in uncut swards at 0 kg ha<sup>-1</sup> N in 1999 (Table 2). Our results support those of Harmoney et al. (2001), who found forage mass of legumes in pastures was 179 kg ha<sup>-1</sup> greater on backslopes than on summits.

Reduced grass competition improved legume survival on backslope positions. In 1998 and 1999, respectively, 132 and 234 plants m<sup>-2</sup> survived in swards cut at 5 cm compared with 21 and 31 plants m<sup>-2</sup> in uncut swards (Table 3). By repeatedly cutting the sward to 5- and 13-cm heights, grass LAI was reduced throughout June and July, months during which legume seedlings were early in development (Table 5). Greater leaf area from grasses limits establishment of legumes in pastures by limiting the photosynthetic photon flux (Trott et al., 1988) and altering the ratio of red to far-red wavelengths (Ballaré et al., 1990) that reaches seedlings at ground

Overall legume survival may have been greater among plots cut at 5 cm on backslopes in 1999 than in 1998 because of reduced soil fertility and competition from existing grasses. We did not measure soil nutrient status, but as mentioned previously, the pasture in which this experiment was established was fertilized and moderately grazed before 1998. During 1998, however, the sod adjacent to the plots was mowed and raked but not fertilized. In 1999, the plots were established ≈3 m away from the 1998 plots. The abundance of litter before seeding was probably similar among 1998 and 1999 plots; however, soil fertility levels may have been lower in 1999 plots. Under conditions of reduced competition and soil fertility, addition of N at 22 kg ha<sup>-1</sup> also may have positively benefited legume survival within the 5-cm cut plots without increasing grass competition or diminishing nodule formation in 1999 (Table 4).

Reducing competition from grass before and after seeding of legumes is necessary to support legume emergence if planting is delayed following N fertilization. Overall legume emergence was not affected by sward height in 1998 but was reduced by 61 to 98% in swards left uncut compared with swards cut at 5 cm in 1999 (Table 2). Wet field conditions delayed planting until 30 d after N application in 1999. Within uncut swards and swards fertilized at 44 to 89 kg ha<sup>-1</sup> N, light availability most likely limited legume emergence. Increased etiolation of legume seedlings as N fertilization rates increased under 13-cm and uncut swards was observed.

<sup>\*\*\*</sup> Denotes significance at the 0.001 probability level.

<sup>†</sup> Denotes significance at the 0.100 probability level.

<sup>‡</sup> SE for comparison of landscape position means by measurement date. § SE for comparison of sward height means by measurement date.

<sup>¶</sup> SE for comparison of N fertilizer rate means by measurement date.

Overall legume survival, however, did not appear to be affected by the delay in planting as survival was similar among uncut swards and swards fertilized at 44 to 89 kg ha<sup>-1</sup> N in 1998 and 1999.

Nitrogen fertilization further escalated competition from grass, and the negative effect on legume establishment was greater when sward heights were not minimized. Nitrogen fertilization is known to favor grasses over legumes and is not a recommended practice to maintain legumes in pastures (Heichel, 1985). When N fertility is high, nodule formation and N<sub>2</sub> fixation of legumes is inhibited (Heichel, 1985), and grasses compete more successfully with legumes (Schwinning and Parsons, 1996). We found that N fertilizer did not affect emergence in 1998 or 1999 when swards were repeatedly cut at 5 cm before and after seeding. Nitrogen fertilizer, however, reduced legume emergence in 13-cm and uncut swards. Overall legume survival was also affected less by N fertilization in 5-cm swards than in 13-cm and uncut swards.

Competition from grass is likely greater on summits than on backslopes because above-ground dry matter production of grass tends to be greater on summits than on backslopes (Harmoney et al., 2001). From 0 to 44 kg ha<sup>-1</sup> N, overall legume survival was on average greater on backslopes than on summits by 64 plants m<sup>-2</sup> in 1998 and 105 plants m<sup>-2</sup> in 1999. Addition of 89 kg ha<sup>-1</sup> N was required to reduce overall legume survival on backslopes to similar densities of survival on summits (Fig. 1). Growth rate of the sward appears to be greater on summits compared with backslopes. Leaf area index of the sward was initially greater and increased at a greater rate on summits than on backslopes from 28 May 1998 to 22 June 1998 and from 22 June 1999 to 16 July 1999.

Though we did not conduct this study across multiple locations or hillslopes, we expect competition from grasses to be greater on summits and overall legume survival to be greater on backslopes among other pastures typical of southeastern Iowa, and that these results would occur regardless of aspect. Aspect did not affect legume abundance or the production of grass dry matter in a previous experiment conducted in southeastern Iowa pastures (Harmoney et al., 2001).

Red clover had the greatest emergence and survival of the three legumes when sown together at similar rates in these cool-season grass pastures. Red clover was the predominant legume regardless of landscape position, N fertilizer rate, or sward height. Alfalfa and birdsfoot trefoil contributed significantly to overall legume survival when swards were cut at 5 cm and N fertilizer rates were 0 to 22 kg ha<sup>-1</sup>. In sown mixtures, the diversity of seedlings that establish may be lower in N-fertilized

grasslands because accumulated litter and lower light penetration inhibits germination and/or survival of seedlings (Tilman, 1993). Previous research has shown that red clover is affected less by low availability of light than alfalfa and birdsfoot trefoil (Gist and Mott, 1957). It remains unknown, however, as to how emergence and survival of these legumes might be affected were they to be sown as a mixture in pastures with different limiting conditions.

Our study shows that legume establishment is limited by seedling survival on summit positions in pastures. Establishment of legumes is successful on backslopes because of less competition from grass. Increased sward height and N fertility enhance grass competitiveness, thus reducing both emergence and survival of legumes in pastures. We recommend legumes be seeded on backslope landscape positions in pastures and that N fertilizer not be applied, and sward heights be reduced before seeding.

#### **REFERENCES**

- Ballaré, C.L., A.L. Scopel, and R.A. Sánchez. 1990. Far-red radiation reflected from adjacent leaves: An early signal of competition in plant communities. Science (Washington, DC) 24:329–332.
- Cuomo, G.J., D.G. Johnson, and W.A. Head, Jr. 2001. Interseeding kura clover and birdsfoot trefoil into existing cool-season grass pastures. Agron. J. 93:458–462.
- Gist, G.R., and G.O. Mott. 1957. Some effects of light intensity, temperature, and soil moisture on the growth of alfalfa, red clover, and birdsfoot trefoil seedlings. Agron. J. 49:33–36.
- Groya, F.L., and C.C. Sheaffer. 1981. Establishment of sod-seeded alfalfa at various levels of soil moisture and grass competition. Agron. J. 73:560–565.
- Harmoney, K.R., K.J. Moore, E.C. Brummer, C.L. Burras, and J.R. George. 2001. Spatial legume composition and diversity across seeded landscapes. Agron. J. 93:992–1000.
- Heichel, G.H. 1985. Symbiosis: Nodule bacteria and leguminous plants. p. 64–71. *In* D.S. Metcalfe (ed.) Forages: The science of grassland agriculture. 4th ed. Iowa State Univ. Press, Ames.
- Heichel, G.H., and K.I. Henjum. 1991. Dinitrogen fixation, N transfer, and productivity of forage legume-grass communities. Crop Sci. 31:202–208.
- McIntosh, M.S. 1983. Analysis of combined experiments. Agron. J. 75:153–155.
- Schwinning, S., and A.J. Parsons. 1996. Analysis of the coexistence mechanisms for grasses and legumes in grazing systems. J. Ecol. 84:799–813.
- Sleugh, B., K.J. Moore, J.R. George, and E.C. Brummer. 2000. Binary legume-grass mixtures improve forage yield, quality, and seasonal distribution. Agron. J. 92:24–29.
- Taylor, R.W., and D.W. Allinson. 1983. Legume establishment in grass sods using minimum-tillage seeding techniques without herbicide application: Forage yield and quality. Agron. J. 75:167–172.
- Tilman, D. 1993. Species richness of experimental productivity gradients: How important is colonization limitation? Ecology 74:2179–2191
- Trott, J.O., K.J. Moore, V.L. Lechtenberg, and K.D. Johnson. 1988. Light penetration through tall fescue in relation to canopy biomass. J. Prod. Agric. 1:137–140.
- Van Soest, P.J. 1982. Nutritional ecology of the ruminant. O & B Books, Corvallis, OR.