University of Nebraska - Lincoln DigitalCommons@University of Nebraska - Lincoln

Faculty Papers and Publications in Animal Science

Animal Science Department

1-1-2007

Effect of Increasing Levels of Corn Bran on Milk Yield and Composition

B. N. Janicek University of Nebraska - Lincoln

Paul J. Kononoff University of Nebraska - Lincoln, pkononoff2@unl.edu

A. M. Gehman University of Nebraska - Lincoln, agehman2@unl.edu

Dakota Gold Research Association, Sioux Falls, SD

M. L. Gibson Dakota Gold Research Association, Sioux Falls, SD

Follow this and additional works at: http://digitalcommons.unl.edu/animalscifacpub



Part of the Animal Sciences Commons

Janicek, B. N.; Kononoff, Paul J.; Gehman, A. M.; Karges, K.; and Gibson, M. L., "Effect of Increasing Levels of Corn Bran on Milk Yield and Composition" (2007). Faculty Papers and Publications in Animal Science. Paper 604. http://digitalcommons.unl.edu/animalscifacpub/604

This Article is brought to you for free and open access by the Animal Science Department at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Papers and Publications in Animal Science by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Short Communication: Effect of Increasing Levels of Corn Bran on Milk Yield and Composition¹

B. N. Janicek,* P. J. Kononoff,*² A. M. Gehman,* K. Karges,† and M. L. Gibson†
*Department of Animal Science, University of Nebraska, Lincoln 68583-0908
†Dakota Gold Research Association, Sioux Falls, SD 57104

ABSTRACT

Thirty-nine lactating Holstein cows (23 multiparous and 16 primiparous) were randomly assigned to 1 of 3 dietary treatments in a crossover design. Dietary treatments differed by the proportion of corn bran [10, 17.5, and 25% dry matter (DM); designated as low, medium, and high] replacing corn silage and alfalfa. The corn bran coproduct contained 8.2% moisture and 12.9% crude protein, 30.4% neutral detergent fiber (NDF), and 45.0% nonfiber carbohydrate, 9.9% ether extract, and 0.70% P (DM basis). The low treatment consisted of 15.8% NDF from forage (fNDF) and 33.1% total NDF; the medium treatment consisted of 12.9% fNDF and 32.5% total NDF; and the high diet contained 9.9% fNDF and 31.8% total NDF. Dry matter intake was not affected by treatment. The percent milk fat decreased by 0.26% with the inclusion of corn bran from 10 to 25% of the diet DM, but total milk fat yield was not affected. In comparison, corn bran increased yield of milk protein 0.12 kg/d when bran increased from 10 to 25% of the diet DM. Total milk yield tended to increase when bran increased from 10 to 25% of the diet DM, but no differences were observed on 3.5% fat-corrected milk. Lastly, feed conversion significantly improved with increasing inclusion: 1.39, 1.39, and 1.55 \pm 0.05 kg of milk/kg of DMI for low, medium, and high, respectively. Observed effects were likely due to the increase in energy intake associated with increasing levels of corn bran.

Key words: corn bran, ethanol coproduct, milk, lactation

The recently passed Energy Policy Act (EPACT, 2005) mandates up to 7.5 billion gallons of renewable fuel to

be used by 2012 and will result in a dramatic increase in the availability of corn milling coproducts. The inclusion of coproducts such as corn distillers grains with solubles (DDGS) into dairy diets has been demonstrated to maintain milk production (Schingoethe et al.. 1999) but are usually fed in limited amounts because of the potential to negatively affect milk fat percentage (Leonardi et al., 2005). Traditional corn-ethanol production employs a system in which the whole corn kernel is ground, cooked, and then fermented to produce ethanol. An alternative to this method is separating the kernel into its 3 major components, namely bran, germ, and endosperm, prior to fermentation. These components are then spared the fermentation processes and may be used for animal feed. Compared with traditional DDGS, corn bran is similar in NDF but lower in CP. Because corn bran is not exposed to the fermentation process, it is generally a better quality nutrient source. The objective of this research was to evaluate the effects of increasing levels of a corn bran coproduct on milk production and composition.

Thirty-nine lactating Holstein cows (23 multiparous and 16 primiparous) were randomly assigned to 1 of 3 dietary treatments in a 3-period crossover design. The experiment was conducted in 2 phases, with 20 animals in phase 1 and 19 animals in phase 2. Cows were housed in individual stalls and milked at 0730 and 1930 h. Cows were fed at 0800 h for ad libitum consumption to allow for approximately 5% refusal. During each of the 21-d periods, cows were offered 1 of 3 rations (Table 1) that differed by the concentration of corn bran that replaced alfalfa hay, alfalfa haylage, and corn silage. Dietary treatments were 1) low, 10% DM bran, 2) medium, 17.5% DM bran, and 3) high, 25% DM bran. The experimental cows were cared for according to the guidelines stipulated by the University of Nebraska-Lincoln Animal Care and Use Committee.

Samples of feed were collected weekly and composited by experimental period. Collected samples were immediately frozen (-20°C) and stored for further analysis. Before analysis samples were dried at 60°C in a forcedair oven and ground (1-mm screen; Wiley mill, Arthur H. Thomas Co., Philadelphia, PA) and analyzed as de-

Received April 11, 2007. Accepted May 4, 2007.

¹A contribution of the University of Nebraska Agricultural Research Division, supported in part by funds provided through Hatch Act. Additional support was provided by Dakota Gold Research Association, Sioux Falls, SD.

²Corresponding author: pkononoff2@unl.edu

4314 JANICEK ET AL.

Table 1. Chemical analysis and diet ingredients

Ingredient (% of DM)			
	Low	Medium	High
Corn silage	23.0	18.8	14.5
Alfalfa hay	8.7	7.1	5.47
Alfalfa haylage	8.7	7.1	5.47
Corn grain	14.7	14.7	14.7
Bran ²	10.2	17.5	25.0
Soybean hulls	12.6	12.6	12.6
Soybean meal	6.9	6.9	6.9
Soypass ³	5.7	5.7	5.7
Limestone	1.7	1.7	1.7
Cottonseed	6.5	6.5	6.5
Sodium bicarbonate	0.70	0.70	0.70
Magnesium oxide	0.17	0.17	0.17
Salt	0.11	0.11	0.11
Vitamin E	0.02	0.02	0.02
$\mathrm{Selplex}^4$	0.01	0.01	0.01
Vitamin A, D, E ⁵	0.13	0.13	0.13
Trace mineral ⁶	0.04	0.04	0.04
Chemical composition			
CP, % of DM	18.4	18.2	18.1
RUP, % of DM	6.6	6.5	6.4
NDF, % of DM	33.1	32.5	31.8
NFC,7 % of DM	38.3	39.0	39.7
Ether extract, % of DM	4.9	5.3	5.7
Phosphorus	0.34	0.38	0.42
Particle size			
>19.0 mm	5.7	4.8	4.5
19.0 to 8.0 mm	26.3	23.7	21.4
8.0 to 1.18 mm	42.9	44.4	43.5
<1.18 mm	25.2	27.1	30.7

¹Rations containing increasing levels of corn bran; low (10% DM), medium (17.5% DM), and high (25% DM).

scribed by Kononoff et al. (2006). For determination of NDF, α -amylase and sodium sulphite were used (Mertens, 2002). The Penn State Particle Separator was used to measure particle size for all rations (Kononoff et al., 2003). Body weight was measured on the last 2 d of each period, but BCS (1 - 5 scale) was measured on the last day of each period. Body condition score was measured by a single trained individual and differed from that described by Wildman et al. (1982) because it was reported to the quarter point. Milk production was measured daily, and milk samples were collected on d 20 and 21 of each period during the AM and PM milking and preserved using 2-bromo-2-nitropropane-1,3 diol. Milk samples were analyzed for fat and true protein (AOAC, 1990) using a B2000 Infrared Analyzer (Bentley Instruments, Chaska, MN) by Heart of America DHIA (Manhattan, KS). Daily DMI and milk yield were averaged weekly.

Milk production, milk composition, intake, and BCS data were analyzed using the MIXED procedure of SAS (Version 9.1, SAS Institute Inc., Cary, NC) according to the following model: $y_{ijklm} = \mu + b_{ij} + \rho_k + \gamma_l + \lambda_m + \pi_n + e_{ijklm}$, where μ is the general mean, b_{ij} is the random effect of the jth cow within the ith sequence, ρ_k is the effect of the kth period, γ_l is the fixed effect of the effect of the lth diet, λ_m is the fixed effect of the mth parity, π_n is the fixed effect of the nth experimental phase, and e_{ijk} reflects the random error in the measurement of the response. Linear, quadratic, and cubic orthogonal contrasts were tested using the CONTRAST statement of SAS.

The corn bran coproduct (Dakota Gold Marketing, Sioux Falls, SD) contained 8.2% moisture and 12.9% CP, 30.4% NDF, and 45.0% NFC, 9.9% ether extract, and 0.70% P when expressed on a DM basis. This chemical analysis outlines the major differences in chemical composition that may exist amount corn milling coproducts. Specifically, compared with the corn bran coproduct, DDGS typically contain similar levels of NDF (38.8%) and fat (10.0%) but higher levels of CP (29.7%; NRC, 2001).

Diets were formulated to contain 3 levels of the bran, which replaced a portion of the forages. The chemical composition of the diets fed were similar (Table 1), containing approximately 18% CP, but increasing the levels of bran resulted in a modest decrease in total diet NDF: 33.1, 32.5, and 31.8% DM for low, medium, and high treatments, respectively. The concentrations of ether extract and NFC were also different (ether extract = 4.9, 5.3, 5.7% DM and NFC = 38.3, 39.0, and 39.7% DM for low, medium, and high, respectively). The diet P content increased from 0.34, 0.38, and 0.42% as corn bran was included in the diet from 10, 17.5, and 25% of the diet DM. The TMR particle size analysis is also presented in Table 1. Ration particle size reflected the amount of bran included in the treatments. Increasing inclusion of bran reduced the proportion of material on the 19.0-mm screen (5.7, 4.8, 4.5%), but increased the proportion retained on the pan (<1.18mm; 25.2, 27.1, 30.7%).

Increasing the level of bran did not affect DMI, which averaged 23.8 ± 0.73 kg/d across treatments (Table 2). Milk yield tended (P = 0.07) to increase linearly with increasing concentrations of bran: 32.7, 33.4, 35.8 ± 1.30 kg/d for low, medium, and high, respectively (Table 2). As a consequence of lack of effect on DMI and effect on milk yield, feed conversion significantly improved with increasing inclusion: 1.39, 1.39, and 1.55 ± 0.05 kg of milk/kg of DMI for low, medium, and high, respectively. The observation that the feeding of corn milling coproducts may result in increased milk yield but not intake is in contrast to that of Owen and Larson (1991).

²Dakota Gold Marketing Inc., Sioux Falls, SD.

³LignoTech, Overland Park, KS.

⁴Alltech Inc., Nicholasville, KY.

 $^{^5 \}rm Formulated$ to supply approximately 120,000 IU/d of vitamin A, 24,000 IU/d of vitamin D, and 800 IU/d of vitamin E.

 $^{^6}Formulated$ to contained 1.0% Ca, 0.50 % P, 0.36% Mg, 1.3% K. 7NFC calculated by difference: 100 – (%NDF + % CP + % ether extract + % ash).

Table 2. The effect of feeding corn bran on milk production and composition of Holstein dairy cows

	$Treatment^1$				P-value	
Item	Low	Medium	High	SEM^2	Linear	Quadratic
DMI, kg/d	23.6	24.2	23.6	0.73	0.91	0.54
Energy intake, ³ Mcal/d	38.4	40.2	41.1	1.30	0.15	0.76
Milk yield, kg/d	32.7	33.4	35.8	1.30	0.07	0.59
Fat, %	3.58	3.50	3.32	0.10	0.06	0.70
Protein, %	2.99	2.97	3.06	0.04	0.17	0.23
MUN, mg/dL	14.8	14.9	13.9	0.34	0.05	0.14
3.5% FCM	32.9	33.57	34.30	1.29	0.40	0.97
Lactose, %	4.78	4.75	4.82	0.05	0.49	0.39
Fat yield, kg/d	1.16	1.18	1.16	0.05	0.94	0.75
Protein yield, kg/d	0.97	1.00	1.09	0.04	0.02	0.44
Feed conversion, kg of milk/kg of DMI	1.39	1.39	1.55	0.05	0.03	0.19
BCS	3.29	3.27	3.27	0.05	0.76	0.80

 $^{^1\}mathrm{Rations}$ containing increasing levels of corn bran low (10% DM), medium (17.5% DM), and high (25% DM).

In that study, DDGS replaced primarily soybean meal and diets were of low protein content. These authors suggested that intakes were increased by feeding DDGS because a greater amount of amino acids were required to maintain maximal milk production. Thus it is possible that in the current experiment intakes were similar across treatments because the coproduct replaced forage, and based on the NRC (2001) model, experimental diets were formulated to be higher in CP and adequate in MP. This suggestion is further supported because a lack of effect of DMI has also been observed when DDGS replace forage in the form of alfalfa haylage (Clark and Armentano, 1993).

A common field concern related to feeding many corn milling coproducts is the high fat content. It is now generally understood that a buildup of the unsaturated fatty acid, linoleic acid, in the rumen may lead to events that can cause milk fat depression (Baumgard et al., 2000). Generally speaking, coproducts are high in fat and are a rich source of linoleic acid. Although bran resulted in a tendency (P = 0.06) for a reduction of the concentration of milk fat, total milk yield was increased, so there were no differences in total milk fat yield. This response is similar to Leonardi et al. (2005), who increased the proportion of DDGS in a ration from 0 to 15% DM. Practically, these data also outline the importance of understanding effects on milk composition and total milk yield before implicating corn milling coproducts as a cause for low milk fat tests. In comparison to percent fat, differences in the concentration of milk protein were not observed (mean = $3.00 \pm 0.04\%$). However, total protein yield was significantly increased for cows consuming more bran (0.97, 1.00, and 1.09 kg/ d for low, medium, and high, respectively) due to greater milk yield. This response is possibly due to improved energy status of cows consuming bran, as energy intake is positively correlated with milk protein synthesis (Grieve et al., 1986). The diets containing bran may have provided more ruminally fermentable OM resulting in higher amounts of propionate for use in gluconeogenesis and amino acids being spared for protein synthesis (Harvatine et al., 2002). No statistical differences were observed on 3.5% FCM with increasing levels of bran.

Compared with other studies evaluating health and reproduction (Windig et al., 2005), the current study contained a small number of animals and should be interpreted with caution. During the second phase of the experiment, 3 animals were diagnosed with displaced abomasums, and data from these animals were removed from the data set. Two animals were diagnosed while consuming the high treatments, and one was diagnosed while consuming the medium treatment. The reason for these observations is unclear but may have been a result of a reduction in effective fiber when the bran replaced forage. The NRC (2001) recommends dairy diets contain at least 25% NDF with at least 19% from forage (fNDF) and an upward adjustment in ration NDF content if fNDF content is reduced. In the current experiment, the low treatment consisted of 15.8% fNDF and 33.1% NDF, which is in the recommended range of minimum total and forage NDF. In comparison, the high diet contained 9.9% fNDF and 31.8% NDF, which is below recommended levels. Practically, health results of the current experiment suggest that the bran coproduct is a feedstuff with high nutrient quality but should be included into dairy diets balanced to contain adequate levels of effective fiber.

²Highest SE of treatment means is shown.

³According to NRC (2001).

4316 JANICEK ET AL.

REFERENCES

- Association of Official Analytical Chemists. 1990. Official Methods of Analysis. 15th ed. AOAC, Arlington, VA.
- Baumgard, L. H., B. A. Corl, D. A. Dwyer, A. Saebo, and D. E. Bauman. 2000. Identification of the conjugated linoleic acid isomer that inhibits milk fat synthesis. Am. J. Physiol. 278:R179–R184.
- Clark, P. W., and L. E. Armentano. 1993. Effectiveness of neutral detergent fiber in whole cottonseed and dried distillers grains compared with alfalfa haylage. J. Dairy Sci. 76:2644–2650.
- Energy Policy Act (EPACT). 2005. Public Law 109-58. http://www.frwebgate.access.gpo.gov/cgi-bin/getdoc.cgi?dbname=109_cong_public_laws&docid=f:publ058.109 Accessed Apr. 2, 2006.
- Grieve, D. G., S. Korver, Y. S. Rijpkema, and G. Hof. 1986. Relationship between milk composition and some nutritional parameters in early lactation. Livest. Prod. Sci. 14:239–254.
- Harvatine, D. I., J. L. Firkins, and M. L. Eastridge. 2002. Whole linted cottonseed as a forage substitute fed with ground or steamflaked corn: Digestibility and performance. J. Dairy Sci. 85:1976–1987.
- Kononoff, P. J., A. J. Heinrichs, and D. A. Buckmaster. 2003. Modification of the Penn State forage and TMR separator and the effects of moisture content on its measurements. J. Dairy Sci. 86:1858–1863.

- Kononoff, P. J., S. Ivan, W. Matzke, R. J. Grant, R. Stock, and T. J. Klopfenstein. 2006. Milk production of dairy cows fed a wet corn gluten feed during the dry period and lactation. J. Dairy Sci. 89:2608–2617.
- Leonardi, C., S. Bertics, and L. E. Armentano. 2005. Effect of increasing oil from distillers grains or corn oil on lactation performance. J. Dairy Sci. 88:2820–2827.
- Mertens, D. R. 2002. Gravimetric determination of amylase-treated neutral detergent fiber in feeds using refluxing in beakers or crucibles: Collaborative study. J. AOAC 85:1217–1240.
- NRC. 2001. Nutrient Requirements of Dairy Cattle. 7th rev. ed. Natl. Acad. Sci., Washington, DC.
- Owen, F. G., and L. L. Larson. 1991. Corn distillers dried grains versus soybean meal in lactation diets. J. Dairy Sci. 74:972–979.
- Schingoethe, D. J., M. J. Brouk, and C. P. Birkelo. 1999. Milk production and composition from cows fed wet corn distillers grains. J. Dairy Sci. 82:574–580.
- Wildman, E. E., G. M. Jones, P. E. Wagner, H. F. Troutt, and T. N. Lesch. 1982. A dairy cow body condition scoring system and its relationship to selected production characteristics. J. Dairy Sci. 65:495–501.
- Windig, J. J., M. P. L. Calus, and R. F. Veerkamp. 2005. Influence of herd environment on health and fertility and their relationship with milk production. J. Dairy Sci. 88:335–347.