

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

USGS Staff -- Published Research

US Geological Survey

2005

Geographic Variation, Genetic Structure, and Conservation Unit Designation in the Larch Mountain Salamander (*Plethodon larselli*)

R. Steven Wagner
Central Washington University

Mark P. Miller
Utah State University, markperrymiller@gmail.com

Charles M. Crisafulli
US Forest Service

Susan M. Haig
U.S. Geological Survey, Susan_Haig@usgs.gov

Follow this and additional works at: <https://digitalcommons.unl.edu/usgsstaffpub>

Wagner, R. Steven; Miller, Mark P.; Crisafulli, Charles M.; and Haig, Susan M., "Geographic Variation, Genetic Structure, and Conservation Unit Designation in the Larch Mountain Salamander (*Plethodon larselli*)" (2005). *USGS Staff -- Published Research*. 674.
<https://digitalcommons.unl.edu/usgsstaffpub/674>

This Article is brought to you for free and open access by the US Geological Survey at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in USGS Staff -- Published Research by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Geographic variation, genetic structure, and conservation unit designation in the Larch Mountain salamander (*Plethodon larselli*)

R. Steven Wagner, Mark P. Miller, Charles M. Crisafulli, and Susan M. Haig

Abstract: The Larch Mountain salamander (*Plethodon larselli* Burns, 1954) is an endemic species in the Pacific northwestern United States facing threats related to habitat destruction. To facilitate development of conservation strategies, we used DNA sequences and RAPDs (random amplified polymorphic DNA) to examine differences among populations of this species. Phylogenetic analyses of cytochrome *b* revealed a clade of haplotypes from populations north of the Columbia River derived from a clade containing haplotypes from the river's southwestern region. Haplotypes from southeastern populations formed a separate clade. Nucleotide diversity was reduced in northern populations relative to southern populations. These results were corroborated by analyses of RAPD loci, which revealed similar patterns of clustering and diversity. Network analyses suggested that northern populations were colonized following a range expansion mediated by individuals from populations located southwest of the river. Changes in the Columbia River's location during the Pliocene and Pleistocene likely released distributional constraints on this species, permitting their northern range expansion. Based on the barrier presented by the Columbia River's present location and differences in haplotype diversity and population structure observed between northern and southern populations, we suggest that designation of separate management units encompassing each region may assist with mitigating different threats to this species.

Résumé : La salamandre de Larch Mountain (*Plethodon larselli* Burns, 1954) est une espèce endémique de la région du nord-ouest pacifique des États-Unis qui est menacée par la destruction de son habitat. Les séquences d'ADN et les RAPD (amplifications aléatoires d'ADN polymorphe) nous ont servi à étudier les différences entre les populations de l'espèce, afin de faciliter la mise au point de stratégies de conservation. Les analyses phylogéniques du cytochrome *b* indiquent l'existence d'un clade d'haplotypes provenant des populations au nord du Columbia dérivés d'un clade contenant des haplotypes provenant de la région du sud-ouest du fleuve. Les haplotypes de la région sud-est forment un clade séparé. La diversité des nucléotides est plus faible dans les populations du nord que dans celles du sud. Les analyses des locus RAPD confirment les résultats et présentent des patrons similaires de regroupement et de diversité. Des analyses de réseau laissent croire que les populations du nord ont fait leur colonisation après une extension d'aire réalisée par des individus provenant des populations établies au sud-ouest du fleuve. Les déplacements du lit du Columbia durant le pliocène et le pléistocène ont réduit certaines contraintes biogéographiques pour l'espèce et ont permis l'expansion d'aire vers le nord. Compte tenu de la barrière que forme le Columbia dans son emplacement actuel et des différences de diversité des haplotypes et de structure de population entre les populations du nord et du sud, nous proposons l'établissement d'unités de gestion distinctes pour chacune des régions pour atténuer les diverses menaces auxquelles fait face cette espèce.

[Traduit par la Rédaction]

Introduction

Phylogeographic studies aid in identifying historical barriers to dispersal and gene flow. Such analyses can substantially contribute to our understanding of the effects of natural and anthropogenic impacts on habitat and population

fragmentation (Avice 1994). Amphibians often have specific ecological requirements and low dispersal rates that make them susceptible to fragmentation by historical and current processes (Blaustein et al. 1994). Indeed, genetic studies of amphibians frequently reveal substantial cryptic genetic diversity attributable to the influence of vicariant events

Received 23 September 2004. Accepted 10 March 2005. Published on the NRC Research Press Web site at <http://cjb.nrc.ca> on 6 May 2005.

R.S. Wagner.¹ Genetics Program, Oregon State University, Corvallis, OR 97331, USA and USGS Forest and Rangeland Ecosystem Science Center, 3200 SW Jefferson Way, Corvallis, OR 97331, USA.

M.P. Miller.² Department of Biology, 5305 Old Main Hill, Utah State University, Logan, UT 84322-5305, USA.

C.M. Crisafulli. US Forest Service Pacific Northwest Research Station, Mount St. Helens National Volcanic Monument, 42218 NE Yale Bridge Road, Amboy, WA 98601, USA.

S.M. Haig. USGS Forest and Rangeland Ecosystem Science Center, 3200 SW Jefferson Way, Corvallis, OR 97331, USA.

¹Present address: Department of Biological Science, Central Washington University, Ellensburg, WA 98926, USA.

²Corresponding author (e-mail: Mark.Miller@usu.edu).


(Good and Wake 1992; Highton 1995; Jockusch 1996; Tilley and Mahoney 1996; Miller et al. 2005). Therefore, to aid amphibian conservation efforts, it is important to understand the role of fragmentation in population differentiation and implement management plans that preserve within-species genetic diversity.

The Larch Mountain salamander (*Plethodon larselli* Burns, 1954) is a completely terrestrial species associated with a variety of habitat types in the Pacific northwestern United States (Crisafulli 1999). This species is considered to be one of the rarest amphibian species in the Pacific Northwest because of its restricted range along a narrow corridor of the Columbia River Gorge and patchy distribution in the northern Oregon Cascade Range and south and central Washington Cascade Range (Howard et al. 1983; Kirk 1983; Herrington and Larsen 1985; Crisafulli 1999). As of 2004, there are only 137 extant sites. They occur in isolated patches of forested, steep talus within the Columbia River Gorge (Burns 1954, 1962, 1964; Herrington and Larsen 1985) and at rock outcrops, cave entrances, and a wide variety of forest and shrub habitats in the Cascade Mountains (Aubry et al. 1987; Darda and Garvey-Darda 1995; Crisafulli 1999). These regions have a substantial disturbance history primarily because of volcanism, glaciation, and flooding during the late Pliocene Epoch (McGroder 1991) and the Quaternary Period (Harris 1988) that could influence contemporary patterns of genetic diversity and population genetic structure in Larch Mountain salamanders.

Major threats to this species include habitat loss and population fragmentation caused by logging, recreational activities, and housing development (Herrington and Larsen 1985). Larch Mountain salamanders are considered a species of concern by federal and state agencies. They are listed by Oregon State as a Sensitive species, Vulnerable status, and by Washington State as a Sensitive species. In Oregon and Washington the species receives special management consideration from the US Forest Service as a Sensitive species and the US Bureau of Land Management as an Assessment species (US Forest Service and US Bureau of Land Management 1994). The Nature Conservancy has classified this species as both globally imperiled and subnationally imperiled. Furthermore, the species is listed in the *IUCN Red List* (International Union for the Conservation of Nature 2004) as Near Threatened. Thus, given contemporary management concerns over the Larch Mountain salamander, we examined population differentiation and genetic structure (both nuclear and mitochondrial) of 15 Larch Mountain salamander populations throughout their range. Furthermore, we evaluate possible historical biogeographical factors that may also influence Larch Mountain salamander genetic structure.

We used mtDNA cytochrome *b* sequences and RAPD (random amplified polymorphic DNA) markers to assess patterns of genetic diversity across the salamander's range. Cytochrome *b* sequences have been used in a wide variety of taxa for designating conservation units (e.g., Baker et al. 1995; Lento et al. 1997; Mundy et al. 1997; Castilla et al. 1998; Walker et al. 1998) and in several salamander species to infer intraspecific phylogenies (e.g., Hedges et al. 1992; Moritz et al. 1992; Tan and Wake 1995; Jackman et al. 1997). The RAPD technique, a procedure that permits examination of large numbers of segregating, putative nuclear loci

Fig. 1. Map showing the locations of Larch Mountain salamander (*Plethodon larselli*) populations examined in this study.



from the genome, has been frequently used in vertebrate conservation studies (e.g., Haig et al. 1994; Fleischer et al. 1995; Haig et al. 1996; Kimberling et al. 1996; Nusser et al. 1996; Haig et al. 1997; Haig 1998) and in herpetological studies (e.g., Gibbs et al. 1994; Prior et al. 1997).

Materials and methods

Tissue sampling and DNA isolation

Larch Mountain salamanders were hand-captured at 15 sites throughout their range (Fig. 1). The distal 1 cm of tail was removed from each individual and stored in liquid nitrogen or on dry ice until it was transferred to an ultra-cold freezer (-80°C). Salamanders from 7 of the sites were returned to their exact collection location after tissue sampling. Individuals captured at the remaining 8 sites were killed with chlorotone solution, fixed in formalin, and stored in buffered ethanol for use as voucher specimens and for studies of salamander morphology and diet. All sampling was performed in accordance with approved Institutional Animal Care and Use Committee protocols under collection permits issued by the US Forest Service. DNA was isolated from each sample using standard phenol-chloroform extraction protocols (Sambrook et al. 1989). The quantity and quality of each DNA sample were determined by fluorimetry and agarose gel electrophoresis, respectively.

Mitochondrial DNA sequencing and analyses

The polymerase chain reaction (PCR) was used to amplify an ~850 bp fragment of the cytochrome *b* gene, using the following primers designed for vertebrates: MVZ-15 (5'-GAACTAATGGCCACACWWTACGNAA-3') and MVZ-16 (5'-AAATAGGAAATATCATTCTGGTTTAAAT-3') (Moritz et al. 1992). Three individuals from each of 14 sampling locations and two individuals from an additional location were randomly selected for sequence analysis (Table 1). Frag-

Table 1. Locations and abbreviations for Larch Mountain salamander (*Plethodon larselli*) populations sampled for genetic analyses.

Site number	Site name	M/R	Haplotype	UTM	
				Northing	Easting
1	Bridal Veil Falls, Oreg.	3/7	A	5044609	564084
2	Multnomah Falls, Oreg.	3/23	B	5047459	569864
3	Herman Creek, Oreg.	3/6	C	5060052	593158
4	Wyeth Campground, Oreg.	3/7	D	5060141	595314
5	Starvation Falls, Oreg.	3/7	C	5060132	602222
6	Cape Horn, Wash.	3/13	E	5047071	563095
7	Dog Creek, Wash.	3/0	F	5062753	603369
8	Lower Copper Creek, Wash.	3/18	G	5070324	560795
9	Zig Zag Creek, Wash.	3/22	H	5077306	569269
10	Ole's Cave, Wash.	3/6	J	5103836	560258
11	Straight Creek, Wash.	3/20	I	5119422	590086
12	Quartz Creek, Wash.	3/19	K	5117283	592217
13	Packwood Palisades, Wash.	3/19	L	5171614	610412
14	Lake Kachess, Wash.	3/0	M	5236995	638500
15	Box Canyon, Wash.	2/0	M	5251500	628100

Note: *M* and *R* are the number of individuals used for mtDNA sequence and RAPD analyses, respectively. Site number corresponds to map location in Fig. 1.

ments were amplified using the following thermalcycling parameters: initial denaturation for 10 min at 93 °C, followed by 40 cycles of 1 min at 93 °C, 1 min at 52 °C, and 2 min at 72 °C. A final extension at 72 °C for 10 min completed the reaction. Each 50- μ L reaction was conducted using 100 ng of template, 0.5 U (1 U \approx 16.67 nkat) of AmpliTaq Gold[®] DNA polymerase (Perkin-Elmer) with the supplied reaction buffer, 2 mmol/L MgCl₂, 1 μ mol/L of each primer, and 100 μ mol/L of each dNTP. PCR amplicons were extracted from a 1% agarose gel using 0.45- μ m Ultrafree-MC centrifugal filters (Millipore); supernatants were transferred to 50- μ m filters (Millipore) to prepare templates for sequencing. A ~400 bp section of the PCR product was sequenced using the primers MVZ-15 and cyt2 (5'-AAACTGCAGCCCCTCAGAATGATATTTGTCCTCA-3'; Kocher et al. 1989) and run on an Applied Biosystems 373A DNA sequencer. Sequences from fragments were aligned by eye using the Genetic Data Environment (Smith et al. 1992) and compared with a GenBank archived cytochrome *b* sequence of *Plethodon elongatus* Van Denburgh, 1916 (accession No. L75821; Moritz et al. 1992).

We used Molecular Evolutionary Genetic Analysis (MEGA) version 2.1 (Kumar et al. 2001) to perform preliminary sequence data explorations and to obtain simple descriptive statistics (average pairwise genetic distances and standard errors) from the haplotypic data. In addition, we used three methods to infer phylogenetic relationships among haplotypes observed in this study. First, we reconstructed a neighbor-joining phylogeny (Saitou and Nei 1987) of haplotypes using PAUP* 4.0b10 (Swofford 1998). Pairwise Jukes–Cantor genetic distances calculated for the haplotypes detected in our analyses were all small (0.002–0.054 substitutions/site) and clearly suggested that relatively short-term evolutionary processes were being observed within *P. larselli*. This indicated that use of more complicated nucleotide substitution models (with their associated greater variances) to account for multiple hits at nucleotide sites was not necessary (Nei and Kumar 2000, p. 112). As a result, we

relied on Jukes–Cantor genetic distances for the neighbor-joining analysis. Second, we used PAUP* to perform maximum parsimony based analyses of our data. Trees were evaluated using a heuristic search and the tree bisection–reconnection algorithm with starting trees obtained from 10 replicates of a random stepwise addition procedure. Bootstrap support for both neighbor-joining and maximum parsimony topologies was obtained from 1000 bootstrap replicates (Felsenstein 1985). A cytochrome *b* sequence from *P. elongatus* (GenBank accession No. L75821; Moritz et al. 1992) was used to root each phylogenetic tree. Finally, we used the computer program TCS (Clement et al. 2000) to produce a haplotype network from the set of sequences observed in *P. larselli*. This procedure, based on the statistical parsimony approach of Templeton et al. (1992), allowed us to infer the genealogy of the set of haplotypes observed among populations.

RAPD procedure and data analyses

RAPD profiles were generated as described in Aagaard et al. (1995). PCR was performed using the following reagent concentrations (25 μ L volume): 1 \times PCR buffer (Promega), 1.8 mmol/L MgCl, 100 μ mol/L of each dNTP, 0.2 μ mol/L primer, 2 ng of template DNA, and 1 U of *Taq* DNA Polymerase (Promega). Thermalcycling was performed using the following parameters: denaturation for 3 min at 93 °C, followed by 45 cycles of denaturation for 1 min at 93 °C, annealing for 1 min at 45 °C, and elongation for 2 min at 72 °C. A final 10-min elongation at 72 °C completed the reaction. Fifteen microlitres of each reaction were loaded in a 2.0% agarose gel and electrophoresed for 4 h (100 V) in TBE (90 mmol/L Tris Base, 90 mmol/L boric acid, 2 mmol/L EDTA, pH 8.0). Amplification products were sized by comparison with a 1-kb DNA ladder (Gibco BRL). The gels were then stained with ethidium bromide (1 μ g/mL) for 30 min and destained for 2 h in deionized water.

Two salamanders from each of four different populations (Straight Creek, Cape Horn, Herman Creek, and Multnomah

Falls; Fig. 1, Table 1) were used to perform preliminary screening of 235 RAPD primers (Oligonucleotide Synthesis Laboratory, University of British Columbia, Vancouver; www.biotech.ubc.ca). This preliminary screening allowed us to identify RAPD primers that yielded distinct, well-separated, and reproducible bands, which were subsequently chosen for final analyses. Reproducibility was assessed in side-by-side RAPD reactions and over multiple RAPD runs. Negative controls were run with all amplifications to check for contamination. Ultimately, 14 RAPD primers that produced a total of 34 distinct, reproducible bands were used for analysis of the complete data set. Information on the specific RAPD primers used for analysis can be obtained from Wagner (2001). Sample sizes from each location for the final data set are provided in Table 1.

We used two congruent approaches for our analyses of the RAPD data. First, owing to the dominant nature of RAPD markers (Lynch and Milligan 1994; Zhivotovsky 1999), we obtained estimates of allele frequencies from the dominant marker data at each locus using the allele frequency estimator of Lynch and Milligan (1994). For purposes of these estimates, each RAPD marker was assumed to be a Mendelian locus whose underlying genotypes corresponded to Hardy–Weinberg expectations. Using these allele frequency estimates, we quantified the extent of genetic differentiation among sampling locations using the F_{ST} analog θ (Weir and Cockerham 1984) with the computer program TFPGA (Miller 1998a). Estimates of θ were obtained for all populations simultaneously (θ_T) as well as separately for groups of populations found north (θ_N) and south (θ_S) of the Columbia River. Ninety-five percent confidence limits for each θ value were obtained through the use of a bootstrap procedure over loci consisting of 1000 resampling replications. In addition, we used TFPGA to calculate pairwise genetic distances between populations using the coancestry distance measure of Reynolds et al. (1983; a logarithmic transformation of pairwise θ estimates), which were subsequently used to perform a UPGMA (unweighted pair group method with arithmetic means) analysis of the interpopulational genetic distance matrix. Node support for the resulting UPGMA dendrogram was obtained through the use of a bootstrap procedure over loci (*sensu* Felsenstein 1985) based on 1000 replicates.

Second, we performed a congruent set of analyses based solely on variation in RAPD marker phenotypes. Rather than relying on potentially biased allele frequencies estimated from dominant marker data (Lynch and Milligan 1994; Zhivotovsky 1999), in these analyses we used only information on the presence or absence of RAPD markers within and among populations. For these analyses, we used the computer program AMOVA-PREP (Miller 1998b) to help prepare input files that were subsequently analyzed using Arlequin 2.001 (Schneider et al. 2000). In Arlequin, we performed an analysis of molecular variance (AMOVA; Excoffier et al. 1992) of marker phenotype variation to obtain an additional measure of overall and pairwise interpopulational genetic differentiation (i.e., F_{ST} values). F_{ST} estimates were calculated based on interindividual genetic distances determined by the number of mismatched RAPD marker phenotypes between individuals. As with the calculation of θ estimates (see above), F_{ST} estimates were obtained for all populations simultaneously (F_{ST}^T) and for sets of popula-

tions located north (F_{ST}^N) and south (F_{ST}^S) of the Columbia River. Furthermore, we obtained F_{ST} estimates for each pairwise combination of populations under analysis. The significance of F_{ST} values (overall and pairwise) was determined via random allocation of individuals and RAPD marker genotypes to populations using 3000 randomization replicates. P values from pairwise contrasts were evaluated at the $\alpha = 0.004$ level to account for nonindependence of genetic distances from the 12 populations examined in the RAPD analyses. As with coancestry distances obtained from TFPGA, pairwise F_{ST} values obtained for populations were subsequently used for UPGMA analyses performed by NTSYS-PC (Rohlf 1994).

Results

mtDNA sequence analyses

Our analyses of cytochrome *b* sequence variation revealed 13 unique haplotypes among the 15 populations examined (Table 2). These sequences were characterized by the presence of 31 variable nucleotide sites (14 parsimony informative) and pairwise percent sequence divergences ranging from 0.2% to 5.2%. In general, haplotypes were location specific with the exception of two sequences (Table 1): haplotype C was found at both Herman Creek and Starvation Falls (Oregon, sites 3 and 5), while Lake Kachess and Box Canyon (Washington, sites 14 and 15) both contained haplotype M. No mtDNA sequence variation was detected within populations. Thus, despite limited sample sizes for sequencing, our analyses suggest that significant differences in haplotype frequencies generally exist among populations. To illustrate, haplotype A (from population 1; Table 1) occurs at a frequency of 0.068 in our data set (three copies of the haplotype from a total of 44 sampled individuals). Under a simple null hypothesis of genetic panmixia, the probability of observing all three copies of this haplotype at the same location is $\sim 0.068^3$, or ~ 0.003 . Clearly, by extension, the probability of observing 11 location-specific haplotypes (Table 1) becomes exceedingly small. Overall, this pattern is consistent with other genetic analyses of salamanders from the northwestern United States (e.g., Jockusch and Wake 2002; Mahoney 2004; Miller et al. 2005) and suggests little contemporary gene flow among populations.


In maximum parsimony analyses, we found three most parsimonious trees (tree length = 103, consistency index = 0.94) that were mainly characterized by rearrangements of haplotypes joined by short branches. Neighbor-joining analysis revealed a single tree with a sum of branch lengths of 0.308. Bootstrap analyses based on both inferential methods produced extremely similar topologies, especially with respect to the relationships of haplotypes found in populations north and south of the Columbia River (Fig. 2). Specifically, southern haplotypes C (from Oregon populations 3 and 5) and D (from Oregon population 4) formed a monophyletic group that was supported by over 98% of bootstrap replicates for both phylogenetic reconstruction techniques. In contrast, all haplotypes detected in populations north of the Columbia River (Washington haplotypes E–M) formed a separate monophyletic group that was supported by over 75% of bootstrap replicates. Interestingly, two southern haplotypes (haplotypes A and B) from populations 1 and 2

Table 3. Pairwise F_{ST} values reflecting the extent of genetic differentiation among 12 Larch Mountain salamander populations used for RAPD analyses in this study.

Population	1	2	3	4	5	6	8	9	10	11	12
1	—										
2	0.18	—									
3	0.51	0.56	—								
4	0.44	0.51	0.21	—							
5	0.47	0.52	0.26	0.08	—						
6	0.42	0.54	0.55	0.51	0.53	—					
8	0.46	0.53	0.54	0.48	0.51	0.15	—				
9	0.41	0.51	0.49	0.44	0.48	0.05	0.08	—			
10	0.38	0.47	0.54	0.48	0.51	0.19	0.16	0.16	—		
11	0.37	0.46	0.52	0.5	0.52	0.14	0.23	0.16	0.14	—	
12	0.39	0.47	0.51	0.46	0.48	0.14	0.08	0.1	0.08	0.08	—
13	0.49	0.57	0.59	0.54	0.56	0.19	0.1	0.15	0.28	0.15	0.12

Note: All values are significantly different from zero at the $\alpha = 0.004$ level, with the exception of those values in bold. These values are nonetheless significant at the $\alpha = 0.05$ level. F_{ST} values are also highly correlated with coancestry distances estimated from the data ($r = 0.95$).

Fig. 3. Haplotype network based on sequence differences among 13 cytochrome *b* haplotypes observed in Larch Mountain salamander. Letters (A–M) indicate one of 13 haplotypes (Table 2) found among 15 populations examined (Fig. 1, Table 1) in Oregon (OR) and Washington (WA). Numbers in parentheses indicate population(s) where haplotypes were detected. Small circles indicate inferred haplotypes that are extinct or were not detected in analyses. Each segment (letter or circle) of the network reflects a single base-pair difference between haplotypes.




smaller than comparable values obtained for the five southern populations analyzed ($\theta_S = 0.41$, UCL = 0.54, LCL = 0.28; $F_{ST}^S = 0.44$, $P < 0.001$).

In addition to high overall statistical indicators of genetic structure, we also observed considerable variation in pairwise genetic distances (coancestry distances and F_{ST} values) between populations. Coancestry distances ranged from 0.05 to 1.1, while pairwise F_{ST} values ranged from 0.05 to 0.59 (Table 3). All pairwise F_{ST} values were significantly different from zero at the $\alpha = 0.004$ level, with the exception of those for Oregon populations 4 and 5 ($F_{ST} = 0.085$, $P = 0.044$), Washington populations 6 and 9 ($F_{ST} = 0.049$, $P = 0.015$), and Washington populations 10 and 12 ($F_{ST} = 0.083$, $P = 0.015$), which nonetheless were significant at the $\alpha = 0.05$ level (Table 3). Pairwise coancestry distances and F_{ST} estimates, however, were highly correlated ($r = 0.95$), indicating that analyses based on estimated allele frequencies and raw marker phenotypes gave qualitatively similar results. Furthermore, UPGMA analyses based on both distance matrices gave comparable results, especially with respect to detection of major population clusters (Fig. 4). UPGMA dendrograms suggested three main clusters of populations with substantial bootstrap support; these clusters largely coincided with well-supported nodes obtained from the phylogenetic relationships of haplotypes detected among populations (Fig. 2). For example, as in our phylogenetic analyses of mtDNA sequence variation, a well-supported cluster (99% of bootstrap replicates) contained all populations north of the Columbia River (Fig. 4). Likewise, Oregon populations 3, 4, and 5, which contained haplotypes C and D (Fig. 2), clustered together in over 80% of bootstrap replicates (Fig. 4). Overall, the main difference between topologies obtained from mtDNA sequence analyses and RAPD markers was the placement of Oregon populations 1 and 2 (containing haplotypes A and B). In UPGMA analyses, these populations clustered together in over 98% of bootstrap replicates (Fig. 4) and, in contrast to the mtDNA sequences, did not show any particularly substantial alliance with northern populations.

Discussion

Our analyses of nuclear and mitochondrial DNA data produced remarkably similar overall patterns (Figs. 2, 4). Specifically, although pairwise analyses of RAPD data

Fig. 4. Midpoint-rooted UPGMA (unweighted pair group method with arithmetic means) dendrograms of (A) pairwise coancestry genetic distances and (B) pairwise F_{ST} values between 12 Larch Mountain salamander populations based on 34 RAPD (random amplified polymorphic DNA) loci. Numbers at branch termini indicate the populations listed in Table 1 and Fig. 1. Letters in parentheses indicate the mitochondrial haplotype (Tables 1, 2) detected at each site and state (OR, Oregon; WA, Washington) for each population. Numbers on branches of the coancestry distance tree reflect the proportions of 1000 bootstrap replicates that supported the topology. Only bootstrap values greater than 0.5 are shown.



suggested that most populations significantly differed from one another, we note that there was substantially less differentiation among populations located north of the Columbia River than among populations to the south. In RAPD analyses, statistical indicators of genetic differentiation for northern populations were low ($\theta_N = 0.13$, UCL = 0.18, LCL = 0.09; $F_{ST}^N = 0.13$, $P < 0.001$) compared with those for southern populations ($\theta_S = 0.41$, UCL = 0.54, LCL = 0.28; $F_{ST}^S = 0.44$, $P < 0.001$). This pattern was also reflected by mtDNA sequence data: average pairwise nucleotide differences among haplotypes were small in the north compared with the south (north, $d = 4.33$, SE = 1.10; south, $d = 8.33$, SE = 2.09). Overall, this pattern may reflect one of two non-exclusive possibilities. First, our data may suggest that the northern region of *P. larselli*'s distribution was more recently established via range expansion, perhaps during the late Pleistocene in concert with glacial retreat. Alternatively, despite detection of significant nuclear differences among populations, our data may also suggest that contemporary gene flow among northern populations is greater and perhaps facilitated through more substantial connectivity among suitable habitats. We propose, however, that the former hypothesis is likely more tenable given the assumed low dispersal ability of many terrestrial salamanders (and *P. larselli* by analogy; see below) and the more patchy distribution of

habitat (and likely salamander populations) in the north compared with the south (Crisafulli 2005). Furthermore, consistent with the range expansion hypothesis, our phylogenetic analyses of mtDNA sequence data show a well-supported northern haplotype clade nested within the southern haplotypes (Fig. 2), with haplotypes from the northern clade forming the distinctive starlike pattern commonly attributed to demographic expansions that occur in concert with range expansions (Avice 2000; Fig. 3). Based on inferences made from the haplotype network (Fig. 3), our data also indicate that individuals from the southwestern populations (or closely related and geographically proximal ones) primarily facilitated the northern range expansion. Overall, our data are extremely consistent with other phylogeographical studies of North American taxa that have detected genetic signatures of northern range expansions following glacial retreat (e.g., Nason et al. 2002; Masta et al. 2003; Starkey et al. 2003; Zamudio and Savage 2003; Ripplinger and Wagner 2005).

Although analyses suggest that individuals from the southwestern region of our study area primarily contributed to the range expansion, we must further provide a mechanism to explain how the Columbia River itself was traversed in this process. The efficacy of rivers as barriers to dispersal of terrestrial plethodontid salamanders has been questioned (Highton 1972). However, based on distributional data, the Columbia River appears to be a barrier for a number of terrestrial salamanders including the Oregon slender salamander, *Batrachoseps wrighti* (Bishop, 1937) (= *B. wrightorum*), and the clouded salamander, *Aneides ferreus* Cope, 1869, for whom the river demarks the northern boundary of their range (Corkran and Thoms 1996). Likewise, the Columbia River defines the southwestern extent of the range of Van Dyke's salamander (*Plethodon vandykei* Van Denburgh, 1906; Jones and Crisafulli 2005). Furthermore, we note that the location of the Columbia River has changed over the long-term course of evolutionary history. The river became repeatedly blocked during the Pliocene and the Pleistocene, a process that significantly deflected the main river channel both north and south of the current main stem on several different occasions (Fecht et al. 1987; Heller et al. 1987). One extreme reroute of the channel altered the course of the river southward towards Mt. Hood (Fig. 1), which may have caused the river to bisect the region between the southwestern (populations 1 and 2) and southeastern populations (populations 3–5) analyzed in this study. Therefore, this process may have released distributional constraints (limits imposed by the river) on southwestern populations, providing an opportunity for a northern range expansion. This hypothesis is also consistent with the high degree of both nuclear and mitochondrial genetic differentiation of southwestern and southeastern populations examined in our study (Figs. 2, 3, 4), as this southern reroute of the river may have acted as a formidable barrier to gene flow that promoted the strong contemporary pattern of divergence between southeastern and southwestern populations. Furthermore, this dramatic hydrogeological event may have subjected populations along the southern bank of the Columbia River to strong bottlenecks that effectively reduced allelic diversity within populations and exacerbated the overall pattern of genetic differentiation in the region. Thus, we suggest that our data

point to the Columbia River as an important overall determinant of patterns of genetic diversity in *P. larselli*. This conclusion is consistent with those reached in a recent study of genetic variation in *B. wrighti* (Miller et al. 2005), which also suggested that changes in the location of the Columbia River may have permitted an expansion of this species' range up to the river's southern bank.

Conservation implications

Larch Mountain salamanders face various threats across their range depending on land ownership and land management objectives. Anthropogenic habitat destruction and fragmentation may have important effects (Herrington and Larsen 1985). In the Columbia River Gorge, road and house construction and development of rock quarries are primary concerns, whereas in the Cascade Mountains, forestry practices are the chief concern. Logging is a particularly acute threat in areas east and northeast of Mount St. Helens, where deep tephra deposits, accumulated over millennia, seal off the salamander's access to cool, moist subterranean microhabitats (Crisafulli et al. 2004). In this tephra fall zone, forest canopy ameliorates forest floor conditions through shading, reduction of desiccating winds, and maintenance of high relative humidity, which provides the microclimatic conditions required by the salamander. In areas outside of deep tephra deposition, unconsolidated surface rock is prevalent and provides the salamanders with access to subterranean retreats during periods of unfavorable surface conditions (i.e., summer drought period). In such rocky areas, forest canopy does not appear to be a critical attribute of Larch Mountain salamander habitat and logging probably poses fewer problems for the salamander, provided that surface rocks and associated microclimates are not altered.

Given their reliance on sites with cool, moist microhabitats (Crisafulli 1999), their narrow physiological tolerances (as with most *Plethodon* species; see Feder 1983; Spotila 1972), and their putative low dispersal capabilities, Larch Mountain salamanders may be constrained in their ability to respond to contemporary disturbances. Indeed, although dispersal and movement of terrestrial salamanders are poorly understood and infrequently quantified, five studies of similar western plethodontid salamanders revealed typical distances moved by individuals, ranging from 1.7 m (*Batrachoseps attenuatus* (Eschscholtz, 1833); Hendrickson 1954) to 23 m (*Ensatina eschscholtzii* Gray, 1850; Stebbins 1954), with a mean distance of 2.5 m (*Plethodon vehiculum* (Cooper, 1860); Hendrickson 1954; Stebbins 1954; Barbour et al. 1969; Barthalmus and Bellis 1972; Ovaska 1988). In addition to the potential for low dispersal, a number of factors related to the reproductive biology of Larch Mountain salamanders could influence species viability when extensive habitat disturbances occur. For example, their reproductive rate may be low because females reach sexual maturity only after 4 years of age, and they appear to have a biennial ovarian cycle with an average clutch size of 7.33 (Herrington and Larsen 1987). Variance in hatching success, juvenile survival, and adult survival are unknown. Furthermore, it has been suggested that the amount of suitable habitat for Larch Mountain salamanders is limited within the Columbia River Gorge itself (Herrington and Larsen 1985) and the central Washington Cascade Range (Crisafulli 1999). Therefore, as-

signment of conservation units to allow for different management options across the range of the Larch Mountain salamander may assist with mitigation of threats to this species.

Previous studies of multiple species have demonstrated strong patterns of genetic differentiation across the Columbia River (Soltis et al. 1997; Arbogast et al. 2001; Monsen and Blouin 2003; Spruell et al. 2003). In contrast, Howard et al. (1983) suggested that populations of Larch Mountain salamanders from each side of the Columbia Gorge were relatively recently diverged (between 4000 and 43 000 years ago) based on low allozyme divergence of two pairs of populations (four populations) located directly across the river from one another. However, their analyses (based on less polymorphic allozyme data) also detected unique alleles for each location examined. Thus, our results may indicate that the Columbia River provides a long-term barrier to dispersal for Larch Mountain salamanders that will ultimately reinforce genetic differentiation of northern and southern populations. Furthermore, we note that morphological differences have been described between northern and southern populations, including variation in number of vomerine teeth and melanophore pigmentation (Brodie 1970). Subsequently, given the concordance of genetic and morphological patterns of divergence, in combination with the river as a long-term barrier to dispersal, we suggest that northern and southern populations are currently on separate evolutionary trajectories, despite the paraphyly of southwestern populations revealed through phylogenetic analyses of mtDNA sequence data (Fig. 2). In addition, our data clearly indicate that northern populations are substantially less differentiated than southern populations. This suggests that extirpation of a southern population would, on average, result in the loss of appreciably more genetic diversity in *P. larselli* relative to extirpation of a northern population. This is especially important considering that although most Columbia River Gorge populations are nominally protected within National Scenic Area reserves, increasing recreational use in this area has the potential to affect already fragmented southern populations (Herrington and Larsen 1985). Furthermore, northern populations are also genetically differentiated (albeit less so than southern populations; Table 3) and face threats from timber harvesting practices. Therefore, designation of separate management strategies for northern versus southern populations may be beneficial to allow for flexibility in mitigating different threats to *P. larselli* across its range.

Acknowledgments

The success of this project was dependent on many individuals to whom we are most grateful. We thank C. Corkran, T. Dove, J. Dwyer, M. Hee, J. Kling, and L. Trippe for help in sample collection. Laboratory assistance was provided by M. Boriss, T. Mullins, M. Rhodes, and S. Warnock. The manuscript benefited from the suggestions of J. Beatty, L. Gorman, M. Hee, D. Kesler, K. Krutovskii, T. Mullins, P. Sanzenbacher, O. Taft, and the Utah State University "Herp Group". This project was funded by the USGS Forest and Rangeland Ecosystem Science Center and the Forest Service, Pacific Northwest Research Station, Olympia Forestry Sciences Laboratory.

References

- Aagaard, J.E., Volmer, S.S., Sorensen, F.C., and Strauss, S.H. 1995. Mitochondrial DNA products among RAPD profiles are frequent and strongly differentiated between races of Douglas-fir. *Mol. Ecol.* **4**: 441–447.
- Arbogast, B.S., Browne, R.A., and Weigl, P.D. 2001. Evolutionary genetics and Pleistocene biogeography of North American tree squirrels (*Tamiasciurus*). *J. Mammal.* **82**: 302–319.
- Aubry, K.B., Senger, C.M., and Crawford, R.I. 1987. Discovery of Larch Mountain salamanders *Plethodon larselli* in the central Cascades Range of Washington. *Biol. Conserv.* **42**: 147–152.
- Avise, J.C. 1994. Molecular markers, natural history and evolution. Chapman and Hall, New York.
- Avise, J.C. 2000. Phylogeography. The history and formation of species. Harvard University Press, Cambridge, Mass.
- Baker, C.S., Perry, A., Chambers, G.K., and Smith, P.J. 1995. Population variation in the mitochondrial cytochrome *b* gene of the orange roughy *Hoplostethus atlanticus* and the Hoki *Macruronus novaezelandiae*. *Mar. Biol. (Berl.)*, **122**: 503–509.
- Barbour, J.W., Hardin, R.W., Schafer, J.P., and Harvey, M.J. 1969. Home range, movements and activity of the dusky salamander, *Desmognathus fuscus*. *Copeia*, 1969: 293–297.
- Bartholomus, G.T., and Bellis, E.D. 1972. Home range, homing and the homing mechanism of the salamander, *Desmognathus fuscus*. *Copeia*, 1972: 632–642.
- Blaustein, A.R., Wake, D.B., and Sousa, W.P. 1994. Amphibian declines: judging stability, persistence and susceptibility of populations to local and global extinctions. *Conserv. Biol.* **8**: 60–71.
- Brodie, E.D., Jr. 1970. Western salamanders of the genus *Plethodon*: systematics and geographic variation. *Herpetologica*, **26**: 468–516.
- Burns, D.M. 1954. A new subspecies of the salamander *Plethodon vandykei*. *Herpetologica*, **10**: 83–87.
- Burns, D.M. 1962. The taxonomic status of the salamander *Plethodon vandykei larselli*. *Copeia*, 1962: 177–181.
- Burns, D.M. 1964. *Plethodon larselli*. Catalogue of American amphibians and reptiles, **13**: 1.
- Castilla, A.M., Fernandez-Pedrosa, V., Backeljau, T., Gonzalez, A., Latorre, A., and Moya, A. 1998. Conservation genetics of insular *Podarcis* lizards using partial cytochrome *b* sequences. *Mol. Ecol.* **7**: 1407–1411.
- Clement, M., Posada, D., and Crandall, K.A. 2000. TCS: a program to estimate gene genealogies. *Mol. Ecol.* **9**: 1657–1659.
- Corkran, C.C., and Thoms, C. 1996. Amphibians of Oregon, Washington and British Columbia. Lone Pine, Renton, Wash.
- Crisafulli, C.M. 1999. Survey protocol for the Larch Mountain salamander (*Plethodon larselli*). In Survey protocols for amphibians under the Survey and Manage provision of the Northwest Forest Plan. Version 3.0. Edited by D.H. Olson. Interagency Publication of the Regional Ecosystem Office, Portland, Oreg. BLM Publication BLM/OR/WA/PT-00/033+1792, US GPO 2000-589-124/04022 Region No. 10. Available from <http://www.or.blm.gov/surveyandmanage/SP/Amphibians99/protoch.pdf> [cited 16 February 2005].
- Crisafulli, C.M. 2005. Larch Mountain salamander, *Plethodon larselli* Burns. In Amphibians of the Pacific Northwest. Edited by L.L.C. Jones, W.P. Leonard, and D.H. Olson. Seattle Audubon Society, Seattle, Wash. In Press.
- Crisafulli, C.M., Clayton, D.R., and Olson, D.H. 2004. Conservation assessment for the Larch mountain salamander (*Plethodon larselli*). Interagency Publication of the Regional Ecosystem Office, Portland, Oreg. BLM Publications, Portland, Oreg.
- Darda, D.M., and Garvey-Darda, P.A. 1995. Geographic distribution: *Plethodon larselli*. *Herpetol. Rev.* **26**: 150.
- Excoffier, L., Smouse, P.E., and Quattro, J.M. 1992. Analysis of molecular variance inferred from metric distances among DNA haplotypes: application to mitochondrial DNA restriction data. *Genetics*, **131**: 479–491.
- Fecht, K.R., Reidel, S.P., and Tallman, A.M. 1987. Paleodrainage of the Columbia River system on the Columbia Plateau of Washington State — a summary. In Selected papers on the geology of Washington. Edited by J.E. Shuster. Washington Division of Geology and Earth Resources Bulletin 77. pp. 219–248.
- Feder, M.E. 1983. Integrating the ecology and physiology of plethodontid salamanders. *Herpetologica*, **39**: 291–310.
- Felsenstein, J. 1985. Confidence limits on phylogenies: an approach using the bootstrap. *Evolution*, **39**: 783–791.
- Fleischer, R.C., Fuller, G., and Ledig, D.B. 1995. Genetic structure of endangered clapper rail (*Rallus longirostris*) populations in southern California. *Conserv. Biol.* **9**: 1234–1243.
- Gibbs, H.L., Prior, K.A., and Weatherhead, P.J. 1994. Genetic analysis of populations of a threatened snake species using RAPD markers. *Mol. Ecol.* **3**: 329–337.
- Good, D.A., and Wake, D.B. 1992. Geographic variation and speciation in the torrent salamanders of the genus *Rhyacotriton* (Caudata: Rhyacotritonidae). *Univ. Calif. Publ. Zool.* **126**: 1–91.
- Haig, S.M. 1998. Molecular contributions to conservation genetics. *Ecology*, **79**: 413–425.
- Haig, S.M., Rhymer, J.M., and Heckel, D.G. 1994. Population differentiation in randomly amplified polymorphic DNA of red-cockaded woodpeckers. *Mol. Ecol.* **3**: 581–595.
- Haig, S.M., Bowman, R., and Mullins, T.D. 1996. Population structure of red-cockaded woodpeckers in south Florida: RAPDs revisited. *Mol. Ecol.* **5**: 725–734.
- Haig, S.M., Gratto-Trevor, C.L., Mullins, T.D., and Colwell, M.A. 1997. Population identification of western hemisphere shorebirds throughout the annual cycle. *Mol. Ecol.* **6**: 413–427.
- Harris, S.L. 1988. Fire mountains of the west: the Cascade and Mono Lake volcanoes. Mountain Press, Missoula, Mont.
- Hedges, S.B., Bogart, J.P., and Maxson, L.R. 1992. Ancestry of unisexual salamanders. *Nature (Lond.)*, **356**: 708–710.
- Heller, P.L., Tabor, R.W., and Suczek, C.A. 1987. Paleographic evolution of the United States Pacific Northwest during Paleogene time. *Can. J. Earth Sci.* **24**: 1652–1667.
- Hendrickson, J.R. 1954. Ecology and systematics of salamanders of the genus *Batrachoseps*. *Univ. Calif. Publ. Zool.* **54**: 1–46.
- Herrington, R.E., and Larsen, J.H., Jr. 1985. Current status, habitat requirements and management of the Larch Mountain salamander *Plethodon larselli* Burns. *Biol. Conserv.* **34**: 169–179.
- Herrington, R.E., and Larsen, J.H., Jr. 1987. Reproductive biology of the larch mountain salamander (*Plethodon larselli*). *J. Herpetol.* **21**: 48–56.
- Highton, R. 1972. Distributional interactions among eastern North American salamanders of the genus *Plethodon*. In The distributional history of the biota of the southern Appalachians. III. Vertebrates. Research division monograph 4, Virginia Polytechnic Institute, Blacksburg, Va. pp. 139–188.
- Highton, R. 1995. Speciation in eastern North American salamanders of the genus *Plethodon*. *Annu. Rev. Ecol. Syst.* **26**: 579–600.
- Howard, J.H., Wallace, R.L., and Larsen, J.H., Jr. 1983. Genetic variation and population divergence in the Larch Mountain salamander (*Plethodon larselli*). *Herpetologica*, **39**: 41–47.
- International Union for the Conservation of Nature. 2004. IUCN red list of threatened species [online]. Available from <http://www.redlist.org> [cited 16 February 2005].

- Jackman, T.R., Applebaum, G., and Wake, D.B. 1997. Phylogenetic relationships of Bolitoglossine salamanders: a demonstration of the effects of combining morphological and molecular data sets. *Mol. Biol. Evol.* **14**: 883–891.
- Jockusch, E.L. 1996. Evolutionary studies in *Batrachoseps* and the other plethodontid salamanders: correlated character evolution, molecular phylogenetics, and reaction norm evolution. Ph.D. thesis, Department of Integrative Biology, University of California, Berkeley, Calif.
- Jockusch, E.L., and Wake, D.B. 2002. Falling apart and merging: diversification of slender salamanders (Plethodontidae: *Batrachoseps*) in the American West. *Biol. J. Linn. Soc.* **76**: 361–391.
- Jones, L.C., and Crisafulli, C.M. 2005. Van Dyke's salamander, *Plethodon vandykei* Van Denburgh. In *Amphibians of the Pacific Northwest*. Edited by L.L.C. Jones, W.P. Leonard, and D.H. Olson. Seattle Audubon Society, Seattle, Wash. In Press.
- Kimberling, D.N., Ferrarira, A.R., Shuster, S.M., and Keim, P. 1996. RAPD marker estimation of genetic structure among isolated northern leopard frog populations in the south-western USA. *Mol. Ecol.* **5**: 521–529.
- Kirk, J.J. 1983. Distribution of *Plethodon larselli* in Oregon with notes on plethodontids. Report to the Oregon Department of Fish and Wildlife, Portland, Oreg.
- Kocher, T.D., Thomas, W.K., Edwards, S.V., Paabo, S., Villablanca, F.X., and Wilson, A.C. 1989. Dynamics of mitochondrial DNA evolution in animals: amplifications and sequencing with conserved primers. *Proc. Natl. Acad. Sci. U.S.A.* **86**: 6196–6200.
- Kumar, S., Tamura, K., Jakobsen, I.B., and Nei, M. 2001. MEGA2: Molecular Evolutionary Genetics Analysis software. *Bioinformatics*, **17**: 1244–1245.
- Lento, G.M., Haddon, M., Chambers, G.K., and Baker, C.S. 1997. Genetic variation of southern hemisphere fur seals (*Arctocephalus* spp.). *J. Hered.* **88**: 202–208.
- Lynch, M., and Milligan, B.G. 1994. Analysis of population genetic structure with RAPD markers. *Mol. Ecol.* **3**: 91–99.
- Mahoney, M. 2004. Molecular systematics and phylogeography of the *Plethodon elongates* species group: combining phylogenetic and population genetic methods to investigate species history. *Mol. Ecol.* **13**: 149–166.
- Masta, S.E., Laurent, N.M., and Routman, E.J. 2003. Population genetic structure of the toad *Bufo woodhousii*: an empirical assessment of the effects of haplotype extinction on nested clade analysis. *Mol. Ecol.* **12**: 1541–1554.
- McGroder, M.F. 1991. Reconciliation of two-sided thrusting, burial metamorphism, and diachronous uplift in the Cascades of Washington and British Columbia. *Geol. Soc. Am. Bull.* **103**: 189–209.
- Miller, M.P. 1998a. Tools for population genetic analysis (TFPGA). Version 1.3 [computer program]. Available from <http://www.marksgeneticsoftware.net/tfpga.htm> [cited 18 February 2005].
- Miller, M.P. 1998b. AMOVA-PREP [computer program]. Available from <http://www.marksgeneticsoftware.net/amovaprep.htm> [cited 18 February 2005].
- Miller, M.P., Haig, S.M., and Wagner, R.S. 2005. Conflicting patterns of genetic structure produced by nuclear and mitochondrial markers in the Oregon slender salamander (*Batrachoseps wrighti*): implications for conservation efforts and species management. *Conserv. Genet.* In Press.
- Monsen, K.J., and Blouin, M.S. 2003. Genetic structure in a montane ranid frog: restricted gene flow and nuclear–mitochondrial discordance. *Mol. Ecol.* **12**: 3275–3286.
- Moritz, C., Schneider, C.J., and Wake, D.B. 1992. Evolutionary relationships within the *Ensatina eschscholtzii* complex confirm the ring species interpretation. *Syst. Biol.* **41**: 273–291.
- Mundy, N.I., Winchell, C.S., and Woodruff, D.S. 1997. Genetic differences between the endangered San Clemente Island loggerhead shrike *Lanius ludovicianus mearnsi* and two neighbouring subspecies demonstrated by mtDNA control region and cytochrome *b* sequence variation. *Mol. Ecol.* **6**: 29–37.
- Nason, J.D., Hamrick, J.L., and Fleming, T.H. 2002. Historical vicariance and postglacial colonization effects on the evolution of genetic structure in *Lophocereus*, a Sonoran Desert columnar cactus. *Evolution*, **56**: 2214–2226.
- Nei, M., and Kumar, S. 2000. Molecular evolution and phylogenetics. Oxford University Press, New York.
- Nusser, J.A., Goto, R.M., Ledig, D.B., Fleischer, R.C., and Miller, M.M. 1996. RAPD analysis reveals low genetic variability in the endangered light-footed clapper rail. *Mol. Ecol.* **5**: 463–472.
- Ovaska, K. 1988. Spacing and movements of the salamander *Plethodon vehiculum*. *Herpetologica*, **44**: 377–386.
- Prior, K.A., Gibbs, H.L., and Weatherhead, P.J. 1997. Population genetic structure in the black rat snake: implications for management. *Conserv. Biol.* **11**: 1147–1158.
- Ripplinger, J., and Wagner, R.S. 2005. Phylogeography of northern populations of the Pacific Chorus Frog, *Pseudacris regilla*. *Northwest. Nat.* **85**: 118–125.
- Reynolds, J., Weir, B.S., and Cockerham, C.C. 1983. Estimation of the coancestry coefficient: basis for a short-term genetic distance. *Genetics*, **105**: 767–779.
- Rohlf, F.J. 1994. NTSYS-pc: Numerical Taxonomy and Multivariate Analysis System. Version 1.8 [computer program]. Exeter Software, Setauket, N.Y.
- Saitou, N., and Nei, M. 1987. The neighbour-joining method: a new method for reconstructing phylogenetic trees. *Mol. Biol. Evol.* **4**: 406–425.
- Sambrook, J., Fritsch, E.F., and Maniatis, T. 1989. Molecular cloning: a laboratory manual. 2nd ed. Cold Spring Harbor Laboratory Press, Plainview, N.Y.
- Schneider, S., Roessli, D., and Excoffier, L. 2000. Arlequin: a software for population genetics data analysis [computer program]. Available from the authors, Genetics and Biometry Laboratory, Department of Anthropology and Ecology, University of Geneva, Geneva, Switzerland.
- Smith, S.W., Wang, C., Gillevet, P.M., and Gilbert, W. 1992. Genetic Data Environment and the Harvard Genome Database: genome mapping and sequencing. Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y. Computer program available from <ftp://ftp.bio.indiana.edu/molbio/unix/GDE/>.
- Soltis, D., Gitzendanner, M., Strenge, D., and Soltis, P. 1997. Chloroplast DNA intraspecific phylogeny of plants from the Pacific Northwest of North America. *Plant Syst. Evol.* **206**: 353–373.
- Spotila, J.R. 1972. Role of temperature and water in the ecology of lungless salamanders. *Ecol. Monogr.* **42**: 95–125.
- Spruell, P., Hemmingsen, A.R., Howell, P.J., Kanda, N., and Allendorf, F.W. 2003. Conservation genetics of bull trout: geographical distribution of variation at microsatellite loci. *Conserv. Genet.* **4**: 17–29.
- Starkey, D.E., Shaffer, H.B., Burke, R.L., Forstner, M.J.R., Iverson, J.B., Janzen, F.J. et al. 2003. Molecular systematics, phylogeography, and the effects of Pleistocene glaciation in the painted turtle (*Chrysemys picta*) complex. *Evolution*, **57**: 119–128.
- Stebbins, R.C. 1954. Natural history of the salamanders of the plethodontid genus *Ensatina*. *Univ. Calif. Publ. Zool.* **54**: 47–124.

- Swofford, D.L. 1998. Phylogenetic Analysis Using Parsimony (PAUP*). Version 4.0b10 [computer program]. Sinauer Associates, Sunderland, Mass.
- Tan, A., and Wake, D. 1995. MtDNA phylogeography of the California newt, *Taricha torosa* (Caudata, salamandridae). *Mol. Phylogenet. Evol.* **4**: 383–394.
- Templeton, A.R., Crandall, K.A., and Sing, C.F. 1992. A cladistic analysis of phenotypic associations with haplotypes inferred from restriction endonuclease mapping and DNA sequence data. III. Cladogram estimation. *Genetics*, **132**: 619–633.
- Tilley, S.G., and Mahoney, M.J. 1996. Patterns of genetic differentiation of *Desmognathus ochrophaeus* complex (Amphibia: Plethodontidae). *Herpetol. Monogr.* **10**: 1–42.
- US Forest Service and US Bureau of Land Management. 1994. Final supplemental environmental impact statement on management of habitat for inter-successional and old growth forest related species within the range of the Northern Spotted Owl. US Forest Service and US Bureau of Land Management, Portland, Ore.
- Wagner, R.S. 2001. Phylogeography, evolution and conservation in forest-associated Pacific Northwest salamanders. Ph.D. thesis, Oregon State University, Corvallis, Ore.
- Walker, D., Moler, P.E., Buhlmann, K.A., and Avise, J.C. 1998. Phylogeographic uniformity in mitochondrial DNA of the snapping turtle (*Chelydra serpentina*). *Anim. Conserv.* **1**: 55–60.
- Weir, B.S., and Cockerham, C.C. 1984. Estimating *F*-statistics for the analysis of population structure. *Evolution*, **38**: 1358–1370.
- Zamudio, K.R., and Savage, W.K. 2003. Historical isolation, range expansion, and secondary contact of two highly divergent mitochondrial lineages in spotted salamanders (*Ambystoma maculatum*). *Evolution*, **57**: 1631–1652.
- Zhivotovsky, L.A. 1999. Estimating population structure in diploids with multilocus dominant DNA markers. *Mol. Ecol.* **8**: 507–514.

This article has been cited by:

1. J. A. Lee-Yaw, D. E. Irwin. 2012. Large geographic range size reflects a patchwork of divergent lineages in the long-toed salamander (*Ambystoma macrodactylum*). *Journal of Evolutionary Biology* n/a. [[CrossRef](#)]
2. J. A. Lee-Yaw, D. E. Irwin. 2012. Large geographic range size reflects a patchwork of divergent lineages in the long-toed salamander (*Ambystoma macrodactylum*). *Journal of Evolutionary Biology* **25**:11, 2276. [[CrossRef](#)]
3. J. A. Lee-Yaw, D. E. Irwin. 2012. Large geographic range size reflects a patchwork of divergent lineages in the long-toed salamander (*Ambystoma macrodactylum*). *Journal of Evolutionary Biology* **25**:11, 2276. [[CrossRef](#)]
4. Tara A. Pelletier, Deborah A. Duffield, Edward A. DeGrauw. 2011. Rangewide Phylogeography of the Western Red-Backed Salamander (*Plethodon vehiculum*). *Northwestern Naturalist* **92**:3, 200-210. [[CrossRef](#)]
5. Sarah E. May, Kim A. Medley, Steve A. Johnson, Eric A. Hoffman. 2011. Combining genetic structure and ecological niche modeling to establish units of conservation: A case study of an imperiled salamander. *Biological Conservation* **144**:5, 1441-1450. [[CrossRef](#)]
6. AARON B. A. SHAFER, CATHERINE I. CULLINGHAM, STEEVE D. CÔTÉ, DAVID W. COLTMAN. 2010. Of glaciers and refugia: a decade of study sheds new light on the phylogeography of northwestern North America. *Molecular Ecology* **19**:21, 4589-4621. [[CrossRef](#)]
7. Anna M. Goebel, Tom A. Ranker, Paul Stephen Corn, Richard G. Olmstead. 2009. Mitochondrial DNA evolution in the *Anaxyrus boreas* species group. *Molecular Phylogenetics and Evolution* **50**:2, 209-225. [[CrossRef](#)]
8. JULIE A. LEE-YAW, JASON T. IRWIN, DAVID M. GREEN. 2008. Postglacial range expansion from northern refugia by the wood frog, *Rana sylvatica*. *Molecular Ecology* **17**:3, 867-884. [[CrossRef](#)]
9. Jennifer A. Dever. 2007. Fine-scale Genetic Structure in the Threatened Foothill Yellow-legged Frog (*Rana Boylii*). *Journal of Herpetology* **41**:1, 168-173. [[CrossRef](#)]
10. D.M.MarshD.M. Marsha(540) 458-8176fax: (540) 458-8012(email: marshd@wlu.edu)R.B.PageR.B. Pagea‡(email: Robert.Page@uky.edu)*T.J.HanlonT.J. HanlonaH.BarekeH. BarekeR.CorritoneR. CorritoneaN.JetterN. JetteraN.G.BeckmanN.G. Beckmana§(email: beckm089@umn.edu)†K.GardnerK. GardneraD.E.SeifertD.E. SeifertaP.R.CabeP.R. Cabea. 2007. Ecological and genetic evidence that low-order streams inhibit dispersal by red-backed salamanders (*Plethodon cinereus*). *Canadian Journal of Zoology* **85**:3, 319-327. [[Abstract](#)] [[Full Text](#)] [[PDF](#)] [[PDF Plus](#)]
11. CRAIG A. STEELE, ANDREW STORFER. 2006. Coalescent-based hypothesis testing supports multiple Pleistocene refugia in the Pacific Northwest for the Pacific giant salamander (*Dicamptodon tenebrosus*). *Molecular Ecology* **15**:9, 2477-2487. [[CrossRef](#)]