

8-24-2012

A new species of *Culcua* Walker (Diptera: Stratiomyidae) from Vietnam

Norman E. Woodley

Smithsonian Entomology Lab, USDA-ARS, norman.woodley@ars.usda.gov

Follow this and additional works at: <http://digitalcommons.unl.edu/insectamundi>



Part of the [Entomology Commons](#)

Woodley, Norman E., "A new species of *Culcua* Walker (Diptera: Stratiomyidae) from Vietnam" (2012). *Insecta Mundi*. 759.
<http://digitalcommons.unl.edu/insectamundi/759>

This Article is brought to you for free and open access by the Center for Systematic Entomology, Gainesville, Florida at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Insecta Mundi by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

INSECTA MUNDI

A Journal of World Insect Systematics

0243

A new species of *Culcua* Walker (Diptera: Stratiomyidae) from Vietnam

Norman E. Woodley
Systematic Entomology Laboratory–PSI–ARS–USDA
Smithsonian Institution NHB–168
P O Box 37012
Washington, DC 20013–7012, USA
norman.woodley@ars.usda.gov

Date of Issue: August 24, 2012

Norman E. Woodley
A new species of *Culcua* Walker (Diptera: Stratiomyidae) from Vietnam
Insecta Mundi 0243: 1–4

Published in 2012 by

Center for Systematic Entomology, Inc.
P. O. Box 141874
Gainesville, FL 32614-1874 USA
<http://www.centerforsystematicentomology.org/>

Insecta Mundi is a journal primarily devoted to insect systematics, but articles can be published on any non-marine arthropod. Topics considered for publication include systematics, taxonomy, nomenclature, checklists, faunal works, and natural history. **Insecta Mundi** will not consider works in the applied sciences (i.e. medical entomology, pest control research, etc.), and no longer publishes book reviews or editorials. **Insecta Mundi** publishes original research or discoveries in an inexpensive and timely manner, distributing them free via open access on the internet on the date of publication.

Insecta Mundi is referenced or abstracted by several sources including the Zoological Record, CAB Abstracts, etc. **Insecta Mundi** is published irregularly throughout the year, with completed manuscripts assigned an individual number. Manuscripts must be peer reviewed prior to submission, after which they are reviewed by the editorial board to ensure quality. One author of each submitted manuscript must be a current member of the Center for Systematic Entomology.

Managing editor: Paul E. Skelley, e-mail: insectamundi@gmail.com

Production editors: Michael C. Thomas, Brian Armitage, Ian Stocks

Editorial board: J. H. Frank, M. J. Paulsen

Subject editors: G.B. Edwards, J. Eger, A. Rasmussen, F. Shockley, G. Steck, Ian Stocks, A. Van Pelt, J. Zaspel

Spanish editors: Julieta Brambila, Angélico Asenjo

Printed copies (ISSN 0749-6737) deposited in libraries of:

CSIRO, Canberra, ACT, Australia
Museu de Zoologia, São Paulo, Brazil
Agriculture and Agrifood Canada, Ottawa, ON, Canada
The Natural History Museum, London, Great Britain
Muzeum i Instytut Zoologiczny PAN, Warsaw, Poland
National Taiwan University, Taipei, Taiwan
California Academy of Sciences, San Francisco, CA, USA
Florida Department of Agriculture and Consumer Services, Gainesville, FL, USA
Field Museum of Natural History, Chicago, IL, USA
National Museum of Natural History, Smithsonian Institution, Washington, DC, USA
Zoological Institute of Russian Academy of Sciences, Saint-Petersburg, Russia

Electronic copies (On-Line ISSN 1942-1354, CDROM ISSN 1942-1362) in PDF format:

Printed CD mailed to all members at end of year.
Florida Virtual Campus: <http://purl.fcla.edu/fcla/insectamundi>
University of Nebraska-Lincoln, Digital Commons: <http://digitalcommons.unl.edu/insectamundi/>
Goethe-Universität, Frankfurt am Main: <http://edocs.ub.uni-frankfurt.de/volltexte/2010/14363/>

Author instructions available on the *Insecta Mundi* page at:
<http://www.centerforsystematicentomology.org/insectamundi/>

Copyright held by the author(s). This is an open access article distributed under the terms of the Creative Commons, Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited. <http://creativecommons.org/licenses/by-nc/3.0/>

A new species of *Culcua* Walker (Diptera: Stratiomyidae) from Vietnam

Norman E. Woodley
Systematic Entomology Laboratory–PSI–ARS–USDA
Smithsonian Institution NHB–168
P O Box 37012
Washington, DC 20013–7012, USA
norman.woodley@ars.usda.gov

Abstract. A new species of *Culcua* Walker (Diptera: Stratiomyidae), *C. lingafelteri* Woodley, **new species**, is described from northern Vietnam. It is diagnosed relative to other species using the recent revision of the genus by Rozkošný and Kozánek (2007). This is the first species of *Culcua* reported from Vietnam.

Introduction

The genus *Culcua* Walker, in the subfamily Pachygastrinae of the family Stratiomyidae (Diptera), is known only from the Oriental Region, where it ranges from eastern India east to Taiwan, the Philippines, and the island of Borneo. The genus currently contains ten species, seven of which were described in a recent revision of *Culcua* (Rozkošný and Kozánek 2007). Most of the species are known from very few specimens and virtually nothing is known about their life histories. A paratype specimen of *C. kovaci* Rozkošný and Kozánek from Thailand has label data stating “on leaves nr. fallen tree”. As members of numerous other taxa of Pachygastrinae are attracted to fallen trees and the majority of species have larvae that live under bark (Kraft and Cook 1961), it seems likely that *Culcua* larvae live under bark as well.


A specimen recently collected in Vietnam, housed in the National Museum of Natural History at the Smithsonian Institution (USNM), was found to be a new species. It is described here and compared to the most similar known species.

Culcua lingafelteri Woodley, new species

Figures 1, 3, 4

Diagnosis. The combination of scutum with appressed setulae silvery white interspersed with long, erect black setulae; all tibiae black; first tarsomere of middle leg yellow; first tarsomere of hind leg black; wing with a dark and hyaline pattern; and abdomen about as broad as long will separate *Culcua lingafelteri* from other species in the genus. In the key to species of Rozkošný and Kozánek (2007: 38) this species will key to *C. argentea* Rozkošný and Kozánek but differs from it by having darker antennae (Figs. 1, 2); longer, more conspicuous pilosity on the eye; the postocular orbit narrower; conspicuous, erect black setulae present on the scutum and scutellum (Figs. 4, 5); and wing cell cua_1 mostly hyaline (Fig. 3).

Description. Female. **Head** (Fig. 1) black. Lower frons and face with grayish tomentum covering surface, face with scattered erect pale setulae; upper frons with margins from ocellar triangle to about half distance to antennae yellowish-white tomentose interspersed with longer erect pale setulae; surface shiny, virtually impunctate, slightly concave. Ocellar tubercle moderately prominent, with erect pale setulae. Postocular orbit visible in profile, narrowest at about upper third, posterodorsally with sharply rounded margin, this and postoccipt with dense erect pale setulae, longest ventrally. Eyes with moderately dense brownish-black setulae. Antenna with scape and pedicel black with blackish setulae that are longest on scape; flagellar complex brownish yellow, short penultimate flagellomere and base of last, arista-like flagellomere dark yellowish, the latter becoming dark apically. Palpus with first segment about half the length of the second, black, with erect dark setulae; second segment nearly cylindrical with rounded apex, brownish, velvety in appearance with two or three short dark setulae at apex.



Figures 1–3. Morphological features of *Culcua* species. **1)** *Culcua lingafelteri*, lateral view of head of female holotype. **2)** *Culcua argentea*, lateral view of head of female paratype. **3)** Wing of *C. lingafelteri* (anal area of wing is folded underneath).

Thorax black, only spines of scutellum yellowish white. Most of thoracic dorsum and pleura with vestiture of semi-appressed silvery white setulae except for median vitta on scutum that widens slightly on posterior half, lateral areas of scutum posterior to the transverse suture, entire scutellum with short, semi-appressed black setulae; anterior half of anepisternum, and posterior half of anepimeron bare and shiny; entire scutum and scutellum with erect dark setulae with a few pale setulae at extreme lateral areas of scutum. Scutellar spines elongate conical, subequal in length and each about half as long as scutellum. Legs black, middle and hind tarsi slightly more brownish; middle tarsus with basal segment dark yellowish, second segment brownish yellow; hind tarsus with first segment indistinctly paler at base. Wing hyaline with dark pattern as in Fig. 3, evenly set with microtrichia except basal part of cell c and basal and narrow anterior part of cell cup bare. Alula ovoid, completely set with microtrichia. Halter whitish yellow. Post-tegula with a few short dark setulae.

Abdomen black, sublateral callosities on tergite 3 strongly developed. Vestiture of tergites 1–3 with short semi-appressed black setulae medially; lateral areas of tergites 1–4 with longer, semi-erect



4



5

Figures 4–5. Lateral habitus view of *Culcua* species. 4) *Culcua lingafelteri*, female holotype. 5) *Culcua argentea*, female paratype.

silvery white setulae on approximately apical halves, forming poorly defined pilose spots; tergite 5 evenly clothed with moderately sparse erect pale setulae. Sternite 1 with moderate, fine dark brownish tomentum and short pale setulae medially and laterally; sternites 2–4 evenly set with semi-appressed silvery setulae; sternite 5 with erect dark setulae. Cerci two-segmented, segment 1 cylindrical, brownish; segment 2 darker, elongate-ovoid, about as long as segment 1; both segments with erect pale setulae with a few dark setulae intermixed.

Dimensions. Body length (excluding antennae) 8.9 mm; wing length 8.2 mm.

Distribution. This species is known only from Vietnam. It is the first species of *Culcua* reported for that country.

Type material. Holotype female (USNM) is labeled (slashes indicate each label): “VIETNAM: Vinh Phuc Prov. Tam Dao National Park, Thong Tin TV tower 21°27'49.01”N, 105°38'47.42”E ~1200m 17–23 June 2011/Steven Lingafelter Eduard Jendek Eduard Vives Pham Hong Thai/HOLOTYPE/*Culcua lingafelteri* Woodley 2012”.

Discussion. As noted in the diagnosis, *Culcua lingafelteri* is most similar to *C. argentea*. In addition to the characters mentioned in the diagnosis, *C. lingafelteri* differs from *C. argentea* by being larger (body length range for *C. argentea* given by Rozkošný and Kozánek (2007: 40) is 6.5–7.9 mm), by having a uniformly black median vitta on the scutum (it is black with a very narrow silvery line medially in *C. argentea*), and having the sublateral callosities on abdominal tergite 3 more strongly developed.

Etymology. The species name is a patronym for Steven W. Lingafelter, my friend and life partner, who collected the specimen.

Acknowledgments

I thank Steve Lingafelter for collecting this and other interesting stratiomyids on his recent trip to Vietnam. I am grateful to Keith Arakaki (Bishop Museum, Honolulu) for arranging the loan of the type series of *Culcua argentea*. Gary Ouellette (Systematic Entomology Laboratory) took the images and prepared the plates for publication. I am grateful to Wayne Mathis (Smithsonian Institution) as well as Michael Pogue and Allen Norrbom (Systematic Entomology Lab) for reviewing the manuscript. USDA is an equal opportunity employer and provider.

Literature Cited

- Kraft, K. J. and E. F. Cook. 1961.** A revision of the Pachygasterinae [sic] (Diptera, Stratiomyidae) of America north of Mexico. Miscellaneous Publications of the Entomological Society of America 3(1): 1–24.
- Rozkošný, R. and M. Kozánek. 2007.** A review of the Oriental *Culcua* with descriptions of seven new species (Diptera: Stratiomyidae). Insect Systematics & Evolution 38(1): 35–50.

Received June 2, 2012; Accepted July 29, 2012.

Subject edited by Gary Steck.