

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Publications from USDA-ARS / UNL Faculty

U.S. Department of Agriculture: Agricultural
Research Service, Lincoln, Nebraska

2009

Response Of Adult Mosquitoes To Light-Emitting Diodes Placed In Resting Boxes And In The Field

Michael T. Bentley
University of Florida

Phillip E. Kaufman
University of Florida

Daniel L. Kline
United States Department of Agriculture

Jerome A. Hogsette
United States Department of Agriculture

Follow this and additional works at: <https://digitalcommons.unl.edu/usdaarsfacpub>



Part of the [Agricultural Science Commons](#)

Bentley, Michael T.; Kaufman, Phillip E.; Kline, Daniel L.; and Hogsette, Jerome A., "Response Of Adult Mosquitoes To Light-Emitting Diodes Placed In Resting Boxes And In The Field" (2009). *Publications from USDA-ARS / UNL Faculty*. 997.
<https://digitalcommons.unl.edu/usdaarsfacpub/997>

This Article is brought to you for free and open access by the U.S. Department of Agriculture: Agricultural Research Service, Lincoln, Nebraska at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Publications from USDA-ARS / UNL Faculty by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

RESPONSE OF ADULT MOSQUITOES TO LIGHT-EMITTING DIODES PLACED IN RESTING BOXES AND IN THE FIELD

MICHAEL T. BENTLEY,¹ PHILLIP E. KAUFMAN,^{1,3} DANIEL L. KLINE,² AND JEROME A. HOGSETTE²

ABSTRACT. The response of adult mosquitoes to 4 light-emitting diode (LED) wavelengths was evaluated using diode-equipped sticky cards (DESCs) and diode-equipped resting boxes at 2 sites in north central Florida. Wavelengths evaluated were blue (470 nm), green (502 nm), red (660 nm), and infrared (IR) (860 nm). When trapping with DESCs, 15 mosquito species from 7 genera (*Aedes*, *Anopheles*, *Coquillettia*, *Culex*, *Mansonia*, *Psorophora*, and *Uranotaenia*) were captured. Overall, approximately 43.8% of all mosquitoes were trapped on DESCs fitted with green LEDs. Significantly more females of *Aedes infirmatus*, *Aedes vexans*, and *Culex nigripalpus* were captured on DESCs fitted with blue LEDs compared with red or IR LEDs. DESCs with blue LEDs captured significantly more *Culex erraticus* females than those with IR LEDs. Using resting boxes, 12 species from 5 genera (*Anopheles*, *Coquillettia*, *Culex*, *Mansonia*, and *Uranotaenia*) were collected. Resting boxes without LEDs captured 1,585 mosquitoes (22.2% of total). The fewest number of mosquitoes (16.7%) were collected from boxes affixed with the blue LEDs. Significantly more *Anopheles quadrimaculatus* females were aspirated from resting boxes fitted with red and IR LEDs than from those with blue or green LEDs, or from the unlit control. Blood-fed mosquitoes were recovered in highest numbers from unlit resting boxes, followed by resting boxes fitted with green, IR, and blue LEDs. *Culex erraticus* accounted for the majority of blood-fed mosquitoes followed by *Coquillettia perturbans*. No blood-fed mosquitoes were recovered from resting boxes fitted with red LEDs.

KEY WORDS Attraction, resting box, wavelength, Culicidae, vision

INTRODUCTION

Since the early 1900s, the effectiveness of techniques to attract and track the movements of hematophagous insects has continued to improve (Crans 1995). Adequate and reliable population sampling is frequently seen as the most important and most difficult step in ecological studies. Sampling outdoor mosquito populations is often difficult, because of behavioral differences between species and the fact that mosquitoes, in general, are not equally distributed within a habitat (Crans 1995). As a result, a variety of trapping systems have been developed that have been further enhanced with artificial, reflected, or filtered lights with great success (Barr et al. 1963, Ali et al. 1989, Service 1993, Burkett and Butler 2005, Hoel 2005). However, traps produce a bias when used in vector surveillance and population monitoring by primarily selecting nonblood-fed, host-seeking mosquitoes. Collections of resting mosquitoes yield a more accurate representative sample of a mosquito population, recovering not only nonblood-fed, host-seeking individuals, but also males, as well as blood-fed

and gravid females (Bradley 1943, Hayes et al. 1958, Loomis and Sherman 1959).

Mosquitoes often rest or seek shelter in protected sites (Crans 1995). The capitalization on this natural phenomenon has allowed researchers to effectively sample mosquitoes during inactive hours using artificial resting boxes (Goodwin 1942). Several laboratory and field studies have been conducted to determine the behavior of adult Diptera in response to visual stimuli, with special attention given to the modification of light wavelength and intensity in Culicidae (Huffaker and Back 1943, Fox 1958, Bidlingmayer 1967, Burkett and Butler 2005). Headlee (1937) first observed the impact of varying light intensities on catch size after noting that increased numbers of mosquitoes were attracted to within a certain proximity of traps. Previous studies have also demonstrated the effects of wavelength variation on field populations of *Culex* and *Psorophora*. Higher proportions of *Culex nigripalpus* Theobald, *Culex erraticus* Dyar and Knab, *Psorophora columbiae* Dyar and Knab, and *Psorophora ciliata* Fabricius were collected with New Jersey light traps modified with incandescent blue lights than with traps modified with yellow, orange, green, red, or white lights (Ali et al. 1989).

Much of what is known today concerning the affinity of Diptera to different light wavelengths can be credited to studies in which scientifically imprecise light sources were used (Brett 1938, Bracken et al. 1962, Bradbury and Bennett 1974, Browne and Bennett 1981, Allan and Stoffolano 1986). The recent development of super-bright

¹ Entomology and Nematology Department, PO Box 110620, Institute of Food and Agriculture Sciences, University of Florida, Gainesville, FL 32611-0620.

² United States Department of Agriculture–Agricultural Research Service, Center for Medical, Agricultural, and Veterinary Entomology, 1600 SW 23rd Drive, Gainesville, FL 32608.

³ To whom correspondence should be addressed. Phillip E. Kaufman Entomology and Nematology Department PO Box 110620 Gainesville, FL 32611-0620.

light-emitting diodes (LEDs) has enabled researchers to refine techniques to more effectively attract mosquitoes by using more precise light sources. For example, the Centers for Disease Control and Prevention (CDC) traps fitted with LEDs have greater light intensity with considerably lower energy requirement than the same traps in which incandescent bulbs are used, as is the normal practice (Burkett et al. 1998).

Although evaluations of trapping technology involving insect wavelength preferences and resting behavior have been studied extensively, the combination of the two has not. Few publications exist describing the attractiveness of LEDs to different mosquito species. Additionally, mosquito abundance in resting boxes when combined with a light source has not been published. Therefore, the objectives of this study were to determine wavelength preference of adult mosquitoes using LEDs, and to evaluate the attractiveness of resting boxes fitted internally with LEDs.

MATERIALS AND METHODS

All field trials were conducted at the University of Florida Horse Teaching Unit (HTU) and the Prairie Oaks subdivision (PO) near Gainesville, FL. Both locations were similar environments previously shown to have productive mosquito developmental sites (Butler, personal communication; Holton 2007). Both locations are adjacent to the Payne's Prairie State Preserve, and are surrounded by a mix of hardwood and pine forest with minimal undergrowth. The LEDs (Digi-Key Corporation, Thief River Falls, MN) used in these studies (part number, wavelength, and millicandela [mcd] rating, as described in Hoel [2005]), were blue (P466-ND, 470 nm, 650 mcd), green (67-1755-ND, 502 nm, 1,500 mcd), red (67-1611-ND, 660 nm, 1,800 mcd), and infrared (IR) (LN77L-ND, 860 nm). All LEDs consisted of a round lens and were 8.6 mm long by 5.0 mm in diam; viewing angles were 30° except for IR (20°). A 180-ohm resistor that was soldered to all LEDs restricted current flow to prevent mechanical failure.

Diode-equipped sticky card (DESC) studies were conducted using 20 × 20 cm wooden boxes with 4 sides and an open top and bottom constructed from exterior-grade plywood. Each of the 4 outside surfaces was painted with flat black paint (Valspar, Wheeling, IL). A 13 × 13 cm black sticky card (EPA no. 057296-WI-001, Atlantic Paste & Glue Corporation, Brooklyn, NY) was affixed to the exterior of each wall with 1 LED centered per vertical side. Black cards were selected to reduce variability of reflected light from LEDs. A hole drilled into the center of each sticky card and the box allowed for insertion of an LED from the inside of the DESC boxes so LEDs protruded outward through glue boards.



Fig. 1. Four sided, light-emitting diode-equipped sticky card box. One 13 × 13 cm sticky card and one diode were affixed to each exterior wall of box.

Boxes were positioned 90 cm above ground (Fig. 1). Every 24 h the outward-facing sides of each DESC box were fitted with a randomly selected LED and a new sticky card, resulting in 32 light exposures per day (8 boxes × 4 sides).

Diode-equipped resting boxes (DERB) with 4 sides, a back wall, and an open front were constructed using the specifications of a standard 30-cm³ resting box, as described by Crans (1995) using exterior-grade plywood (Fig. 2a). Box exteriors were painted with flat black, and interiors were painted barn red (Valspar, Wheeling, IL) (Fig. 2b). A hole drilled through the back center wall allowed for the insertion of an LED. For each DERB trial, 4 sites contained 5 treatments, 1 of 4 LED colors (described previously) and an unlit control, resulting in a total of 5 resting boxes per site, 20 resting boxes per location (HTU and PO). Placement of resting boxes within a site was randomized at the start of the trial, and treatments (LED color) were randomized daily wherein LEDs were moved from box to box. Boxes were staged in a slightly offset line so that boxes in front did not disrupt the passage of the LED emission from those to the rear. All boxes were out of direct sunlight, and approximately 4 m apart, with the open sides facing west.

Three CO₂-baited CDC light traps (model 512; John W. Hock Company, Gainesville, FL) with




Fig. 2. Resting boxes. (A) Rear view of resting box showing protective diode housing. (B) Resting boxes were painted interiorly barn red and the exterior was black.

the incandescent bulb removed were operated simultaneously with the DESC and DERB trials and were used to provide a measure of background mosquito populations at each trial location. One CDC trap was placed at the HTU location and 2 were placed at the PO location. All traps were set 120 cm above ground using a shepherd's hook. Carbon dioxide was provided from a 9-kg compressed gas cylinder at a flow rate of 250 ml/min using a 5-psi single-stage regulator equipped with an inline microregulator (no. 007) and an inline filter (Clarke Mosquito Control, Roselle, IL). Flow rates were confirmed

using a Gilmont Accucal® flowmeter (Gilmont Instrument Company, Barrington, IL).

Power to DESC LEDs and CDC traps were provided by 6 v, 12 ampere-hour (A-h) rechargeable gel cell batteries, which were changed every 24–48 h. Resting-box LEDs were powered by 4 rechargeable AA batteries, which were changed every 5 days. Temperature and humidity conditions were obtained from the Gainesville Regional Airport weather station through the online NOAA database (NOAA 2008).

Resting-box and sticky-card trials at both locations occurred in 20 trap-night clusters between July and September 2006, and between May and September 2007 for a total of between 120 and 157 trap nights, depending upon location and trial type (DESC or DERB). Twenty-four hours after deployment (1 trap night), sticky cards and CDC collection nets were removed and mosquitoes were aspirated from boxes. Treatments were randomized and traps reset. Mosquitoes collected in traps were brought back to the laboratory where they were counted by species and sex using the dichotomous keys of Darsie and Morris (2003).

Mosquito preference for LED wavelengths was evaluated using a multi-factorial ANOVA (SAS Institute 2001). For analysis, all data were normalized using the SQRT ($n + 1$) transformation; however, actual values are provided in text and tables. The model included the fixed effects of location, site, and LED treatment; the interaction term, location \times LED treatment and the random effect trial. In instances where either the interaction term or the trial effect was significant, the data were analyzed separately by location or trial. Tukey's standardized test ($\alpha = 0.05$) was used to separate treatment means.

RESULTS

Diode-equipped sticky card boxes

In the DESC trial 452 mosquitoes, including 15 mosquito species from 7 genera, were captured from both locations (Table 1). Significantly more female *Aedes infirmatus* Dyar and Knab ($F_{3, 2,864} = 3.49$; $P = 0.0150$), *Aedes vexans* Meigen ($F_{3, 2,544} = 4.00$; $P = 0.0074$) and *Cx. nigripalpus* Theobald ($F_{3, 2,544} = 4.66$; $P = 0.0030$) were captured on DESCs fitted with blue diodes than on those with red or IR diodes (Table 2). Numerically, DESCs affixed with IR diodes caught the fewest female *Ae. infirmatus*, *Ae. vexans*, and *Cx. nigripalpus*.

Coquillettidia perturbans males were recovered in significantly higher numbers on DESCs containing green diodes on than those with the blue or IR diodes ($F_{3, 2,226} = 3.86$; $P = 0.0091$). Significantly more *Cq. perturbans* female mosquitoes were captured on DESCs with green diodes

Table 1. Relative total mosquito collection abundance from Centers for Disease Control and Prevention (CDC), diode-equipped sticky card, and resting-box trapping systems at 2 locations near Gainesville, FL.

Mosquito species	Trap type ¹		
	CDC ²	DESC ³	RB ⁴
<i>Aedes albopictus</i> ♂	0	1	0
<i>Aedes albopictus</i> ♀	174	0	0
<i>Aedes canadensis</i> ♀	6	0	0
<i>Aedes fulvus pallens</i> ♀	18	0	0
<i>Aedes infirmatus</i> ♂	0	1	0
<i>Aedes infirmatus</i> ♀	6,835	16	1
<i>Aedes sollicitans</i> ♀	7,330	0	0
<i>Aedes taeniorhynchus</i> ♀	18	0	0
<i>Aedes triseriatus</i> ♂	0	1	0
<i>Aedes triseriatus</i> ♀	96	11	1
<i>Aedes vexans</i> ♂	0	4	0
<i>Aedes vexans</i> ♀	10,577	22	0
<i>Anopheles crucians</i> ♂	0	1	15
<i>An. crucians</i> ♀	6,663	2	8
<i>An. quadrimaculatus</i> ♂	0	1	82
<i>An. quadrimaculatus</i> ♀	269	1	90
<i>Coquillettidia perturbans</i> ♂	0	52	218
<i>Cq. perturbans</i> ♀	63,833	80	109
<i>Culex erraticus</i> ♂	0	4	3,455
<i>Cx. erraticus</i> ♀	19,773	21	2,985
<i>Cx. nigripalpus</i> ♂	0	13	6
<i>Cx. nigripalpus</i> ♀	52,885	23	22
<i>Cx. quinquefasciatus</i> ♀	30	0	0
<i>Cx. salinarius</i> ♂	0	9	27
<i>Cx. salinarius</i> ♀	4,168	8	13
<i>Cx. territans</i> ♂	0	4	21
<i>Cx. territans</i> ♀	0	3	38
<i>Mansonia titillans</i> ♂	0	9	14
<i>Ma. titillans</i> ♀	20,720	69	17
<i>Psorophora ciliata</i> ♀	32	0	0
<i>Psorophora columbiae</i> ♀	1,485	3	0
<i>Psorophora ferox</i> ♀	412	0	0
<i>Uranotaenia lowii</i> ♂	64	37	1
<i>Ur. lowii</i> ♀	0	6	23
<i>Ur. sapphirina</i> ♂	704	50	4
<i>Ur. sapphirina</i> ♀	1,485	3	0

¹ CDC = Modified CDC light-trap; DESC = diode-equipped sticky-card box; RB = resting box.

² Trapping occurred simultaneously with DESC and RB trials, 380 trap nights.

³ Trapping occurred from 16 August to 27 September 2006 and from 5 May to 13 September 2007, 260 trap nights.

⁴ Trapping occurred from 21 July to 14 August 2006, and from 5 May to 26 September 2007, 448 trap nights.

than on DESCs fitted with red or IR diodes ($F_{3, 2,864} = 4.66$; $P = 0.0003$) (Table 2). *Mansonia titillans* females were only captured at the HTU location in 2006. These mosquitoes preferred DESCs fitted with either blue or green LEDs over those with red or IR LEDs ($F_{3, 313} = 6.18$; $P = 0.0004$) (Table 2). Trap comparisons for all other species were not significant.

Approximately 91,766 female mosquitoes were captured using modified CDC traps over 360 trap nights. Overall, 21 species from 7 genera were captured (Table 1).

Resting boxes

A total of 7,157 adult mosquitoes, including 12 species from 6 genera, were recovered from resting boxes (Table 1). Although no significant

differences were observed, 23% and 22% of all mosquitoes were recovered in IR and no-light DERBs, respectively. The fewest mosquitoes (16.7%) were recovered in blue DERBs. Significantly more *Anopheles quadrimaculatus* females were aspirated from resting boxes fitted with red LEDs than from all other treatments ($F_{4, 6,315} = 2.47$; $P = 0.0429$) (Table 3).

The year effect was significant for *Cx. erraticus* males and females ($F_{4, 1,126} = 2.40$; $P = 0.0476$). During 2006, no preferences were observed among diode treatments for either sex of this species. However, in 2007, significantly more *Cx. erraticus* females were aspirated from resting boxes fitted with blue, green, and red LEDs, and from no-light boxes compared with boxes containing IR LEDs ($F_{4, 5,577} = 8.41$; $P < 0.0001$) (Table 3).

Table 2. Mean (SE) number of mosquitoes per trap night attracted to light-emitting diode wavelengths affixed to sticky cards positioned externally on wooden boxes during 24-h trapping intervals at 2 locations near Gainesville, FL.

Species	n ²	Mean (SE) ¹			
		Blue ³	Green ³	Red ³	Infrared ³
<i>Aedes infirmatus</i> ♀	720	0.014 (0.004)a	0.004 (0.002)ab	0.003 (0.002)b	0.001 (0.001)b
<i>Aedes vexans</i> ♀	640	0.019 (0.006)a	0.013 (0.004)ab	0.002 (0.002)b	<0.001 (<0.001)b
<i>Coquillettidia perturbans</i> ♂	560	0.014 (0.006)b	0.043 (0.010)a	0.011 (0.006)ab	0.016 (0.005)b
<i>Coquillettidia perturbans</i> ♀	720	0.031 (0.007)ab	0.049 (0.009)a	0.024 (0.007)b	0.008 (0.003)b
<i>Culex erraticus</i> ♀	320	0.034 (0.012)a	0.016 (0.007)ab	0.013 (0.006)ab	0.003 (0.003)b
<i>Culex nigripalpus</i> ♀	640	0.019 (0.007)a	0.016 (0.005)ab	0.002 (0.002)b	<0.001 (<0.001)b
<i>Mansonia titillans</i> 2006 ♀	80	0.313 (0.068)a	0.363 (0.110)a	0.100 (0.038)b	0.050 (0.025)b

¹ Means within rows followed by the same letter were not significantly different ($P < 0.05$, Tukey's standardized test [SAS Institute 2001]).

² n = number of trap nights where total mosquito species capture was ≥ 1 mosquito per 20 trap nights.

³ Blue diode = 470 nm, green diode = 502 nm, red diode = 660 nm, infrared diode = 860 nm.

Notable differences between treatment responses were observed among the 177 blood-fed mosquitoes recovered from resting boxes. Unlit resting boxes (control) accounted for the greatest number of blood-fed mosquitoes recovered, followed by resting boxes fitted with green, IR, and blue LEDs (Table 4). No blood-fed mosquitoes were recovered from resting boxes fitted with red LEDs. Among mosquito species, 74% of all blood-fed mosquitoes recovered were *Cx. erraticus*.

Approximately 100,653 female mosquitoes were trapped in the 3 CDC traps over 448 trap nights. Overall, 21 species from 8 genera were captured (Table 1). CDC traps were functional for >91% of trap nights.

DISCUSSION

Overall, capture of mosquitoes on DESCs was greatest with blue LEDs, followed by green, red, and IR LEDs. Similarly, previous studies have shown host-seeking mosquito wavelength preference to be in the blue-green range (400–600 nm), with diminishing attraction as wavelengths in-

crease in length (>600 nm) (Ali et al. 1989, Burkett et al. 1998).

Our findings agree with Bargren and Nibley's (1956) observations, where considerably more *Ae. vexans* were captured on DESCs fitted with blue diode treatments. However, we observed no significant differences in wavelength preference for *Culex salinarius*. In contrast to Burkett et al. (1998), considerably more *Cx. nigripalpus* were captured on DESCs fitted with blue diodes than on those with red diodes. These results suggest a spectral sensitivity for *Cx. nigripalpus* females at the higher end of the blue spectrum (>450 nm), with decreasing sensitivity for wavelengths from 450 nm toward the lower end of the red spectrum (<640 nm).

Browne and Bennett (1981) and Ali et al. (1989) determined that shorter, blue-green wavelengths (400–600 nm and 430–550 nm, respectively) attracted significantly more *Cq. perturbans* than did longer wavelengths. These results were comparable to our observations where significantly more *Cq. perturbans* were captured on DESCs affixed with lower-spectrum green diodes (502 nm), and fewer were captured on DESCs

Table 3. Mean (SE) mosquitoes/trap night attracted to 4 light-emitting diode wavelengths placed in resting boxes near Gainesville, FL.

Species	n ³	Mean (SE) ^{1,2}				No Light
		Blue	Green	Red	Infrared	
<i>Anopheles quadrimaculatus</i> 2006–2007 ♀	1,268	0.004 (0.003)b	0.013 (0.007)b	0.032 (0.013)a	0.007 (0.005)b	0.015 (0.005)b
<i>Culex erraticus</i> 2006 ♂	288	2.154 (0.333)	2.452 (0.396)	3.009 (0.453)	3.868 (0.617)	3.154 (0.501)
<i>Culex erraticus</i> 2007 ♂	1,120	0.022 (0.005)	0.028 (0.006)	0.024 (0.005)	0.021 (0.005)	0.032 (0.007)
<i>Culex erraticus</i> 2006 ♀	148	2.851 (0.558)	2.980 (0.546)	3.223 (0.616)	3.967 (0.710)	3.932 (0.740)
<i>Culex erraticus</i> 2007 ♀	1,120	0.113 (0.012)a	0.087 (0.011)a	0.083 (0.010)a	0.038 (0.007)b	0.104 (0.012)a

¹ Means within rows followed by the same letter were not significantly different ($P < 0.05$, Tukey's standardized test [SAS Institute 2001]). Means in rows without means separation letters were not significantly different.

² Blue diode = 470 nm, green diode = 502 nm, infrared diode = 860 nm, red diode = 660 nm, and no light indicates no diode treatment.

³ n = number of trap nights in which total mosquito species captured was ≥ 1 per 20 trap nights. Trapping occurred from July 2006 to September 2007.

Table 4. Total blood-fed mosquitoes attracted to light-emitting diodes of 4 different wavelengths placed in resting boxes compared with resting boxes without lights near Gainesville, FL.

Mosquito species	Diode treatment ¹					Total
	Blue	Green	Red	Infrared	No light	
<i>Anopheles crucians</i>	0	3	0	2	2	7
<i>Anopheles quadrimaculatus</i>	0	0	0	2	0	2
<i>Coquillettidia perturbans</i>	7	9	0	5	7	28
<i>Culex erraticus</i>	26	33	0	30	43	132
<i>Culex salinarius</i>	1	4	0	1	1	7
<i>Culex territans</i>	1	1	0	0	1	3
Total	35	50	0	40	54	179

¹ Blue diode = 470 nm, green diode = 502 nm, red diode = 660 nm, infrared diode = 860 nm, and no light indicates no diode treatment.

fitted with higher-spectrum IR (860-nm) diodes. Through the use of more scientifically accurate LEDs, our results demonstrated a stronger preference of *Cq. perturbans* for green wavelengths (502 nm) than for blue (470 nm), suggesting wavelength attraction of this species was nearer to the green range of the blue-green spectrum (>500 nm).

Ali et al. (1989) reported higher capture rates for *Cx. erraticus* when using blue-colored bulbs (430–490 nm) compared with red-colored bulbs (620–720 nm) of comparable intensity. Similarly, we captured significantly more *Cx. erraticus* females on DESCs affixed with blue diodes than on DESCs fitted with IR diodes, but observed no significant preferences between blue, green, or red diodes. However, we recovered more male *Cx. erraticus* (26%) from resting boxes fitted with IR LEDs, and more females (23%) from unlit boxes. Therefore, wavelength preferences for host-seeking *Cx. erraticus* may range in the lower bands of the spectrum (<470 nm), whereas resting *Cx. erraticus* may be averse to these wavelength bands.

No blood-fed *An. quadrimaculatus* were recovered from DERBs fitted with red LEDs, suggesting a physiological stage-dependent behavior. Therefore, the addition of >660-nm LEDs to resting boxes may enhance efficacy of sampling host-seeking *An. quadrimaculatus* populations, whereas their inclusion may repel blood-fed *An. quadrimaculatus*.

Although not significant, the trends observed in DERB trials suggest that resting mosquitoes may choose the higher wavelengths of light and dark boxes over the lower wavelengths of light that attract host-seeking mosquito species. Lower wavelengths are preferred for phototactic, host-seeking mosquito species trapped at night, such as *Cx. erraticus*, *Cx. nigripalpus*, and *Ps. columbiae* (Bargren and Nibley 1956, Ali et al. 1989, Burkett et al. 1998, Burkett and Butler 2005). These behavioral differences between host-seeking mosquitoes and those captured in resting boxes may be the product of variation in wavelength attraction between the physiological

stages and deserves further attention. Additionally, our use of narrow wavelengths may have excluded mosquitoes preferring longer or shorter wavelength bands than those selected.

Although no blood-fed mosquitoes were captured using DESCs, 179 blood-fed mosquitoes were recovered from resting boxes. *Culex erraticus* was the most common blood-fed mosquito recovered, accounting for 74% of all blood-fed mosquitoes captured. This high recovery rate is likely because of the elevated presence of *Cx. erraticus* among all mosquito species recovered from resting boxes throughout this study. Among treatments, 43 *Cx. erraticus* were recovered from resting boxes fitted with IR LEDs. Although resting boxes fitted with IR LEDs recovered the highest number of blood-fed mosquitoes, it is important to note that no blood-fed mosquitoes were recovered from resting boxes fitted with red LEDs. Our observations may suggest a variation in wavelength preferences among host-seeking and blood-fed mosquitoes. These findings provide important information that warrants further investigation into behavioral phototaxis differences among physiological stages of mosquitoes.

The results of our study suggest that, in the absence of alternative host stimuli, wavelengths in the lower green (502 nm) spectral range would be optimal for targeting a broad range of mosquito species. Additionally, the use of LEDs as opposed to wavelength filters or colored bulbs provides a more precise and energy efficient delivery system for spectrally sensitive insects.

Although the CO₂-baited CDC trap captured considerably more host-seeking mosquitoes, the DERB and DESC provide an alternative trapping system. Given their accuracy in specific wavelength achievement, small size, and minimal power usage, LEDs can be used as light sources for various trap designs where access and equipment to target sites are limited. Their demonstrated effectiveness for attracting mosquitoes without the aid of supplemental host attractants further eliminates the need for and costs of heavy tanks (CO₂) or volatile chemicals

(lactic acid, octenol). The durability of the LED-based equipment also reduces time-consuming maintenance.

ACKNOWLEDGMENTS

The authors thank J. Butler, S. TenBroeck, J. Calahan, and the residents of the Prairie Oaks Subdivision for allowing access to their properties. The authors would also like to acknowledge J. Pitzer and P. Obenauer for their involvement and assistance. This study was supported by the University of Florida Agricultural Experiment Station federal formula funds, Project No. FLA-04598, received from Cooperative State Research, Education, and Extension Service, US Department of Agriculture. Thank you to J. Cilek for reviewing a draft of this publication.

REFERENCES CITED

- Ali A, Nayar JK, Knight JW, Stanley BH. 1989. Attraction of Florida mosquitoes (Diptera: Culicidae) to artificial light in the field. *Proc Calif Mosq Vector Control Assoc* 57:82–88.
- Allan SA, Stoffolano JG. 1986. Effects of hue and intensity on visual attraction of adult *Tabanus nigrovittatus* (Diptera: Tabanidae). *J Med Entomol* 23:83–91.
- Bargren WC, Nibley C. 1956. Comparative attractiveness of colored lights of equal intensity to specific species of mosquitoes. 3rd Area Med Lab, Fort McPherson, GA.
- Barr AR, Smith TA, Boreham M, White KE. 1963. Evaluation of some factors affecting the efficiency of light traps on collecting mosquitoes. *J Econ Entomol* 56:123–127.
- Bidlingmayer WL. 1967. A comparison of trapping methods for adult mosquitoes: species response and environmental influence. *J Med Entomol* 4:200–220.
- Bracken GK, Hanec W, Thorsteinson AJ. 1962. The orientation of horseflies and deerflies (Tabanidae: Diptera). II. The role of some visual factors in the attractiveness of decoy silhouettes. *Can J Zool* 40:685–695.
- Bradbury WC, Bennett GF. 1974. Behavior of adult *Simuliidae* (Diptera). I. Response to color and shape. *Can J Zool* 52:251–259.
- Bradley GH. 1943. Determination of densities of *Anopheles quadrimaculatus* on the wing. *Proc Annu Meet N J Mosq Extermination Assoc* 30:22–27.
- Brett GA. 1938. On the relative attractiveness to *Aedes aegypti* of certain coloured cloths. *Trans R Soc Trop Med Hyg* 32:113–124.
- Browne SM, Bennett GF. 1981. Response of mosquitoes (Diptera: Culicidae) to visual stimuli. *J Med Entomol* 18:505–521.
- Burkett DA, Butler JF. 2005. Laboratory evaluation of colored light as an attractant for female *Aedes aegypti*, *Aedes albopictus*, *Anopheles quadrimaculatus* and *Culex nigripalpus*. *Fla Entomol* 88:383–389.
- Burkett DA, Butler JF, Kline DL. 1998. Field evaluation of colored light-emitting diodes as attractants for woodland mosquitoes and other Diptera in north central Florida. *J Amer Mosq Control Assoc* 14:186–195.
- Crans WJ. 1995. Resting boxes as mosquito surveillance tools. *Proc New Jersey Mosq Control Assn* 82:53–57.
- Darsie RF, Morris CD. 2003. Keys to the adult females and fourth instar larvae of the mosquitoes of Florida (Diptera, Culicidae). Volume 1. De Leon Springs, FL: E. O. Painter Printing Co.
- Fox I. 1958. The mosquitoes of the international airport, Isla Verde, Puerto Rico, as shown by light traps. *Mosq News* 18:117–124.
- Goodwin MN. 1942. Studies on artificial resting places of *Anopheles quadrimaculatus* Say. *J Natl Malaria Soc* 1:93–99.
- Hayes RO, Bellamy RE, Reeves WC, Willis MJ. 1958. Comparison of four sampling methods for measurement of *Culex tarsalis* adult populations. *Mosq News* 20:174–178.
- Headlee TJ. 1937. Some facts underlying the attraction of mosquitoes to sources of radiant energy. *J Econ Entomol* 30:309–312.
- Hoel D. 2005. *Response of Aedes albopictus (Diptera: Culicidae) to traps, attractants, and adulticides in north central Florida*. [Ph.D. dissertation]. University of Florida, Gainesville, FL.
- Holton AC. 2007. *Evaluation of different commercial lures and horse odors as an attractant and their abilities to increase mosquito trap numbers at the University of Florida Horse Teaching Unit*. [M.S. thesis]. University of Florida, Gainesville, FL.
- Huffaker CB, Back RC. 1943. A study of methods of sampling mosquito populations. *J Econ Entomol* 36:561–569.
- Loomis EC, Sherman EJ. 1959. Comparison of artificial shelters and light traps for measurement of *Culex tarsalis* and *Anopheles freeborni* populations. *Mosq News* 19:232–237.
- NOAA [National Oceanic and Atmospheric Administration]. 2008. *Gainesville weather information 2006–2007*. [Internet]. Gainesville, FL: NOAA. [accessed September 3, 2007]. Available from: <http://forecast.weather.gov/MapClick.php?CityName=Gainesville&state=FL&site=JAX&textField1=29.6742&textField2=-82.3363&e=0>.
- SAS Institute. 2001. *SAS/STAT user's manual*, Version 8.2. Cary, NC: SAS Institute.
- Service MW. 1993. *Mosquito ecology: field sampling methods*. 2nd ed. London: Elsevier Applied Science.