

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Publications from USDA-ARS / UNL Faculty

U.S. Department of Agriculture: Agricultural
Research Service, Lincoln, Nebraska

3-2011

Comparison of various configurations of CDC-type traps for the collection of *Phlebotomus papatasi* Scopoli in southern Israel

Daniel L. Kline

United States Department of Agriculture-ARS-Center for Medical

Jerome Hogsette

United States Department of Agriculture-ARS-Center for Medical

Gunter Muller

Kuvin Centre for the Study of Infectious and Tropical Diseases

Follow this and additional works at: <http://digitalcommons.unl.edu/usdaarsfacpub>



Part of the [Agricultural Science Commons](#)

Kline, Daniel L.; Hogsette, Jerome; and Muller, Gunter, "Comparison of various configurations of CDC-type traps for the collection of *Phlebotomus papatasi* Scopoli in southern Israel" (2011). *Publications from USDA-ARS / UNL Faculty*. 1020.

<http://digitalcommons.unl.edu/usdaarsfacpub/1020>

This Article is brought to you for free and open access by the U.S. Department of Agriculture: Agricultural Research Service, Lincoln, Nebraska at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Publications from USDA-ARS / UNL Faculty by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Comparison of various configurations of CDC-type traps for the collection of *Phlebotomus papatasi* Scopoli in southern Israel

Daniel L. Kline¹, Jerome A. Hogsette¹, and Günter C. Müller²

¹United States Department of Agriculture-ARS-Center for Medical, Agricultural, and Veterinary Entomology, Gainesville, FL, U.S.A.

²Department of Microbiology and Molecular Genetics, IMRIC, Kuvim Centre for the Study of Infectious and Tropical Diseases, Faculty of Medicine, Hebrew University, Jerusalem, Israel, 91120

Received 22 June 2010; Accepted 21 January 2011

ABSTRACT: We conducted two experiments to determine the best CDC-trap configuration for catching male and female *Phlebotomus papatasi*. First, visual features were evaluated. Standard CDC traps were modified to have black or white catch bags, black or white lids, or no lids and these were tried in different combinations. Significantly more male sand flies were caught by darker traps; significantly more females were captured by traps with either all black or a combination of black and white features. Attraction may be due to dark color or contrast in colors. CDC traps with suction and the following features were also evaluated: no light; incandescent light; ultraviolet (UV) light; combination of black color, heat and moisture; CO₂ alone, or a combination of black color, heat, moisture, and CO₂ simultaneously, all in upright and inverted positions, with the opening for insect entry always 50 cm above the ground. Significantly more females than males were caught by all traps (standard and inverted) except the control traps with suction only. Traps with CO₂ caught more sand flies than traps without CO₂. Traps with black color, heat and moisture captured significantly more sand flies than the control traps, but with the addition of CO₂, these traps catch significantly more sand flies than the other traps evaluated. Inverting traps increased the catch for like traps by about two times. **Journal of Vector Ecology 36 (Supplement 1): S212-S218. 2011.**

Keyword Index: sand flies, visual targets, UV light, inverted traps, carbon dioxide, heat, color.

INTRODUCTION

Phlebotomine sand flies have a wide distribution, though mainly in the tropics and subtropics (Lane 1993). Towards the north, they reach south west Canada (Young and Perkins 1984), and in the south they are found until latitude 40°S (Killick-Kendrick 1999). Both female and males are dependent on sugar as an energy source (Schlein and Warburg 1986), but females also need additional blood meals for egg production every few days (Killick-Kendrick 1999). This is the reason for frequent contacts between vector and host, and why phlebotomine sand flies are such a nuisance as well as vectors of numerous diseases (Comer and Tesh 1991; Ashford 2001, Birtles 2001).

The dynamics of sand fly attraction to hosts is rather complex and little is known compared to mosquitoes and other biting flies (Gibson and Torr 1999). As for most biting flies, carbon dioxide (CO₂) is the single most potent attractant for sand flies (Pinto et al. 2001), but they are also attracted by host odors alone as shown in several experiments in the laboratory and field (Killick-Kendrick et al. 1986, Morton and Ward 1989, Dougherty et al. 1999). Moreover, temperature and humidity gradients seem to play a role in sand fly attraction (Nigam and Ward 1991) and there is also evidence that optical aspects are important for host detection (Mellor et al. 1996).

Almost every attempt to study natural behavior or applied control of sand flies in the field involves sampling

the population in one way or another. For adult sand flies, either small CDC-like traps or sticky traps (sheets of paper or plastic covered with a viscous adhesive such as motor oil) are commonly used (Alexander 2000). Non-attractive traps, like simple sticky papers and unlighted, unbaited CDC traps only catch flies from their immediate area and accordingly, tend to yield relatively low numbers of sand flies. Several productive trapping methods and collecting procedures have been standardized for sampling sand flies. Selection of an appropriate method depends on the objectives and type of study to be performed, species, sex or physiological state of the insects required, and any constraints on preservation and transportation of the specimens (Service 1993, Alexander 2000).

A more active and selective way is to attract biting flies, in most cases females, to all kind of baits including animals and humans (Sharp et al. 1984, Andrade et al. 2008), as well as elements of them like clothes, hair, urine, feces, etc. (Allan et al. 2006, Kline 1998). More commonly, isolated or combined attractive features, like CO₂, visual targets, chemical lures, heat, moisture, and movement, are now used to increase trap catches (Kline 2006, Bernier et al. 2003, Murphy et al. 2001). Light, especially in the long-wave ultraviolet (UV) range, is generally regarded as an attractant. However, it often causes disorientation of nocturnally active flying insects (Nowinszky 2004). With their orientation thus compromised, both male and female sand flies are drawn towards the direction of the light source and are unable to

Report Documentation Page

Form Approved
OMB No. 0704-0188

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE JUN 2010		2. REPORT TYPE		3. DATES COVERED 00-00-2010 to 00-00-2010	
4. TITLE AND SUBTITLE Comparison of various configurations of CDC-type traps for the collection of Phlebotomus papatasi Scopoli in southern Israel				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) U.S. Department of Agriculture -Agricultural Service (USDA/ARS),Center for Medical, Agricultural and Veterinary Entomology,1600 SW 23rd Dr,Gainesville,FL,32608				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT We conducted two experiments to determine the best CDC-trap configuration for catching male and female Phlebotomus papatasi. First, visual features were evaluated. Standard CDC traps were modified to have black or white catch bags, black or white lids, or no lids and these were tried in different combinations. Significantly more male sand flies were caught by darker traps; significantly more females were captured by traps with either all black or a combination of black and white features. Attraction may be due to dark color or contrast in colors. CDC traps with suction and the following features were also evaluated: no light; incandescent light; ultraviolet (UV) light; combination of black color, heat and moisture; CO2 alone, or a combination of black color, heat, moisture, and CO2 simultaneously, all in upright and inverted positions, with the opening for insect entry always 50 cm above the ground. Significantly more females than males were caught by all traps (standard and inverted) except the control traps with suction only. Traps with CO2 caught more sand flies than traps without CO2. Traps with black color, heat and moisture captured significantly more sand flies than the control traps, but with the addition of CO2, these traps catch significantly more sand flies than the other traps evaluated. Inverting traps increased the catch for like traps by about two times. Journal of Vector Ecology 36 (Supplement 1): S212-S218. 2011.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

avoid the capture mechanisms of traps.

A large body of literature exists in which numerous methods and trap designs for the collection of sand flies are discussed (for a review see Alexander 2000). However, little attention has been paid to the fact that a few small changes in the design and presentation of CDC-like traps might increase the catch size and change the sex ratio. This is important if data from different areas need to be compared or if sex ratios are used as an indication of possible breeding sites (Felicangeli 2004). The purpose of this study was to evaluate the impact of selected modifications of the standard CDC trap on the number and sex ratio of *P. papatasi* captured.

MATERIALS AND METHODS

Study site

The study was carried out in mid-autumn near Jericho, about 10km north of the Dead Sea, at an altitude of about 300 m below sea level. This region is an extreme desert and belongs to the Saharo-Arabian phyto-geographical zone (Danin 1988). The annual precipitation of 50 to 100 mm is restricted to short winter rains and average daily temperature ranges from ca. 20° C between late September and April to >30°C from May through August (Ashbel 1951). The traps were evaluated in a neglected date plantation where *P. papatasi* is the dominant sand fly species and others, like *P. sergenti*, are rare or absent (Faiman et al. 2009, Müller and Schlein 2004, Schlein et al. 2001).

The experiments were conducted in the dry autumn when the annual winter and spring vegetation was already dry. About 20% of the remaining natural vegetation inside the plantation was scattered shrubs and semi-shrubs, including *Suaeda asphaltica* (Boiss.), *S. fruticosa* Forsk., *Atriplex halimus* (L), *A. leucoclada* Boiss. (Chenopodiaceae) and *Prosopis farcta* (Macbride) (Mimosaceae). Along the periphery of the oasis, groups of *Tamarix nilotica* (Ehrenb.) (Tamaricaceae) Bge. trees and shrubs, like *Alhagi graecorum* Boiss. (Papilionaceae) and *Salsola tetrandia* (Chenopodiaceae) Forssk., were restricted to small water catchments. No flowering plants, honeydew or honeydew-producing insects of any kind were found in the area at the time of the experiments.

Traps and experimental description

Efficacy of the modified CDC traps was evaluated in two experiments conducted from early September to early October, 2006. The modified traps (based on the CDC trap model 512, John Hock, Gainesville, FL, U.S.A.) were operated simultaneously and continuously, along an unpaved road crossing the plantation, with a distance of 20 m between each trap location. Traps were hung on bamboo tripods so the opening for insect entry was 50 cm above the ground (in both upright and inverted configurations). Traps were rotated clockwise between the trap locations at 17:00 daily to eliminate positional bias. Insects captured in traps during the night were removed at 07:00. Traps were powered by 6 volt motorcycle batteries which were recharged daily

with a generator.

Experiment I-Visual Features

Traps were evaluated as visual targets by illuminating them with incandescent bulbs and operating them in the upright (normal) position with variations in lid presence/absence and lid and catch bag color. Catch bags were used either in their original white color or were stained black with a commercial textile dye determined to have no repellent effects. Trap lids were used either in their original black color or were painted white. Stained and painted catch bags and lids, respectively, and unaltered catch bags and lids were submerged in a clear outdoor water pond 1 mo prior to use to eliminate any possible odors from the stain and paint. The list of modified traps is as follows: 1) White catch bag/no lid; 2) White catch bag/white lid; 3) White catch bag/black lid; 4) Black catch bag/no lid; 5) Black catch bag/white lid; 6) Black catch bag/black lid.

Experiment II-Trap Orientation

Traps, either upright or inverted, were evaluated in combination with selected attraction features. For a control, we used suction only with no light. To evaluate the effect of incandescent light, the original CDC light trap configuration was maintained. To evaluate the effect of UV light, the incandescent bulb was replaced with a small portable money checker (Tragbarer Geldschein-Prüfer mit Leuchte, model 751778 – 62, Conrad Electronics, Munich, Germany) equipped with a 4 watt, 6 volt UV tube attached horizontally, 3 cm above the opening for insect entry on the trap body (similar to the CDC Model 1212, John Hock, Gainesville FL, U.S.A). The UV unit was connected to a separate 6 volt motorcycle battery.

Heat was generated by heat film (Westham Innovations LTD., Tel Aviv, Israel) placed beneath a metal jacket of 4 mm iron sheet which fit tightly around the entire trap body. The modified trap bodies were then covered with a heavy non-glossy black paper. The surface temperature of the covered trap bodies, which was set at 41° C, was verified with an infrared thermometer (CEM DT8862 Professional 12:1 IR Infrared Dual Laser Thermometer, Meter Shack) gun. Moisture was supplied from sheets of 80 x 80 cm filter papers folded fan-like, with their tightly folded side inserted in beakers of water (Müller and Schlein 2006). Traps equipped with CO₂ used a bottled supply with a flow rate of 250 ml/min. The CO₂ lines were affixed to the body of the traps so CO₂ was released into the airflow 5 cm above the opening for insect entry.

The combinations of evaluated features were as follows: 1) Upright trap with suction only as control; 2) Inverted trap with suction only as control; 3) Upright trap with UV light; 4) Inverted trap UV light; 5) Upright trap with black body, heat and moisture; 6) Inverted trap with black body, heat and moisture; 7) Upright trap with black body, heat moisture, and CO₂; 8) Inverted trap with black body, heat moisture, and CO₂; 9) Upright trap with CO₂ only; 10) Inverted trap with CO₂ only.

Table 1. Mean *P. papatasi* adults captured on CDC traps modified to have black or white lids and catch bags ($n = 20$).

Trap	Sand flies captured ¹
Bag Black-Lid Black	14.5 ± 1.7a
Bag White-Lid Black	12.1 ± 1.6ab
Bag Black-Lid White	11.3 ± 1.7ab
Bag Black-No Lid	10.7 ± 1.5ab
Bag White-No Lid	9.0 ± 0.8b
Bag White-Lid White	5.5 ± 0.5c

¹Means followed by the same letter are not significantly different ($P < 0.05$; Ryan-Einot-Gabriel-Welsch Multiple Range Test [SAS Institute 2003]).

Statistical analysis

Data were first normalized by conversion to $\log_{10}(n+1)$ then subjected to ANOVA (SAS 2003) using the following main effects model statements: Total = Treatment Sex Replication, where the dependent variable represented numbers of sand flies captured, Treatment was one of the modified traps, Sex was either male or female, and Replication was an indication of trap location on one of the consecutive trapping days of each study. Means were separated with the Ryan-Einot-Gabriel-Welsch Multiple Range Test (REGWQ), and unless otherwise stated, $P < 0.05$ (SAS 2003). Although $\log_{10}(n+1)$ values were used for the analyses, actual values are reported in the text, figures and tables.

RESULTS

Experiment I-Visual Features

The main effects model was significant for the dependent variable ($F=8.54$, d.f.=15,119, $P<0.0001$). Means for the total numbers of flies captured ranged from 14.5 to 5.5 and overall, traps captured significantly more females

than males.

There were no significant differences between the mean numbers of sand flies captured by any of the three traps having black catch bags, and the trap with white catch bag and black lid (Table 1). The all-black trap caught significantly more sand flies than the trap with no lid and a white catch bag, and the all-white trap caught significantly less sand flies than all of the other traps evaluated (Table 1). With trap preference ignored, significantly more females (14.1 ± 0.9) than males (6.9 ± 0.4) were captured overall. There were significantly more females than males captured by all traps except the all-white trap and the white catch bag/no lid trap (Table 2). The all-white trap and the white catch bag/no lid trap captured significantly fewer females than the four other traps, however, significance groupings for males were less clearly defined (Table 2). The all-black trap captured numerically more male sand flies than all other traps evaluated, and significantly more male sand flies than the black catch bag/no lid and all-white traps.

Experiment II-Trap Orientation

The main effects model was significant for the dependent variable ($F=71.99$, d.f.=19,199, $P<0.0001$). Means for the total numbers of flies captured ranged from 996.3 to 2.7 and overall, traps captured significantly more females than males.

All traps with CO_2 captured significantly more sand flies than the trap configurations not using CO_2 ; however, traps with CO_2 plus black bodies, heat and moisture captured significantly more sand flies than the traps with CO_2 alone (Table 3). When CO_2 is added to the black body, heat and moisture combination, the sand fly catch increases significantly, indicating the importance of CO_2 as an attractant. All added features significantly increased the trap catches when compared with the controls, but there were no significant differences between like traps resulting from trap inversion when sex is overlooked (Table 3). With trap preference ignored, significantly more females (435.4 ± 62.6) than males (61.8 ± 9.2) were captured overall.

With the exception of the control traps, all trap

Table 2. Mean numbers (\pm SE) of female and male *P. papatasi* adults captured on CDC traps modified to have black or white lids and catch bags ($n = 10$).

Trap	Females ¹	Males
Bag Black-Lid Black (BB-LB)	19.4 ± 2.5aX	9.6 ± 1.2bX
Bag White-Lid Black (BW-LB)	17.3 ± 2.1aX	6.9 ± 0.9bXYZ
Bag Black-Lid White (BB-LW)	16.5 ± 2.2aX	6.0 ± 1.0bXYZ
Bag Black-No Lid (BB-NL)	15.7 ± 1.8aX	8.5 ± 1.3aXY
Bag White-No Lid (BW-NL)	9.4 ± 1.1aY	5.6 ± 0.9bYZ
Bag White-Lid White (BW-LW)	6.4 ± 0.8aY	4.5 ± 0.6aZ

¹Means for females and males for like traps in rows (lower case) and for females or males in columns (upper case) followed by the same letter are not significantly different ($P < 0.05$; Ryan-Einot-Gabriel-Welsch Multiple Range Test [SAS Institute 2003]).

configurations captured significantly more females than males (Table 4). Significantly more females were captured by traps with CO₂ than by those without CO₂, and the inversion of traps resulted in significant increases in catches of female sand fly in like traps, notably the traps with CO₂ alone and the traps with black bodies, heat and moisture, but no CO₂ (Table 4). With males, traps with CO₂ and traps with UV light captured significantly more sand flies than the other traps evaluated (Table 4). In fact, there were no

significant differences in males captured between the traps with UV light and those with CO₂ alone. Inversion of traps had no significant effect on male trap catches between like traps (Table 4).

DISCUSSION

The literature is replete with studies indicating that many biting flies are attracted to optical targets (for a review see Allan et al. 1987). In previous studies, *P. papatasi* and *Lutzomyia* spp. sand flies were attracted, if given a choice of colors, mainly to red LEDs (Hoel et al. 2007, Mann et al. 2009). Also, mosquitoes are attracted to different types of LEDs but it appears that colors are species, or at least ecotype, specific (Burkett et al. 1998, Burkett and Butler 2005). Bearing in mind that sand flies have a similar spectral sensitivity as mosquitoes (Muir et al. 1992), it would not be surprising if in the future color preferences for different sand fly species are also found. Because these previously reported studies were conducted with colors of light produced by LEDs and bulbs, while in our study we used colored surfaces, results might vary.

To the best of our knowledge, this is the first study to demonstrate that female sand flies are significantly attracted to black color, a trait which is very wide spread among hematophagous Diptera (Gibson and Torr 1999). However because our traps were not entirely black, this attraction could be due to the contrast between the light and dark surfaces in the traps. An attraction by the females to the traps having contrasting colors rather than to the all-white trap was significant (Table 2), and this could be a general trait among sand flies. Additional knowledge about this factor might lead to improvements in the CDC trap.

Carbon dioxide is a strong long-range attractant for most female biting flies (Gibson and Torr 1999). Therefore it is not surprising that traps with CO₂, regardless of trap

Table 3. Mean numbers (\pm SE) of *P. papatasi* adults captured on CDC traps with selected modifications ($n = 20$).

Trap	Sand flies captured ¹
CNTRL trap black body, heat, moisture and CO ₂ inverted,	996.3 \pm 187.4a
CNTRL trap black body heat moisture, and CO ₂	628.1 \pm 125.6a
CNTRL trap with CO ₂ inverted.	473.4 \pm 108.5b
CNTRL trap with CO ₂	262.4 \pm 61.5b
CNTRL with UV inverted	41.8 \pm 4.0c
CNTRL trap with UV light	36.2 \pm 3.7c
CNTRL trap black body, heat and moisture inverted	26.5 \pm 5.3d
CNTRL trap black body, heat and moisture	15.0 \pm 2.7d
CNTRL trap inverted	3.4 \pm 0.6e
CDC with suction and incandescent light	2.7 \pm 0.5e

¹Means followed by the same letter are not significantly different ($P < 0.05$; Ryan-Einot-Gabriel-Welsch Multiple Range Test [SAS Institute 2003]).

Table 4. Mean numbers (\pm SE) of female and male *P. papatasi* adults captured on CDC traps with selected modifications ($n = 10$).

Trap	Females ¹	Males
CNTRL trap black body, heat, moisture and CO ₂ inverted	1740.8 \pm 153.9aU	251.7 \pm 38.1bW
CNTRL trap black body heat moisture	1086.8 \pm 138.9aUV	169.4 \pm 24.0bW
CNTRL trap with CO ₂ inverted	880.5 \pm 112.9aV	66.3 \pm 12.1bX
CNTRL trap with CO ₂	484.7 \pm 70.2aW	40.1 \pm 6.0bX
CNTRL trap black body, heat and moisture inverted	46.1 \pm 5.6aX	6.9 \pm 1.3bY
CNTRL with UV inverted	45.4 \pm 5.6aX	38.1 \pm 5.7bX
CNTRL trap with UV light	39.4 \pm 5.7aXY	33.0 \pm 4.8bX
CNTRL trap black body, heat and moisture	23.6 \pm 3.5aY	6.4 \pm 1.1bY
CNTRL trap inverted	3.5 \pm 1.0aZ	3.2 \pm 0.7aYZ
CDC with suction and incandescent light	3.0 \pm 0.8aZ	2.4 \pm 0.5aZ

¹Means for females and males for like traps in rows (lower case) and for females or males in columns (upper case) followed by the same letter are not significantly different ($P < 0.05$; Ryan-Einot-Gabriel-Welsch Multiple Range Test [SAS Institute 2003]).

orientation, captured large numbers of sand flies (Table 3). The black trap body-heat-moisture combination provides about a 10-fold increase over the control traps in female sand flies captured (Table 4), but when CO₂ was added to that combination the female captures increased by 38 to 45 times. Similar almost synergistic effects were produced by combining CO₂ with red LEDs and a 1-octen-3-ol/1-hexen-3-ol lure for *Lutzomyia shannoni* (Mann et al. 2009) and combining CO₂ with octenol for *P. papatasi* (Beavers et al. 2004).

The traps with UV light increased the females captured by ca. 12 times over the control traps (Table 4). While this is not competitive with traps using CO₂, Burkett et al. (2007) reported excellent results with a similar UV-CDC device when evaluated against other non-CO₂ traps. Unfortunately, this device captured considerably more *Sergentomyia* spp. than *P. papatasi*.

In a recent study near our experimental area, Faiman et al. (2009) demonstrated that inverted CDC traps, with openings 10 cm above the ground and baited with dry ice caught an average of 1.6 times more female and 1.7 times more male *P. papatasi* as upright CDC traps, baited with dry ice with the trap opening 40 cm above the ground. In our study, we observed a similar effect for *P. papatasi* if traps were baited with anything but suction alone (control)(Table 4). Burkett et al. (2007) also had favorable results with an updraft CDC trap and it captured a higher percentage of *P. papatasi* than the other traps evaluated.

Faiman et al. (2009) speculated that the higher catches with inverted traps might be related to the proximity of the trap openings to the ground and a more dense and more focused CO₂ plume (Cooperband and Carde, 2006). Our trap openings were 50 cm above the ground regardless of trap orientation but in an additional experiment we observed no significant differences between inverted traps with openings 10 cm and 50 cm above the ground (data not shown). This suggests that the results cannot be explained only by the difference in height. Sand flies are supposedly flying only short distances, very low and moving along the ground often in short jumps (Killick-Kendrick et al. 1986, Doha et al. 1991, Alexander and Young 1992). In previous studies with repellents in southern Israel, we observed that most *P. papatasi* bite the lower extremities mainly below the knee (unpublished data of the authors). If female sand flies are approaching potential hosts in an upward movement, they may be more easily caught if the suction is at the bottom of a trap than at the top. This might also explain why significantly more females, but not males, were caught in inverted traps baited with materials characteristic of potential hosts (black color, heat, moisture, CO₂) compared to upright traps baited similarly. Males are attracted to hosts for the opportunity of mating with host-seeking females. They are probably reacting differently from females and are not trying to find a suitable area for blood feeding (Lane et al. 1990, Memmott 1991, 1992) which might explain the smaller catches.

In summary, when exposed to traps with black and white components, adults of *P. papatasi* are more attracted

to darker traps or traps with more contrast. When exposed to traps having some characteristics of a live host, CO₂ plays a strong role in attraction, with smaller degrees of attraction observed from other components. Inversion of traps can result in a 1.5- to 2-fold increase in trap catch.

Acknowledgments

This study was supported in part by funds provided to the U.S. Department of Agriculture from the Department of Defense through the Deployed War-Fighter Protection Program (DWFP).

REFERENCES CITED

- Alexander, B. 2000. Sampling methods for phlebotomine sand flies. *Med. Vet. Entomol.* 14:109-122.
- Alexander, B. and D. Young. 1992. Dispersal of phlebotomine sand flies (Diptera: Psychodidae). *Mem. Inst. Oswaldo Cruz.* 87: 397-403.
- Allan, S.A., J.F. Day, and J.D. Edman. 1987. Visual ecology of biting flies. *Annu. Rev. Entomol.* 32:297-316.
- Allan, S.A., U.R. Bernier, and D.L. Kline. 2006. Laboratory evaluation of avian odors for mosquito (Diptera: Culicidae) attraction. *J. Med. Entomol.* 43: 225-231.
- Andrade, A.J., M.R. Andrade, E.S. Dias, M.C. Pinto, and I.E. Eiras. 2008. Are light traps baited with kairomones effective in the capture of *Lutzomyia longipalpis* and *Lutzomyia intermedia*? An evaluation of synthetic human odor as an attractant for phlebotomine sand flies (Diptera: Psychodidae: Phlebotominae). *Mem. Inst. Oswaldo Cruz.* 103: 337-343.
- Ashford, R.W. 2001. *Phlebotomus* fevers. In: M. W. Service (ed.) *The Encyclopedia of Arthropod-Transmitted Infections*. pp. 397-401. CABI Publishing, Wallingford, U.K.
- Ashbel, D. 1951. *Bio-climatic Atlas of Israel*. Meteorology Dept. of the Hebrew University. Jerusalem. 151pp. (Hebrew and English).
- Beavers, G.M., H.A. Hanafi, and E.A. Dykstra. 2004. Evaluation of 1-octen-3-ol and carbon dioxide as attractants for *Phlebotomus papatasi* (Diptera: Psychodidae) in southern Egypt. *J. Am. Mosq. Contr. Assoc.* 20: 130-133.
- Bernier, U.R., D.L. Kline, K.H. Posey, M. Booth, R.A. Yost, and D.R. Barnard. 2003. Synergistic attraction of *Aedes aegypti* (L.) to binary blends of L-lactic acid and acetone, dichloromethane, or dimethyl disulfide. *J. Med. Entomol.* 40: 653-656.
- Birtles, R.J. 2001. Carrion's disease. In: M. W. Service (ed.) *The Encyclopedia of Arthropod-Transmitted Infections*. 104-106. CABI Publishing, Wallingford, U.K.
- Burkett, D.A., J.F. Butler, and D.L. Kline. 1998. Field evaluation of colored light-emitting diodes as attractants for woodland mosquitoes and other Diptera in north central Florida. *J. Am. Mosq. Contr. Assoc.* 14: 186-195.
- Burkett, D.A. and J.F. Butler. 2005. Laboratory evaluation of

- colored light as an attractant for female *Aedes aegypti*, *Aedes albopictus*, *Anopheles quadrimaculatus* and *Culex nigripalpus*. Fla. Entomol. 88: 383-389.
- Burkett, D.A., R. Knight, J.A. Dennett, V. Sherwood, E. Rowton, and R.E. Coleman. 2007. Impact of phlebotomine sand flies on U.S. military operations at Tallil Air Base, Iraq: 3. Evaluation of surveillance devices for the collection of adult sand flies. J. Med. Entomol. 44: 381-384.
- Comer, J.A. and R.B. Tesh. 1991. Phlebotomine sand flies as vectors of vesiculovirus: a review. Parasitologia. 33: 143-150.
- Cooperband, M.F. and R.T. Carde. 2006. Orientation of *Culex* mosquitoes to carbon dioxide-baited traps: flight manoeuvres and trapping efficiency. Med. Vet. Entomol. 20: 11-16.
- Danin, A. 1988. Flora and vegetation of Israel and adjacent areas. In: Yom-Tov & Tchernov (eds.). *The Zoogeography of Israel*. pp. 251-276. Junk Publishers, Dordrecht, Netherlands.
- Doha, S., M.G. Shehata, S.M. El Said, and B. El Sawaf. 1991. Dispersal of *Phlebotomus papatasi* (Scopoli) and *P. langeroni* Nitzulescu in El Hammam, Matrouh Governorate, Egypt. Ann. Parasitol. Hum. Comp. 66: 69-76.
- Dougherty, M.J., P.M. Guerin, R.D. Ward, and J.G.C. Hamilton. 1999. Behavioral and electrophysiological responses of the phlebotomine sand fly *Lutzomyia longipalpis* (Diptera: Psychodidae) when exposed to canid host odor kairmones. Phys. Entomol. 24: 251-262.
- Faiman, R., R. Cuño, and A. Warburg. 2009. Comparative efficacy of three suction traps for collecting phlebotomine sand flies (Diptera: Psychodidae) in open habitats. J. Vector Ecol. 34: 114-118.
- Feliciangeli, M.D. 2004. Natural breeding places of phlebotomine sandflies. Med. Vet. Entomol. 18: 71-80.
- Gibson, G. and S.J. Torr. 1999. Visual and olfactory response of haematophagous Diptera to host stimuli. Med. Vet. Entomol. 13: 2-23.
- Hoel, D.F., J.F. Butler, E.Y. Fawaz, N. Watany, S.S. El-Hossary, and J. Villinski. 2007. Response of phlebotomine sand flies to light-emitting diode-modified light traps in southern Egypt. J. Vector Ecol. 32: 302-308.
- Killick-Kendrick, R. 1999. The biology and control of phlebotomine sand flies. Clinics Dermatol. 17: 279-289.
- Killick-Kendrick, R., T.J. Wilkes, M. Bailly, I. Bailly, and L.A. Righton. 1986. Preliminary field observations on the flight speed of a phlebotomine sand fly. Trans. R. Trop. Med. Hyg. 80: 138-142.
- Kline, D.L. 1998. Olfactory responses and field attraction of mosquitoes to volatiles from Limburger cheese and human foot odor. J. Vector Ecol. 23: 186-194.
- Kline, D.L. 2006. Traps and trapping techniques for adult mosquito control. J. Am. Mosq. Contr. Assoc. 22: 490-496.
- Lane, R.P. 1993. Sandflies (Phlebotominae). In: R.P. Lane and R.W. Crosskey (eds.) *Medical Insects and Arachnids*. Chapman & Hall, London. 78-119.
- Lane, R.P., M.M. Pile, and F.P. Amerasinghe. 1990. Anthropophagy and aggregation behavior of the sandfly *Phlebotomus argentipes* in Sri Lanka. Med. Vet. Entomol. 4:79-88.
- Mann, R.S., P.E. Kaufman, and J.F. Butler. 2009. *Lutzomyia* spp. (Diptera: Psychodidae) response to olfactory attractant and light emitting diode modified Mosquito Magnet X (MM-X) traps. J. Med. Entomol. 46: 1052-1061.
- Mellor, H.E., J.G.C. Hamilton, and M. Anderson. 1996. Spectral sensitivity in the eyes of male and female *Lutzomyia longipalpis* sand flies. Med. Vet. Entomol. 10: 371-374.
- Memmott, J. 1991. Sand fly distribution and abundance in a tropical rain forest. Med. Vet. Entomol. 5: 403-411.
- Memmott, J. 1992. Patterns of sand fly distribution in a tropical forest: a causal hypothesis. Med. Vet. Entomol. 6: 188-194.
- Morton, I.E. and R.D. Ward. 1989. Laboratory response of female sand flies (*Lutzomyia longipalpis*) to a host and male pheromone source over distance. Med. Vet. Entomol. 3: 219-223.
- Muir, L.E., B.H. Kay, and M.J. Thorne. 1992. *Aedes aegypti* (Diptera: Culicidae) vision: response to stimuli from the optical environment. J. Med. Entomol. 29: 445-450.
- Müller, G.C. and Y. Schlein. 2004. Nectar and honeydew feeding of *Phlebotomus papatasi* in a focus of *Leishmania major* in Neot Hakikar oasis. J. Vector Ecol. 29: 154-158.
- Müller G.C. and Y. Schlein. 2006. Sugar questing mosquitoes in arid areas gather on scarce blossoms that can be used for control. Intl. J. Parasitol. 36: 1077-1080.
- Murphy, M.W., R.F. Dunton, M.J. Perich, and W.A. Rowley. 2001. Attraction of *Anopheles* (Diptera: culicidae) to volatile chemicals in Western Kenya. J. Med. Entomol. 38: 242-244.
- Nigam, Y. and R.D. Ward. 1991. Male sand fly pheromone and artificial host as attractants for female *Lutzomyia longipalpis* (Diptera: Psychodidae). Phys. Entomol. 16: 305-312.
- Nowinszky, L. 2004. Nocturnal illumination and night flying insects. Appl. Ecol. Environ. Res. 2: 17-52.
- Pinto, M.C., D.H. Campbell-Lendrum, and A.L. Lozovei. 2001. Phlebotomine sandfly responses to carbon dioxide and human odour in the field. Med. Vet. Entomol. 15: 132-139.
- SAS Institute. 2003. SAS/STAT User's Guide. Version 9.1, Cary, NC: SAS Institute.
- Schlein, Y. and A. Warburg. 1986. Phytophagy and the feeding cycle of *Phlebotomus papatasi* (Diptera: Psychodidae) under experimental conditions. J. Med. Entomol. 23:11-15.
- Schlein Y., R.L. Jacobson, and G.C. Müller. 2001. Sand fly feeding on noxious plants: a potential method for the control of leishmaniasis. Am. J. Trop. Med. Hyg. 65: 300-303.

- Service, M.W. 1993. *Mosquito Ecology: Field Sampling Methods*. 2nd ed. London Chapman & Hall. 583pp.
- Sharp, B. L., F.G. Quicke, and E.J. Jansen 1984. Aspects of the behavior of five anopheline species in the endemic malaria area of Natal. *J. Entomol. Soc. S. Afr.* 47: 251-258.
- Young, D.G. and Perkins, P.V. 1984. Phlebotomine sand flies of North America (Diptera: Psychodidae). *J. Am. Mosq. Contr. Assoc.* 44: 263-304.