

2017

Key to the Korean species of armored scale insects (Hemiptera: Diaspididae)

Soo-Jung Suh

Plant Quarantine Technology Center/QIA

Follow this and additional works at: <http://digitalcommons.unl.edu/insectamundi>



Part of the [Ecology and Evolutionary Biology Commons](#), and the [Entomology Commons](#)

Suh, Soo-Jung, "Key to the Korean species of armored scale insects (Hemiptera: Diaspididae)" (2017). *Insecta Mundi*. 1031.
<http://digitalcommons.unl.edu/insectamundi/1031>

This Article is brought to you for free and open access by the Center for Systematic Entomology, Gainesville, Florida at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Insecta Mundi by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

INSECTA MUNDI

A Journal of World Insect Systematics

0526

Key to the Korean species of armored scale insects
(Hemiptera: Diaspididae)

Soo-Jung Suh
Plant Quarantine Technology Center/QIA
167, Yongjeon 1-ro, Gimcheon-si,
Gyeongsangbuk-do, South Korea 39660

Date of Issue: February 24, 2017

Soo-Jung Suh

Key to the Korean species of armored scale insects (Hemiptera: Diaspididae)

Insecta Mundi 0526: 1–11

ZooBank Registered: urn:lsid:zoobank.org:pub:42A72836-95E0-4805-802A-30B7CE6EBE81

Published in 2017 by

Center for Systematic Entomology, Inc.

P. O. Box 141874

Gainesville, FL 32614-1874 USA

<http://centerforsystematicentomology.org/>

Insecta Mundi is a journal primarily devoted to insect systematics, but articles can be published on any non-marine arthropod. Topics considered for publication include systematics, taxonomy, nomenclature, checklists, faunal works, and natural history. **Insecta Mundi** will not consider works in the applied sciences (i.e. medical entomology, pest control research, etc.), and no longer publishes book reviews or editorials. **Insecta Mundi** publishes original research or discoveries in an inexpensive and timely manner, distributing them free via open access on the internet on the date of publication.

Insecta Mundi is referenced or abstracted by several sources including the Zoological Record, CAB Abstracts, etc. **Insecta Mundi** is published irregularly throughout the year, with completed manuscripts assigned an individual number. Manuscripts must be peer reviewed prior to submission, after which they are reviewed by the editorial board to ensure quality. One author of each submitted manuscript must be a current member of the Center for Systematic Entomology.

Chief Editor: David Plotkin, e-mail: insectamundi@gmail.com

Assistant Editor: Paul E. Skelley, e-mail: insectamundi@gmail.com

Head Layout Editor: Eugenio H. Nearn

Editorial Board: J. H. Frank, M. J. Paulsen, Michael C. Thomas

Review Editors: Listed on the **Insecta Mundi** webpage

Manuscript Preparation Guidelines and **Submission Requirements** available on the **Insecta Mundi** webpage at: <http://centerforsystematicentomology.org/insectamundi/>

Printed copies (ISSN 0749-6737) annually deposited in libraries:

CSIRO, Canberra, ACT, Australia

Museu de Zoologia, São Paulo, Brazil

Agriculture and Agrifood Canada, Ottawa, ON, Canada

The Natural History Museum, London, UK

Muzeum i Instytut Zoologii PAN, Warsaw, Poland

National Taiwan University, Taipei, Taiwan

California Academy of Sciences, San Francisco, CA, USA

Florida Department of Agriculture and Consumer Services, Gainesville, FL, USA

Field Museum of Natural History, Chicago, IL, USA

National Museum of Natural History, Smithsonian Institution, Washington, DC, USA

Zoological Institute of Russian Academy of Sciences, Saint-Petersburg, Russia

Electronic copies (Online ISSN 1942-1354, CDROM ISSN 1942-1362) in PDF format:

Printed CD or DVD mailed to all members at end of year. Archived digitally by Portico.

Florida Virtual Campus: <http://purl.fcla.edu/fcla/insectamundi>

University of Nebraska-Lincoln, Digital Commons: <http://digitalcommons.unl.edu/insectamundi/>

Goethe-Universität, Frankfurt am Main: <http://nbn-resolving.de/urn/resolver.pl?urn:nbn:de:hebis:30:3-135240>

Copyright held by the author(s). This is an open access article distributed under the terms of the Creative Commons, Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited. <http://creativecommons.org/licenses/by-nc/3.0/>

Layout Editor for this article: Eugenio H. Nearn

Key to the Korean species of armored scale insects (Hemiptera: Diaspididae)

Soo-Jung Suh

Plant Quarantine Technology Center/QIA
167, Yongjeon 1-ro, Gimcheon-si,
Gyeongsangbuk-do, South Korea 39660
suhsj97@gmail.com; suhsj97@korea.kr

Abstract. A list of all 80 species of armored scale insects (Hemiptera: Diaspididae) in 30 genera that have been identified or documented in Korea, along with a dichotomous key to separate them, is provided.

Key words. Diaspidids, checklist, identification, indigenous species

Introduction

Armored scale insects (Hemiptera: Diaspididae) are the largest and most diverse family of scale insects, with 2,587 species known worldwide (García et al. 2016). They are highly evolved and specialized plant parasites, and generally have a cryptic way of life because of their small size and limited mobility. The female Diaspididae has two nymphal stages; first-instar nymphs (crawlers) have well-developed legs and are known as the dispersal stage; second-instar nymphs and adult females (third-instar) are legless, wingless, feed on sap, and are covered by a waxy shield incorporating the exuviae of previous instars; adult males (when present) have legs and one pair of wings and do not feed; many reproduce asexually and have no males (Rosen 1990).

Armored scale insects occur in every part of the world where plants grow. There is hardly any country in the world today where diaspidids do not cause serious pest problems, and their success invading new territories has made them a constant quarantine threat in many parts of the world (Rosen 1990). They are one of the most economically important groups of insects found on food crops and ornamental plants.

In the Republic of Korea, armored scale insects such as the Asiatic rose scale, *Aulacaspis rosarum* Borchsenius and the euonymus scale, *Unaspis euonymi* (Comstock), are considered major pests because they cause severe damage to trees and shrubs due to their high populations. In addition, more and more non-indigenous armored scale insects such as *Aulacaspis yasumatsui* Takagi (intercepted 29 times) and *Lepidosaphes laterochitinsa* Green (intercepted 65 times) are frequently intercepted on imported plants over the past 20 years and are considered to be potential invasive species (Suh 2016c).

Due to their importance, a survey was conducted between 2006 and 2015 to regularly update a list of armored scale insects found in the Republic of Korea. Recent articles (Paik 2000; Suh and Hodges 2007; Lee 2010; Suh 2011, 2012, 2013, 2014, 2015, 2016a, 2016b; Yu and Suh 2013; Suh and Evans 2016) in addition to information reported herein, have increased the number of armored scale species known to occur in the Republic of Korea to 80 species belonging to 30 genera.

The purpose of this paper is to provide an identification key to the genera and species of armored scale insects known to occur in Korea, based on characters of the adult female. This information will not only enable researchers to identify the species known to occur in Korea, but also aid in the recognition and early detection of newly introduced species.

Materials and Methods

This paper provides a dichotomous key and a list concerning 80 species of armored scale insects reported from Korea. But during the survey over the past ten years, nine species such as *Chionaspis salicis* (Linnaeus) and *Comstockaspis macroporana* (Takagi) were not collected (Table 1), although they have been documented in the Korean fauna of armored scale insects. While developing the key, the author was not able to examine specimens of these species from Korea. Terminology for the morphological structures used in the key follows that of Miller and Davidson (2005).

Results

Table 1 presents a list of the armored scale insects that were collected and reported from the Republic of Korea, along with their regional distribution in Korea. A dichotomous key to the Korean armored scale insects is provided.

Key to armored scale insects of Korea (slide-mounted adult females)

1.	Dorsum of pygidium with areolate pattern	2
—	Dorsum of pygidium without areolate pattern	3
2(1).	With perispiracular pores near posterior spiracles; pygidial macroducts short, approximately 5 times longer than width of dermal orifice	<i>Pseudaonidia paeoniae</i> (Cockerell)
—	Without perispiracular pores near posterior spiracles; pygidial macroducts long, approximately 10 times longer than width of dermal orifice	<i>Pseudaonidia duplex</i> (Cockerell)
3(1).	Usually occurring on bamboos	4
—	Rarely occurring on bamboos	9
4(3).	Median lobes fused; lacking two-barred macroducts	<i>Odonaspis secreta</i> (Cockerell) (in part)
—	Median lobes not fused; two-barred macroducts present	5
5(4).	Median lobes yoked basally, with small space between them; medial lobule of second lobe larger than lateral lobule	<i>Pinnaspis buxi</i> (Bouché) (in part)
—	Median lobes not basally yoked, separated by space at least a half width of lobe or far apart from each other; medial lobule of second lobe about same size as, or smaller than 1 lateral lobule	6
6(5).	Apically toothed marginal plates absent anterior to 2nd lobes; median to 3rd lobes very small and pointed	<i>Unachionaspis tenuis</i> (Maskell)
—	Apically toothed marginal plates present anterior to 2nd lobes; median, second and third lobes without the above combination of features	7
7(6).	With less than 10 perivulvar pores on each side of body; occurring on the leaves	<i>Kuwanaspis hikosani</i> (Kuwana)
—	With more than 20 perivulvar pores on each side of body; occurring on stems	8
8(7).	Abdominal segment 1 with transverse row of macroducts ...	<i>Kuwanaspis howardi</i> (Cooley)
—	Abdominal segment 1 without transverse row of macroducts	<i>Kuwanaspis pseudoleucaspis</i> (Kuwana)
9(3).	With at least one pore near the posterior or anterior spiracles	10
—	Without pores near the spiracles	63
10(9).	Body elongate, oval, or turbinate; not as described below	16
—	Body elongate; head and/or anterior two thoracic segments rectangular in shape, wider than remainder of body	11
11(10).	With ducts on dorsum of abdominal segment 2	12
—	Without ducts on dorsum of abdominal segment 2	14

- 25(19). With at least a few macroducts on medial or submarginal areas of pygidium; not pupillarial (adult female developing with second-instar nymph exuvium) **26**
 — Without macroducts on medial or submarginal areas of pygidium; pupillarial ***Fiorinia japonica* Kuwana**
- 26(25). Setae or gland spines absent between the median lobes **27**
 — Setae or gland spines present between the median lobes **31**
- 27(26). With a series of small dorsal macroducts (wider than microducts in gland spines) on the submedial area of the abdomen **28**
 — Without a series of small dorsal macroducts on the submedial area of the abdomen **30**
- 28(27). Median lobes strongly divergent, with a straight medial margin; with small dorsal macroducts on the submedial area of abdominal segments 1 to 5 **29**
 — Median lobes only slightly divergent, with a rounded medial margin; with small dorsal macroducts on the submedial area of abdominal segments 1 to 4; on many hosts including *Salix* and *Populus* (Salicaceae) ***Chionaspis salicis* (Linnaeus)**
- 29(28). With small dorsal macroducts on the submedial area of the prothorax and mesothorax; primarily on *Alnus* (Betulaceae) ***Chionaspis alnus* Kuwana**
 — Without small dorsal macroducts on the submedial area of the prothorax and mesothorax; on *Quercus* (Fagaceae) ***Chionaspis saitamaensis* Kuwana**
- 30(27). With dorsal macroducts on abdominal segment 6; primarily on *Wisteria* (Fagaceae) ***Chionaspis wistariae* Cooley**
 — Without dorsal macroducts on abdominal segment 6; on *Acer* (Sapindaceae) ***Chionaspis acer* (Takagi and Kawai)**
- 31(26). Body oval or turbinate, length usually less than 1.5 times greatest width **32**
 — Body elongate, length usually 1.5 times or more than greatest width **33**
- 32(31). Third space usually with 1 gland spine; at least 1 bifurcate or trifurcate gland spine in second, third, or fourth space; antennae with sclerotized projections apically; with 5–22 (10) small macroducts on each side of the metathorax and segment 1; on many hosts ***Pseudaulacaspis pentagona* (Targioni-Tozzetti)**
 — Third space usually with 2 or more gland spines; simple gland spine in second, third, or fourth space; antennae rounded apically; with 0–15 (5) small macroducts on each side of the metathorax and segment 1; primarily on *Prunus* (Rosaceae) ***Pseudaulacaspis prunicola* (Maskell)**
- 33(31). With perispiracular pores near posterior spiracles **34**
 — Without perispiracular pores near posterior spiracles **35**
- 34(33). With usually two rows of submarginal macroducts on abdominal segments 2 and 3; with 4–11 pores near anterior spiracles and 1–5 pores near posterior spiracles; found on *Zelkova* and *Ulmus* (Ulmaceae) ***Pseudaulacaspis ulmicola* Tang**
 — With one row of submarginal macroducts on abdominal segments 2 and 3; with 13–27 pores near anterior spiracles and 6–10 pores near posterior spiracles; only found on *Carpinus* (Betulaceae) ***Pseudaulacaspis latiloba* (Takagi and Kawai)**
- 35(33). Without a duct on abdominal segment 6; antennae usually widely apart, width between antennae wider than distance between median lobes; only found on *Quercus* (Fagaceae) ***Pseudaulacaspis kiushiuensis* (Kuwana)**

- With duct on abdominal segment 6; antennae usually close together, width between antennae narrower than distance between median lobes; found on many hosts *Pseudaulacaspis cockerelli* (Cooley)
- 36(18). Median lobes round, with medial margin about the same length as lateral margin 39
- Median lobes with lateral margin conspicuously longer than medial margin 37
- 37(36). Pygidium with transverse sclerosis at the base of median lobe; bosses absent from abdomen 38
- Pygidium without transverse sclerosis at the base of median lobe; bosses present on abdomen *Andaspis recurrens* Takagi and Kawai
- 38(37). Pygidial marginal macroducts numbering 5 on each side; without submedian dorsal ducts on the pygidium *Andaspis crawii* Cockerell
- Pygidial marginal macroducts numbering 6 on each side; with submedian dorsal ducts on the pygidium *Andaspis kashicola* (Takahashi)
- 39(36). Second lobes bilobate with two lobules 40
- Second lobes simple with one lobule 59
- 40(39). Body elongate, body length more than two times maximum width 41
- Body turbinate, body length less than two times maximum width 56
- 41(40). Pygidium with three pairs of lobes; third lobes distinct 42
- Pygidium with two pairs of lobes; if third lobes present, then small and pointed 44
- 42(41). Macroducts present in submarginal and medial areas of abdominal segment 7; third lobes bilobate; not found on grasses 43
- Macroducts absent from submarginal and medial areas of abdominal segment 7; third lobes simple; only found on grasses *Duplachionaspis divergens* (Green)
- 43(42). Perivulvar pores absent; second lobes usually protruding beyond the median lobes; head, thorax and first abdominal segment sclerotized in mature females *Unaspis yanonensis* (Kuwana)
- Perivulvar pores present; second lobes not protruding beyond the median lobes; head, thorax and first abdominal segment not sclerotized *Unaspis euonymi* (Comstock)
- 44(41). Usually occurring on conifers 45
- Rarely occurring on conifers 48
- 45(44). Perivulvar pores in 8 groups, 3 auxiliary groups being present cephalad of usual 5 groups *Lepidosaphes pini* (Maskell)
- Perivulvar pores in 4-5 groups 46
- 46(45). With minute spinules on head *Lepidosaphes okitsuensis* Kuwana
- Without minute spinules on head 47
- 47(46). With lateral spurs between abdominal segments 1 and 2 *Lepidosaphes japonica* (Kuwana)
- Without lateral spurs between abdominal segments 1 and 2 *Lepidosaphes pallida* (Maskell)

48(44). With lateral spurs or tubercles on metathorax and/or on abdominal segments	49
— Without lateral spurs or tubercles	54
49(48). Eye spur like	<i>Lepidosaphes pinnaeformis</i> (Bouché)
— Eye not spur like	50
50(49). With bosses or cicatrices on dorsal submargin of one or more abdominal segments	<i>Lepidosaphes ulmi</i> (Linnaeus)
— Without bosses or cicatrices	51
51(50). With minute spinules on head	52
— Without minute spinules on head	53
52(51). Without lateral spurs on metathorax	<i>Lepidosaphes tubulorum</i> Ferris
— With lateral spurs on metathorax	<i>Lepidosaphes ussuriensis</i> (Borchsenius)
53(51). Cephalothorax elongate, little narrowing anteriorly; thoracic region and first abdominal segment becoming heavily sclerotized dorsally at full maturity ..	<i>Lepidosaphes gloverii</i> (Packard)
— Cephalothorax gradually narrowing anteriorly; thoracic region remaining membranous, but abdominal segments becoming sclerotized dorsally at full maturity	<i>Lepidosaphes yanagicola</i> Kuwana
54(48). Dorsal macroducts present in a row across median area on abdominal segments 3 and 4; 40 or fewer perivulvar pores	55
— Dorsal macroducts absent from median area on abdominal segments 3 and 4; 87-140 perivulvar pores	<i>Lepidosaphes towadensis</i> Takagi and Kawai
55(54). Pygidium broadly rounded; median lobes separated by a space narrower than width of one median lobe; on many hosts including <i>Malus</i> , <i>Prunus</i> , and <i>Pyrus</i> (Rosaceae)	<i>Lepidosaphes conchiformis</i> (Gmelin)
— Pygidium more or less trapezoidal; median lobes separated by a space as wide as one median lobe; primarily on <i>Camellia</i> (Theaceae)	<i>Lepidosaphes kamakurensis</i> (Kuwana)
56(40). Submarginal areas of posterior pygidial segments with at least 2 barrel-shaped macroducts ..	57
— Submarginal areas of posterior pygidial segments without macroducts	<i>Carulaspis juniperi</i> (Bouché)
57(56). Submedial macroducts present on abdominal segments 2 to 5; without notches on medial margin of the median lobes; primarily found on cactus	<i>Diaspis echinocacti</i> (Bouché)
— Submedial macroducts absent from abdominal segments 2 to 5; with notches on medial margin of the median lobes; rarely found on cactus	58
58(57). Submarginal macroducts numbering only two on each side; anterior spiracles with approximately three associated perispiracular pores	<i>Diaspis boisduvalii</i> Signoret
— Submarginal macroducts numbering more than two on each side; anterior spiracles with approximately 14 associated perispiracular pores	<i>Diaspis bromeliae</i> (Kerner)
59(39). Dermal pockets absent between posterior spiracle and body margin	60
— Dermal pockets present between posterior spiracle and body margin	61
60(59). Pygidial lobes conical projections (v-shape); pygidial plates exceeding the lobes; pupillarial ...	<i>Cryptoparlatores leucaspis</i> Lindinger

- Pygidial lobes U-shape; pygidial plates not exceeding the lobes; not pupillarial *Parlatoria pergandii* Comstock
- 61(59). Eye spur-like, apically pointed *Parlatoria proteus* (Curtis)
- Eye variable, usually low dome-shaped 62
- 62(61). With about 60 perivulvar pores; most plates in spaces between first 3 lobes with 2 microducts *Parlatoria theae* Cockerell
- With about 25 perivulvar pores; most plates in spaces between first 3 lobes with 1 microduct *Parlatoria camelliae* Comstock
- 63(9). Median lobes with basal sclerosis 79
- Median lobes without basal sclerosis 64
- 64(63). With perivulvar pores 65
- Without perivulvar pores 75
- 65(64). Paraphyses conspicuous, most as long as or longer than the length of the median lobes 66
- Paraphyses absent or inconspicuous; all paraphyses shorter than median lobes 69
- 66(65). Paraphyses \cap -shape between median lobes, between median and second lobes, and between the second and third lobes *Hypaspidiotus jordani* (Kuwana)
- Paraphyses straight-shape between median lobes, between median and second lobes, and between the second and third lobes 67
- 67(66). With at least 1 cluster of macroducts on submarginal areas of prepygidial segments 68
- Without a cluster of macroducts on submarginal areas of prepygidial segments *Chrysomphalus dictyospermi* (Morgan)
- 68(67). With a cluster of macroducts on submarginal areas of abdominal segment 2 *Chrysomphalus aonidum* (Linnaeus)
- With a cluster of macroducts on submarginal areas of abdominal segments 2 and 3 *Chrysomphalus bifasciculatus* Ferris
- 69(65). Occurring on conifers 70
- Rarely occurring on conifers 72
- 70(69). Second lobes about same size as, or slightly smaller than median lobes, normally same shape as median lobes; anal opening same or narrower than median lobes 71
- Second lobes conspicuously smaller than median lobes, different shape from median lobes; anal opening wider than median lobes *Hemiberlesia lataniae* (Signoret) (in part)
- 71(70). Perivulvar pores present in five groups; median lobes and second lobes broadly rounded; with submarginal macroduct on abdominal segment 1 *Dynaspidiotus tsugae* (Marlatt)
- Perivulvar pores present in four groups; median lobes and second lobes narrowly rounded; without submarginal macroduct on abdominal segment 1 *Dynaspidiotus pseudomeyeri* (Kuwana)
- 72(69). Anal opening large (wider and longer than median lobes) and separated from bases of median lobes by not more than 2 times than its longitudinal diameter *Hemiberlesia lataniae* (Signoret) (in part)
- Anal opening small (narrower and shorter than median lobes) and separated from bases of median lobes by more than 2 times than its longitudinal diameter 73

- 73(72). Third lobes with well-developed sclerotized point *Selenomphalus distylii* Takagi
 — Third lobes simple without a sclerotized point 74
- 74(73). Median lobes closely appressed by less than one-fifth of their width; on *Castanea* and *Quercus* (Fagaceae) *Diaspidiotus cryptoxanthus* (Cockerell)
 — Median lobes separated by distinct space by more than one-half of their width; primarily on *Camellia* (Theaceae) *Diaspidiotus degeneratus* (Leonardi)
- 75(64). Second and third lobes about same size as median lobes 76
 — Second and/or third lobes conspicuously smaller than median lobes 77
- 76(75). Scleroses associated with apophysis anterolaterad of the vulva present on venter of pygidium *Aonidiella citrina* (Coquillett)
 — Scleroses associated with apophysis anterolaterad of the vulva absent from venter of pygidium *Aonidiella taxus* Leonardi
- 77(75). Median lobes with lateral and medial notches; simple shaped plates anterior of seta marking abdominal segment 6; usually occurring on *Pinus* (Pinaceae) *Diaspidiotus makii* (Kuwana)
 — Median lobes with usually lateral notch; bifurcate shaped plates anterior of seta marking abdominal segment 6; occurring on various hosts 78
- 78(77). Anal opening large (wider than median lobes); with marginal macroducts on abdominal segment 3 *Comstockaspis macroporana* (Takagi)
 — Anal opening small (narrower than median lobes); without marginal macroducts on abdominal segment 3 *Comstockaspis perniciosa* (Comstock)
- 79(63). Plates lateral to third lobe dentate, edges sclerotized, with fleshy processes; dorsal marginal setae on the second and third lobes lanceolate (thick and dorsoventrally flattened) *Octaspidiotus stauntoniae* (Takahashi)
 — Plates lateral to third lobe not dentate or sclerotized; dorsal marginal setae on the second and third lobes slender to thick, not flattened 80
- 80(81). Pygidial macroducts long (1.5 times distance between posterior apex of anal opening and base of median lobes), normal; usually occurring on leaves 81
 — Pygidial macroducts short (less than 0.5 times distance between posterior apex of anal opening and base of median lobes), minute; usually occurring on roots 82
- 81(80). Second lobes normally protruding beyond median lobes *Aspidiotus destructor* Signoret
 — Second lobes not protruding beyond median lobes *Aspidiotus cryptomeriae* Kuwana
- 82(80). With perivulvar pores *Rhizaspidiotus canariensis* (Lindinger)
 — Without perivulvar pores *Aspidiella phragmitis* (Takahashi)

Discussion

The purpose of the present work is to provide a checklist and a key for the identification of Korean armored scale insects. This information will aid in the correct identification of each species which is essential for assessing the threat that each species poses and the management strategy that is to be taken. Armored scale insects have received much attention throughout the world due to the damage caused in economically important food crops and ornamental plants and also to the strong possibility of becoming invasive species with international trades for plants.

Of the nine scale species known to occur only in greenhouses, five species were just found on potted seedlings imported into Korea during the survey. They are considered to have failed to establish in the exterior environments. In addition this work provides correct information on the current status of nine greenhouse diaspidid species.

To date, eighty species of armored scale insects in 30 genera have been recorded from the Republic of Korea. However, it can be expected that more species of armored scale insects are likely to be found in Korea if taxonomic studies continue to be performed.

Acknowledgments

I am grateful to Drs. Gregory A. Evans (USDA/APHIS/NIS, Washington, DC, USA) and Takumasa Kondo (Corporación Colombiana de Investigación Agropecuaria (Corpoica), Centro de Investigación Palmira, Palmira, Valle, Colombia) for reviewing the manuscript. Funding for this project came from QIA.

Literature Cited

- García, M., B. Denno, D. R. Miller, G. L. Miller, Y. Ben-Dov, and N. B. Hardy. 2016.** ScaleNet: A literature-based model of scale insect biology and systematics. (Available at ~ <http://scalenet.info>. Last accessed November 2016.)
- Lee, Y. J. 2010.** Family Diaspididae. p. 80-81. *In*: M.K. Paek (ed.). Checklist of Korean insects. Nature and Ecology; Seoul. 598 p.
- Miller, D. R., and J. A. Davidson. 2005.** Armored scale insect pests of trees and shrubs (Hemiptera: Diaspididae). Cornell University Press; Ithaca, NY. 442 p.
- Paik, J. C. 2000.** Economic Insects of Korea 6, Homoptera (Coccinea), Insecta Koreana Suppl. 13. National Institute of Agricultural Science and Technology; Seoul. 193 p.
- Rosen, D. 1990.** Armored Scale Insects, Their Biology, Natural Enemies and Control (series title: World Crop Pests, Vol. 4A). Elsevier Amsterdam; the Netherlands. 688 p.
- Suh, S. J. 2011.** Two new records of *Andaspis* armored scales (Hemiptera: Diaspididae) from Korea. Korean Journal of Applied Entomology 50: 75–77.
- Suh, S. J. 2012.** Notes on pupillarial species of armored scale insects from Korea (Hemiptera: Diaspididae). Korean Journal of Applied Entomology 51: 73–77.
- Suh, S. J. 2013.** On the armored scales, genus *Aulacaspis* Cockerell (Hemiptera: iaspidae), of Korea. Insect Mundi 0295: 1–8.
- Suh, S. J. 2014.** Updated list of *Pinnaspis* armored scales (Hemiptera: Diaspididae) of Korea. Insect Mundi 0347: 1–9.
- Suh, S. J. 2015.** New records of armored scale insects, *Pseudaulacaspis* MacGillivray (Hemiptera: Diaspididae), in Korea. Insect Mundi 0403: 1–8.
- Suh, S. J. 2016a.** New records of *Chionaspis* armored scales (Hemiptera: Diaspididae) in Korea. Insect Mundi 0489: 1–5.
- Suh, S. J. 2016b.** Armored scales (Hemiptera: Diaspididae) from warm temperate forests in Korea. Insect Mundi 0500: 1–6.
- Suh, S. J. 2016c.** Armoured scale insects (Hemiptera: Diaspididae) intercepted at the ports of entry in the Republic of Korea over the last 20 years. Bulletin OEPP/EPPO Bulletin 46: 313–331.
- Suh, S. J., and G. A. Evans. 2016.** A new record of the armored scale genus *Carulaspis* MacGillivray (Hemiptera: Diaspididae) from Korea and its aphelinid parasitoids (Hymenoptera: Aphelinidae). Insect Mundi 0466: 1–6.
- Suh, S. J., and G. S. Hodges. 2007.** Identification of armored scales (Hemiptera: Diaspididae) on bamboos in Korea. Journal of Asia-Pacific Entomology 10: 1–3.
- Yu, H. M., and S. J. Suh. 2012.** A new record for the genus *Duplachionaspis* armored scale (Hemiptera: Diaspididae) from Korea. Korean Journal of Applied Entomology 52: 63–67.

Received January 1, 2017; Accepted January 18, 2017.

Review Editor Marcus Guidoti.

Table 1. Checklist of taxa treated in the identification key with distribution data in Korea (Codes for collecting regions in the Republic of Korea are as follows: GG, Gyeonggido; GW, Gangwondo; CB, Chungcheongbukdo; CN, Chungcheongnamdo; GB, Gyengsangbukdo; GN, Gyengsangnamdo; JB, Jeollabukdo; JN, Jeollanamdo; JJ, Jeju. The nomenclature used here for the Diaspididae follows that of the scale insect database ScaleNet (García et al. 2016). The column headed 'NC' indicates if the species was collected in the Republic of Korea during the survey (2006 to 2015). The asterisk mark (*) indicates 'unknown' (specimens not found outside of greenhouses in the Republic of Korea).

Scientific Name	Distribution within Korea									
	GG	GW	CB	CN	GB	GN	JB	JN	JJ	NC
<i>Andaspis crawii</i> (Cockerell)									•	•
<i>Andaspis kashicola</i> (Takahashi)					•					•
<i>Andaspis recurrens</i> Takagi and Kawai									•	
<i>Aonidiella citrina</i> (Coquillett)										•
<i>Aonidiella taxus</i> Leonardi						•				•
<i>Aspidiella phragmitis</i> (Takahashi)										•
<i>Aspidiotus cryptomeriae</i> Kuwana		•			•	•		•	•	
<i>Aspidiotus destructor</i> Signoret						•				•
<i>Aulacaspis difficilis</i> (Cockerell)						•		•		
<i>Aulacaspis latissima</i> (Cockerell)						•				•
<i>Aulacaspis rosae</i> (Bouché)	•	•	•			•				•
<i>Aulacaspis rosarum</i> Borchsenius	•				•	•	•	•		
<i>Aulacaspis spinosa</i> (Maskell)	•			•	•	•		•	•	
<i>Aulacaspis yabumikiei</i> Kuwana						•		•	•	
<i>Carulaspis juniperi</i> (Bouché)				•						
<i>Chionaspis acer</i> (Takagi and Kawai)	•						•			
<i>Chionaspis alnus</i> Kuwana					•	•				
<i>Chionaspis saitamaensis</i> Kuwana						•		•		
<i>Chionaspis salicis</i> (Linnaeus)										•
<i>Chionaspis wistariae</i> Cooley	•			•		•		•		
* <i>Chrysomphalus aonidum</i> (Linnaeus)	•									
<i>Chrysomphalus bifasciculatus</i> Ferris				•		•		•	•	
* <i>Chrysomphalus dictyospermi</i> (Morgan)										•
<i>Comstockaspis macroporana</i> (Takagi)										•
<i>Comstockaspis perniciosa</i> (Comstock)	•		•		•	•	•		•	
<i>Cryptoparlatoarea leucaspis</i> Lindinger									•	
<i>Diaspidiotus cryptoxanthus</i> (Cockerell)	•					•		•		
<i>Diaspidiotus degeneratus</i> (Leonardi)						•		•	•	
<i>Diaspidiotus makii</i> (Kuwana)						•				
* <i>Diaspis boisduvalii</i> Signoret										•
* <i>Diaspis bromeliae</i> (Kerner)									•	
* <i>Diaspis echinocacti</i> (Bouché)									•	
<i>Duplacionaspis divergens</i> (Green)								•		
<i>Dynaspidiotus pseudomeyeri</i> (Kuwana)	•			•	•	•	•	•	•	
<i>Dynaspidiotus tsugae</i> (Marlatt)										•
<i>Fiorinia japonica</i> Kuwana						•		•		
* <i>Hemiberlesia lataniae</i> (Signoret)	•									
<i>Hypaspidiotus jordani</i> (Kuwana)						•		•	•	
<i>Kuwanaspis hikosani</i> (Kuwana)		•				•		•		
<i>Kuwanaspis howardi</i> (Cooley)						•		•		
<i>Kuwanaspis pseudoleucaspis</i> (Kuwana)	•					•			•	
<i>Lepidosaphes conchiformis</i> (Gmelin)						•	•	•	•	
<i>Lepidosaphes gloverii</i> (Packard)										•
<i>Lepidosaphes japonica</i> (Kuwana)				•						

Scientific Name	Distribution within Korea										
	GG	GW	CB	CN	GB	GN	JB	JN	JJ	NC	
<i>Lepidosaphes kamakurensis</i> (Kuwana)						•		•			
<i>Lepidosaphes okitsuensis</i> Kuwana						•					
<i>Lepidosaphes pallida</i> (Maskell)	•	•		•	•	•	•	•	•		
<i>Lepidosaphes pini</i> (Maskell)	•	•	•	•	•	•	•	•	•		
<i>Lepidosaphes pinnaeformis</i> (Bouché)			•			•		•	•		
<i>Lepidosaphes towadensis</i> Takagi and Kawai										•	
<i>Lepidosaphes tubulorum</i> Ferris				•							
<i>Lepidosaphes ulmi</i> (Linnaeus)										•	
<i>Lepidosaphes ussuriensis</i> (Borchsenius)	•			•	•	•	•		•		
<i>Lepidosaphes yanagicola</i> Kuwana	•	•						•	•		
<i>Lopholeucaspis japonica</i> (Cockerell)	•	•	•	•	•	•	•	•	•		
<i>Octaspidotus stauntoniae</i> (Takahashi)						•		•	•		
<i>Odonaspis secreta</i> (Cockerell)	•					•	•	•	•		
<i>Parlatoria camelliae</i> Comstock						•					
<i>Parlatoria pergandii</i> Comstock										•	
* <i>Parlatoria proteus</i> (Curtis)						•					
<i>Parlatoria theae</i> Cockerell	•			•	•	•	•	•	•		
<i>Pinnaspis aspidistrae</i> (Signoret)	•	•				•			•		
* <i>Pinnaspis buxi</i> (Bouché)										•	
<i>Pinnaspis chamaecyparidis</i> Takagi						•	•	•			
<i>Pinnaspis hikosana</i> Takagi				•							
* <i>Pinnaspis strachani</i> (Cooley)										•	
<i>Pinnaspis uniloba</i> (Kuwana)								•			
<i>Pseudaonidia duplex</i> (Cockerell)						•		•	•		
<i>Pseudaonidia paeoniae</i> (Cockerell)						•		•	•		
<i>Pseudaulacaspis cockerelli</i> (Cooley)	•	•	•	•	•	•	•	•	•		
<i>Pseudaulacaspis kiushiuensis</i> (Kuwana)	•					•			•		
<i>Pseudaulacaspis latiloba</i> (Takagi and Kawai)						•		•			
<i>Pseudaulacaspis pentagona</i> (Targioni-Tozzetti)	•	•	•	•	•	•	•	•	•		
<i>Pseudaulacaspis prunicola</i> (Maskell)	•					•	•	•			
<i>Pseudaulacaspis ulmicola</i> Tang	•										
<i>Rhizaspidotus canariensis</i> (Lindinger)		•									
<i>Selenomphalus distylii</i> Takagi										•	
<i>Unachionaspis tenuis</i> (Maskell)	•				•	•	•	•	•		
<i>Unaspis euonymi</i> (Comstock)	•	•	•	•	•	•	•	•	•		
<i>Unaspis yanonensis</i> (Kuwana)						•			•		

