

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Publications from USDA-ARS / UNL Faculty

U.S. Department of Agriculture: Agricultural
Research Service, Lincoln, Nebraska

2013

Soil type and moisture regime control microbial C and N mineralization in grassland soils more than atmospheric CO₂-induced changes in litter quality

Virginia L. Jin

USDA - Agricultural Research Service, virginia.jin@ars.usda.gov

Richard L. Haney

USDA - Agricultural Research Service

Philip A. Fay

USDA-ARS, philip.fay@ars.usda.gov

H. Wayne Polley

USDA-ARS, wayne.polley@ars.usda.gov

Follow this and additional works at: <https://digitalcommons.unl.edu/usdaarsfacpub>

Jin, Virginia L.; Haney, Richard L.; Fay, Philip A.; and Polley, H. Wayne, "Soil type and moisture regime control microbial C and N mineralization in grassland soils more than atmospheric CO₂-induced changes in litter quality" (2013). *Publications from USDA-ARS / UNL Faculty*. 1157.
<https://digitalcommons.unl.edu/usdaarsfacpub/1157>

This Article is brought to you for free and open access by the U.S. Department of Agriculture: Agricultural Research Service, Lincoln, Nebraska at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Publications from USDA-ARS / UNL Faculty by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.



Soil type and moisture regime control microbial C and N mineralization in grassland soils more than atmospheric CO₂-induced changes in litter quality

Virginia L. Jin^{a,*}, Richard L. Haney^b, Philip A. Fay^b, H. Wayne Polley^b

^aUSDA – Agricultural Research Service, Agroecosystem Management Research Unit, 137 Keim Hall, University of Nebraska – East Campus, Lincoln, NE 68583-0937, USA

^bUSDA – Agricultural Research Service, Grassland, Soil and Water Research Laboratory, 808 E. Blackland Road, Temple, TX 76702, USA

ARTICLE INFO

Article history:

Received 19 September 2012

Received in revised form

28 November 2012

Accepted 30 November 2012

Available online 25 December 2012

Keywords:

Litter decomposition

C mineralization

Net N mineralization

Grassland

Global change

Bouteloua curtipendula

ABSTRACT

Global change-induced alterations in litter quality and soil moisture regime will likely impact grassland C and N dynamics, but how these changes interact with edaphic properties across the landscape is unclear. We measured the effects of litter quality, soil type, soil moisture level, and soil drying-rewetting frequency (D-RW) on microbial C and N mineralization of litter and soil organic matter (SOM) in a full-factorial, controlled incubation experiment. Four levels of litter quality (no litter; or litter from *Bouteloua curtipendula* grown under 280, 380, 500 $\mu\text{L L}^{-1}$ CO₂) were surface-applied to three contrasting soils common to Blackland Prairie landscapes: an upland Mollisol, a lowland Vertisol, and a fluvial Alfisol. Different soil moisture regimes were tested by incubating soils at four moisture levels (air-dry, 25%, 35%, or 50% water-holding capacity, WHC) and by drying-rewetting soils 0, 1, 2, 4 or 8 times over the 112-d incubation period. Litter additions stimulated microbial activity, increasing total CO₂ production (i.e. C mineralized from litter + SOM decomposition) up to 17× more than no-litter controls (average 3×) and decreasing net N mineralization up to –3× less (average –0.5×) due to greater microbial N immobilization. Neither C nor N mineralization, however, was affected by litter quality. For all soils, litter decomposition increased with increasing WHC and D-RW frequency, but the average percent of total CO₂ derived from litter was a negative function of SOM content. Similarly, net N mineralization also was positively correlated with soil WHC and affected most strongly by soil type (Alfisol < Mollisol < Vertisol). Net N mineralization responses to D-RW events was also soil-specific, with Alfisol soils showing no response and Mollisol and Vertisol soils decreasing after 4 D-RW events. Our results suggest that predicted changes in rainfall patterns and its interactions with soil type across the landscape will control short-term C and N mineralization responses in grasslands to a greater extent than atmospheric CO₂-induced changes in litter C:N ratio for this common species of prairie grass.

Published by Elsevier Ltd.

1. Introduction

Global changes such as rising atmospheric CO₂ and altered precipitation regimes are expected to impact grassland productivity by directly or indirectly altering the availability of water, nutrients, and energy (i.e. carbon, C; Dukes et al., 2005; Soussana and Lüscher, 2007; Leakey et al., 2009). The interaction of these global changes with edaphic characteristics on soil microbial activity likely will affect whether grasslands sequester or release C. Little data, however, are available on more spatially-explicit responses to global changes, such as those driven by landscape variation in soils.

Small-scale spatial variation in soils caused by vegetation can significantly affect plant-soil responses to elevated atmospheric CO₂. For example, microbially-mediated C and N processes in soil microsites under shrubs are accelerated under elevated atmospheric CO₂ compared to soils in plant-free interspace microsites in a desert rangeland, particularly when water is less limiting (Billings et al., 2004; Jin and Evans, 2007; Schaeffer et al., 2007; Jin et al., 2011). In more mesic rangelands and grasslands, plant growth responses are mediated largely by elevated CO₂-induced increases in plant water-use efficiency and soil moisture availability (Morgan et al., 2004; Körner, 2006; Polley et al., 2008) and, in some cases, greater N mineralization (Dijkstra et al., 2007, 2010). Furthermore, in a Blackland prairie system, soil type strongly affects how plant productivity and grassland species composition respond to increased soil water availability and subsequent changes to soil N

* Corresponding author. Tel.: +1 402 472 5137; fax: +1 402 472 0516.

E-mail address: Virginia.jin@ars.usda.gov (V.L. Jin).

availability under elevated CO₂ (Polley et al., 2003, 2012a, 2012b; Fay et al., 2012).

Soil CO₂ effluxes are high in tallgrass prairies compared to other temperate terrestrial ecosystems because of their relatively high plant productivity, annual precipitation, and soil organic matter (SOM) content (Mielenick and Dugas, 2000). Changes in the amount and variability of precipitation, however, are likely to regulate soil CO₂ effluxes and their underlying drivers (Fay et al., 2011) and responses to elevated atmospheric CO₂ (Jastrow et al., 2000). For example, grassland soil CO₂ fluxes decreased with increased time interval between rainfall events (Fay et al., 2000; Harper et al., 2005), in part due to reduced soil moisture and plant productivity and expected C input to soils (Fay et al., 2008).

More than half of the CO₂ emitted by grassland soils is from microbial SOM decomposition (Wan and Luo, 2003; Millard et al., 2008). Litter inputs and litter decomposition, however, confer substantial variability to overall soil CO₂ fluxes and the transfer of plant C into SOM. Increases in litter inputs may lead to more C stored as SOM (Franck et al., 1997; Gorissen and Cotrufo, 2000; Jastrow et al., 2000), and changes in litter quality affect the partitioning of C into different SOM pools (Melillo et al., 1982; Cotrufo et al., 1998; Guo and Gifford, 2002). Increased C inputs, however, can stimulate SOM decomposition (i.e. priming effect) and limit grassland C storage potential (Kuzyakov et al., 2000; Xie et al., 2005; Niklaus and Fallool, 2006; Carney et al., 2007). Further, the relationship between litter C:N ratio, which typically increases in plants grown under elevated CO₂, and litter decomposition rate is highly variable (Franck et al., 1997; Gorissen and Cotrufo, 2000; Norby et al., 2001). Changes in the quantity and/or quality of plant litter inputs into soil can result in either net mineralization or immobilization of N and feed back to affect litter decomposition (Torbert et al., 2000). Interactions between altered litter inputs and precipitation regime, therefore, could affect litter decomposition and grassland soil C storage.

The objectives of this study were to evaluate how global change-induced alterations in litter quality and soil moisture regime interact with edaphic characteristics to affect microbial C and N mineralization. Microbial C mineralization was quantified as the decomposition of organic C in litter and SOM to CO₂. Microbial N mineralization, the conversion of organic N to inorganic N (i.e. ammonium + nitrate), was quantified as the net change in soil inorganic N over the incubation period. A positive net change indicates microbial mineralization of N, and a negative net change indicates microbial immobilization of N. We removed the confounding influences of root respiration, climatic variation in the

field, and the cumulative impact of CO₂ enrichment on soils by incubating plant litter grown under three CO₂ levels with soils that had no exposure to elevated atmospheric CO₂ (Paul et al., 2001; Collins et al., 2000). Because short-term C mineralization rapidly quantifies litter quality differences (Haney et al., 2001; Haney and Franzleubbers, 2009), controlled incubations were conducted using four litter treatments (no litter; or litter from *Bouteloua curtipendula* grown under 280, 380, 500 μL L⁻¹ CO₂) on three contrasting soils common to Blackland Prairie landscapes (Alfisol, Mollisol, Vertisol). Total CO₂ production and net N mineralization were measured in soils incubated in a full-factorial experiment with four levels of soil moisture and five levels of drying–rewetting frequency to represent different precipitation regimes. In addition, water-extractable organic C and organic N concentrations in pre- and post-incubated soils were measured to evaluate whether changes in these potentially labile pools correlated with microbial C and N activities. We hypothesized that: (1) litter additions would stimulate microbial activities (decomposition, net N mineralization) in all soils; (2) differences in litter C:N ratio due to growth-CO₂ level would be reflected by differences in microbial activity; (3) C and N mineralization would increase with greater soil moisture; and (4) soil-specific properties would interact with soil moisture regime as the dominant factor controlling litter decomposition and net N mineralization.

2. Materials and methods

2.1. Soil and litter collection

Soils (0–15 cm) from three soil orders commonly found in the Blackland Prairie region of Texas were collected in Spring 2008. These soils represent a broad range of texture, N and C contents, and hydrologic properties for grasslands in the southern portion of the U.S. Central Plains. An Alfisol (Bastsil series), Mollisol (Austin series), and Vertisol (Houston Black series) were collected from U.S. Army Corp of Engineers Stillhouse Hollow Lake area (Belton, TX), the USDA-ARS's Grassland, Soil, and Water Research Laboratory (GSWRL; Temple, TX), and the GSWRL watershed network (Riesel, TX), respectively. Bastsil soils are loamy fine sands (fine-loamy, siliceous, active, thermic Udic Paleustalfs) formed of alluvial sediments and commonly found on stream terraces. Austin soils are silty clays (fine-silty, carbonatic, thermic Udorthentic Haplustolls), typical of erosional uplands. Houston Black soils are heavy shrink-swell clays (very-fine, smectitic, thermic Udic Haplusterts) that dominate lowland areas. Selected soil properties are shown in Table 1. Fresh

Table 1

Initial properties for soils (0–15 cm) and *Bouteloua curtipendula* litter used in laboratory incubations (mean ± se; n = 3). Different letters for each property indicate significant differences between soil series or between growth-CO₂ levels (P ≤ 0.05). Soil texture data from Fay et al. (2009).

Soil order	Unit	Alfisol	Mollisol	Vertisol
Soil series name	–	Bastsil	Austin	Houston Black
Sand	%	71.9 ± 0.9a	11.7 ± 0.4b	10.9 ± 0.6b
Silt	%	20.8 ± 0.5a	45.3 ± 0.7b	43.0 ± 0.6b
Clay	%	7.3 ± 0.4a	39.3 ± 1.0b	49.8 ± 1.4c
Total inorganic C	g C kg ⁻¹	0.3 ± 0.1a	68.2 ± 1.0b	64.6 ± 1.0c
Total soil organic C (SOC)	g C kg ⁻¹	4.1 ± 0.4a	17.8 ± 0.4b	21.8 ± 0.6c
Total N	g N kg ⁻¹	0.6 ± 0.1a	1.7 ± 0.1b	5.6 ± 0.1c
Water-extractable SOC	mg C kg ⁻¹	137 ± 62a	226 ± 24b	376 ± 38c
Water-extractable organic N (ON)	mg N kg ⁻¹	16.3 ± 0.2a	3.9 ± 0.3b	9.8 ± 0.3c
Water-extractable inorganic N (IN)	mg N kg ⁻¹	7.4 ± 0.4a	9.0 ± 0.5b	7.9 ± 0.3a
<i>B. curtipendula</i> growth-CO ₂ level		280 μL L ⁻¹	380 μL L ⁻¹	500 μL L ⁻¹
Initial litter C	mg C g ⁻¹	377 ± 3a	397 ± 1b	384 ± 2c
Initial litter N	mg N g ⁻¹	5.6 ± 0.1a	7.3 ± 0.1b	18.9 ± 2.0c
Initial litter C:N	–	67.3 ± 0.3a	53.1 ± 2.0b	22.7 ± 1.9c
C added in 0.25 g dry litter	mg C	94	99	96
N added in 0.25 g dry litter	mg N	1.4	1.8	4.7

soils were passed through a 2 mm sieve, large organic particles and/or coarse roots removed by hand, and air-dried before use.

Litter from *B. curtipendula* (Michx.) Torr., a widely distributed mid-grass prairie species, was harvested from the USDA-ARS's Lysimeter CO₂ Gradient (LYCOG) experiment (Temple, TX USA; 31°05'N, 97°20'W; Polley et al., 2008; Fay et al., 2009). Plants growing in Alfisol monoliths at 280, 380, or 500 μL L⁻¹ CO₂ sections within LYCOG were harvested at the end of the 2007 growing season as standing dead plant material. Only Alfisol monoliths yielded sufficient grass litter for this experiment at all CO₂ levels. Aboveground biomass was clipped at 5 cm above the soil surface and dried to constant mass at 60 °C. For each growth-CO₂ level, dried litter was cut into 1-cm segments, homogenized, then surface-applied to soils prior to incubation.

2.2. Incubation design

Soils were incubated in triplicate over a 112-day incubation period (16 weeks) at 25 ± 1 °C in the dark using a full factorial design with 3 soil types, 4 litter treatments (no litter; low, ambient, and high growth-CO₂), 4 soil moisture levels (air-dry, 25%, 35%, 50% mean daily water holding capacity, or WHC), and 5 dry-rewet frequencies (D-RW; 0, 1, 2, 4, 8 events). Soil moisture treatments were selected to represent a broad range of current and potential precipitation regimes, including prolonged drought (i.e. constant air-dry) to frequent abundant precipitation (i.e. 50% WHC up to 8 D-RW events). In addition, the standard 25 °C incubation temperature used here closely approximates surface soil temperatures experienced in this region during the growing season (i.e. 4-year average of monthly soil T at 5 cm depth was ~26.5 °C from March through October; Neitsch et al., 2011). The 16-week incubation period was similar to those used in other soil and litter incubation studies (Fierer and Schimel, 2002; Gordon et al., 2008; Marhan et al., 2008; Rasmussen et al., 2008).

Soils in the air-dry treatment were maintained at that level throughout the incubation (i.e. D-RW frequency of 0 only), resulting in a total of 576 soil beakers incubated (192 beakers per soil type). For each incubation sample, 40 g of dry soil was added to a 50-mL polypropylene disposable beaker. Dry litter (0.25 g) was surface-applied to soils (i.e. no incorporation) to better represent field inputs of aboveground litter to soil. Corresponding rates of litter C and N applied per treatment level are shown in Table 1. Each soil beaker was placed in a gas-tight 1-quart canning jar containing a 10 mL alkali trap (1.0 M KOH) to absorb CO₂ and sufficient water to maintain humidity. Alkali traps were replaced on the third day after each re-wetting treatment, then every 7 or 10 days until the next D-RW event throughout the incubation period. Jar-blank traps were also collected in triplicate at each time point from non-soil incubation jars that contained all materials except soils (i.e. trap, water, empty beaker) to provide blank corrections for CO₂ trapped from air inside the incubation jar itself.

Total CO₂ production (C_{total} ; mg CO₂-C kg⁻¹ dry soil) from mineralization of soil organic matter (C_{SOM}) and litter (C_{litter}) was determined by adding 2 M BaCl₂ to alkali traps and titrating to neutral pH with 1.0 M HCl (Anderson, 1982). All CO₂ values from soil incubation jars were first blank-corrected using non-soil incubation values. Litter-derived CO₂ was then calculated as the difference between litter-amended soils and no-litter controls (i.e. $C_{litter} = C_{total} - C_{SOM}$). We assumed that litter addition had no priming effect on SOM decomposition.

Because soil textures differed considerably, soil moisture levels were standardized across all soil types using water holding capacity (WHC, %). 100% WHC was determined for each soil type as the gravimetric water content of soil that was saturated and allowed to drain freely over 6 h in a filter funnel. Air-dry treatments received

no water additions. Moist soil treatments were maintained at an average daily level of 25%, 35%, or 50% WHC over the incubation period. Water holding capacity treatment level was maintained by adding deionized water to soils whenever alkali traps were replaced. Calculation of average daily %WHC included days when soils were undergoing drying-rewetting (D-RW) treatments.

Soils exposed to D-RW treatments underwent 0, 1, 2, 4, or 8 drying events spaced evenly over the 112-day incubation period. Each dry-down event lasted 7 days (i.e. 1 D-RW event = 7 days drying; 2 D-RW events = 14 days drying; 4 D-RW events = 28 days drying; and 8 D-RW events = 56 days drying). Because soils exposed to more D-RW events experienced a greater number of days below the target daily %WHC level, these soils were rewetted to a slightly higher WHC level after each dry-down event to maintain the target daily average %WHC over the 112-day incubation period (*sensu* Fierer and Schimel, 2002) (Fig. 1). For each drying event, jar lids were removed to allow air-flow in the incubator to dry the soil. By day 7 of the dry-down period, soil %WHC approximated air-dry %WHC. The CO₂ produced during the total dry-down period was estimated by measuring one-day CO₂ at days 2, 4, 7 and linearly interpolating daily CO₂ production for intervening days. The CO₂ produced during the total dry-down period was then calculated by summing daily CO₂ production over the 7 days.

2.3. Soil and plant analyses

Soil concentrations of total organic C and N, water-extractable organic C and N, and water-extractable inorganic N (NH₄⁺-N + NO₂⁻-N + NO₃⁻-N) were measured in pre- and post-incubated soils. Total soil organic C (SOC), soil total N (TN), and litter C and N was determined by dry combustion at 600 °C, and total soil C was measured by combustion at 900 °C (Vario Max CHN, Elementar, Hanau, Germany) (Fay et al., 2009). Soil inorganic C was calculated as the difference between total C and SOC.

Soils were extracted with deionized water (1:10), shaken for 1 h, centrifuged at 2500 rpm for 15 min, then the supernatant filtered through Whatman #1 filter paper. Dissolved inorganic C was removed from a 25 mL aliquot of each filtered extract by acidifying with 1 N HCl prior to measuring total dissolved organic C and total dissolved N (Apollo 9000, Teledyne Tekmar, Mason, Ohio).




Fig. 1. Water holding capacity (WHC, %) for all soils and all drying treatments at the mean daily 25% WHC level over the 112-day incubation period. Values for the mean daily 35% and 50% WHC treatments followed similar patterns (data not shown).

Inorganic N was measured in non-acidified filtered extracts using continuous flow colorimetry (Flow Solution IV, OI Analytical, College Station, Texas). Organic N concentrations were calculated as the difference between total N and inorganic N. Net N mineralization (mg N kg^{-1}) was calculated as the difference in inorganic N between pre- and post-incubated soils.

2.4. Statistical analyses

Variables measured on pre-incubated soils and litter were analyzed using one-way analyses of variance (ANOVAs; PROC GLM) to test for differences in soil type and growth CO_2 level, respectively. In post-incubated soils, four-way ANOVAs were used to test the fixed effects of soil type, litter, WHC, and D-RW treatments on C_{total} , C_{litter} , and on the relative contribution of litter mineralization (%) to total cumulative CO_2 production. Final post-incubation soil C and N concentrations (total and water-extractable) and changes in water-extractable C and N concentrations (i.e. post-incubated – pre-incubated value) were also analyzed with four-way ANOVAs. Type III statistics are reported. Data were tested for normality using the Shapiro–Wilk statistic and transformed when necessary. Post-hoc multiple comparisons between significant treatment means were tested for differences using the Fisher's least significant difference (LSD) procedure. Linear regressions were used to examine correlations between cumulative CO_2 production and net N mineralization vs. total and water-extractable C and N. All statistical tests were performed using SAS 9.1 (SAS, Inc., Cary, NC, USA).

3. Results

3.1. Pre-incubated soils and litter

Initial soil values for almost all physical and chemical properties differed by soil type ($P \leq 0.05$; Table 1). Initial C and N concentrations of *B. curtipendula* litter differed by growth CO_2 treatment level ($P \leq 0.05$; Table 1). Although there were small but significant differences in litter C due to growth CO_2 , differences in litter C:N ratio were driven by changes in litter N. Initial litter N values increased significantly with growth CO_2 concentration. As a result, litter C:N ratios decreased from 67 to 23 with increasing growth CO_2 concentration ($P \leq 0.05$).

3.2. Litter and SOM decomposition

Differences in litter C:N did not affect total C mineralization (C_{total} ; $\text{mg CO}_2\text{-C kg}^{-1}$ dry soil) from any soil under any soil moisture regime (WHC, D-RW) over the 112-day incubation. As a result, data were pooled across all litter additions, and results were re-analyzed using only two litter treatment levels: no litter ($n = 3$) vs. litter added ($n = 9$). Total CO_2 production was affected by all treatment interactions ($F_{16,480} = 2.40$; $P_{\text{Soil} \times \text{WHC} \times \text{D-RW} \times \text{Litter}} = 0.002$) (Fig. 2), but variability in total CO_2 production was dominated by soil ($F_{2,480} = 15,033.2$), litter ($F_{1,480} = 2786.63$), WHC ($F_{2,480} = 2702.06$), the two-way interaction between soil and WHC ($F_{4,480} = 1070.60$), and D-RW ($F_{4,480} = 369.29$) (all $P < 0.0001$). Total




Fig. 2. Carbon mineralized from soils and litter as: (a–c) total CO_2 produced ($\text{mg C} \cdot \text{kg soil}^{-1}$) from soil + litter, C_{total} ; (d–f) the percent of total CO_2 produced from litter ($C_{\text{litter}}/C_{\text{total}} \times 100$); and (g–i) the percent of litter remaining after 112 days. Litter addition treatments were averaged because growth CO_2 treatment was not significant ($n = 9$ per point). Standard errors are present but obscured by symbols when variability is low. Open symbols in 25% WHC panels indicate air-dried soil values.

CO₂ production was highest in finer-textured Vertisol soils, and lowest in coarser-textured Alfisol soils, and soil differences became more pronounced as %WHC increased (Fig. 2a–c).

Litter additions significantly increased total CO₂ production in all soils up to 17× (average 3×) more than no-litter controls at the same WHC and dry-rewet frequency (controls not shown). Litter decomposition dominated total CO₂ production in the coarse-textured, low organic matter Alfisol soils (54%–90% of total C mineralized, average 77%; Fig. 2d–f). In Mollisol soils, litter decomposition contributed to approximately half of total CO₂ production (41%–66%; average 52%). In the fine-textured, high organic matter Vertisol soils, total CO₂ produced was dominated by SOM decomposition, with litter decomposition contributing a minor but consistent percentage (13%–27%; average 20%). The total amount of litter decomposed increased (i.e. lower % litter C remaining; Fig. 2g–i) in all soils over the 112-day incubation as both %WHC and D-RW frequency increased, with significantly more litter decomposed in the finer-textured Mollisol and Vertisol compared to the Alfisol at 50% WHC.

3.3. Net N mineralization

Similar to total CO₂ production, differences in litter C:N did not affect net N mineralization for any soil or soil moisture regime over the 112-day incubation, so the litter C:N levels data were pooled re-analyzed using only two levels, no litter vs. litter added. Net N

mineralization was affected by two-way interactions soil and litter ($F_{2,480} = 3.18$; $P_{\text{Soil} \times \text{Litter}} = 0.043$), WHC and litter ($F_{3,480} = 4.59$; $P_{\text{WHC} \times \text{Litter}} = 0.004$), and the three way interaction between soil, WHC, and D-RW ($F_{16,480} = 9.72$; $P_{\text{Soil} \times \text{WHC} \times \text{D-RW}} < 0.0001$). Net N mineralization was most strongly affected by soil type ($F_{2,480} = 488.35$), WHC ($F_{3,480} = 372.25$), the soil × WHC interaction ($F_{4,480} = 130.68$), litter ($F_{1,480} = 125.76$), and D-RW ($F_{4,480} = 19.22$) (all $P < 0.0001$).

Net N mineralization generally was lowest in the Alfisol and highest in the Vertisol, except at 25% WHC where net N mineralization was lowest in the Mollisol. Soil differences in net N mineralization grew more pronounced as %WHC increased (Fig. 3a–f). Adding litter decreased net N mineralization in all soils up to –3× (average –0.5×), indicating enhanced microbial N immobilization (Fig. 3g–i). Greatest levels of N immobilization occurred in the coarser-textured Alfisol compared to finer-textured Mollisol and Vertisol soils.

Net N mineralization in all soils averaged across all D-RW levels increased with %WHC (Fig. 3). The responses to D-RW frequency differed among soil type and WHC level (Fig. 3). Net N mineralization in the Alfisol were consistently low and did not respond to D-RW frequency at any WHC level, but both finer-textured soils showed D-RW frequency effects. In Mollisol soils, net N mineralization did not respond to D-RW frequency in no-litter controls, but decreased with D-RW frequency when soils were amended with litter at 25% WHC. In both no-litter and litter-amended Mollisol




Fig. 3. Net N mineralized (mg N · kg soil⁻¹) over the 112-day incubation in: (a–c) soils only; (d–f) soil + litter; and (g–i) due to litter only. Litter addition treatments were averaged because growth CO₂ treatment was not significant ($n = 9$ per point). Values below dotted lines indicate net N immobilization. Standard errors are present but obscured by symbols when variability is low. Open symbols in 25% WHC panels indicate air-dried soil values.

soils at 35% WHC, net N mineralization increased up to 2–4 D-RW events then stabilized, but dropped after 4 D-RW events at 55% WHC. In Vertisol soils, net N mineralization increased with D-RW frequency at 25% WHC, decreased with D-RW frequency at 35% WHC, and decreased after 2–4 D-RW events at 55% WHC.

3.4. Changes in water-extractable DOC

Changes in water-extractable DOC (ΔDOC ; post-incubated – pre-incubated concentrations) were calculated to evaluate whether decomposition of this potentially labile C pool was correlated with measured C and N mineralization. A positive ΔDOC value indicates a net gain in DOC over the 112-day incubation, whereas a negative value indicates a net loss of DOC. Changes in water-extractable DOC were affected by the three-way interaction between soil, WHC, and D-RW ($F_{6,555} = 4.27$; $P_{\text{Soil} \times \text{WHC} \times \text{D-RW}} < 0.0001$) and the two-way interaction between soil and litter growth- CO_2 level ($F_{6,555} = 43.03$; $P_{\text{Soil} \times \text{Litter } \text{CO}_2} < 0.0001$). Mean ΔDOC ($\pm\text{se}$) were soil-specific but indicated DOC losses in all soils over the 112-day incubation. Litter additions stimulated greater losses of DOC compared to no-litter controls, with greatest losses occurring in the Alfisol ($-149 \pm 10\%$) and the Mollisol ($-131 \pm 21\%$), and lowest losses in the Vertisol ($-24 \pm 16\%$). DOC losses were enhanced by increasing %WHC and D-RW frequency. Further, DOC losses were higher in all soils amended with litter grown at $500 \mu\text{L L}^{-1}$ compared to 280 or $380 \mu\text{L L}^{-1}$ (Fig. 4). Litter growth- CO_2 treatment differences in ΔDOC , however, did not translate to differences in total CO_2 production or net N mineralization in any soil at any WHC or D-RW level. Instead, all soils showed that DOC losses correlated with increases in total CO_2 production (Alfisol: $-7.2x - 128.9$, $R^2 = 0.6127$, $P < 0.0001$; Mollisol: $-4.3x + 959.3$, $R^2 = 0.2676$, $P < 0.0001$; Vertisol: $-11.9x + 3159.4$, $R^2 = 0.4168$, $P < 0.0001$; Fig. 5a). Only Alfisol soils also showed decreased N mineralization with greater DOC losses ($0.13x + 33.6$, $R^2 = 0.5738$, $P < 0.0001$; Fig. 5b).

Changes in water-extractable organic N (ΔDON ; post-incubated – pre-incubated concentrations) were not affected by litter additions or changes in WHC or D-RW level, and were not correlated to either total CO_2 production or net N mineralization (data not shown). Averaged over all treatments, mean ΔDON ($\pm\text{se}$) was -12.3 ± 0.4 , 1.9 ± 0.7 , and $4.4 \pm 1.3 \text{ mg N kg}^{-1}$ in Alfisol, Mollisol, and Vertisol soils, respectively.




Fig. 4. Change in water-soluble organic C concentration (ΔDOC , post-incubated – pre-incubated soil values; $\text{mg C} \cdot \text{kg soil}^{-1}$). Values are averaged over all WHC and D-RW treatments for each soil ($\pm\text{se}$). Different letters within each soil type indicate significant differences in litter treatments ($P \leq 0.05$).




Fig. 5. Relationship between water-soluble organic C concentration (ΔDOC , post-incubated – pre-incubated soil values; $\text{mg C} \cdot \text{kg soil}^{-1}$) and: (a) total CO_2 produced ($\text{mg C} \cdot \text{kg soil}^{-1}$; Alfisol: $-7.2x - 128.9$, $R^2 = 0.6127$, $P < 0.0001$; Mollisol: $-4.3x + 959.3$, $R^2 = 0.2676$, $P < 0.0001$; Vertisol: $-11.9x + 3159.4$, $R^2 = 0.4168$, $P < 0.0001$); and (b) net N mineralized ($\text{mg N} \cdot \text{kg soil}^{-1}$; Alfisol: $0.13x + 33.6$, $R^2 = 0.5738$, $P < 0.0001$). Open symbols are no-litter control soils, closed symbols are litter-amended soils.

4. Discussion

Understanding the local mechanisms controlling soil–plant interactions and potential ecosystem feedbacks is critical for predicting landscape-level impacts of global change drivers. In this study, edaphic properties and average soil moisture exerted greater control than variability in soil moisture or litter quality on C and N mineralization in these temperate grassland soils. Litter additions increased total CO_2 production in all soils compared to no-litter controls, but decreased net N mineralized by presumably increasing microbial N immobilization, partially supporting our first hypothesis. Net changes in water-extractable soil organic C (i.e. ΔDOC) over the 112-d incubation period indicated the greatest DOC losses (i.e. consumption or immobilization of DOC) in all soils amended with litter grown under elevated CO_2 , but this did not translate to growth CO_2 -specific changes in C or N mineralization, lending no support to our second hypothesis. Instead, increasing losses in DOC were broadly correlated with greater total CO_2 production in all soils and decreased N mineralization in Alfisol soils. Overall, soil type and soil water availability far outweighed the minor effects of drying-rewetting (D-RW) frequency, and soil-specific interactions controlled microbial activities, supporting our third and fourth hypotheses.

4.1. Effects of growth CO_2 level on litter quality

Litter C:N ratios typically are higher in plants grown at elevated than ambient CO_2 because N is diluted by the additional carbohydrates accumulated at high CO_2 (Gorissen and Cotrufo, 2000;

Loladze, 2002; Taub and Wang, 2008). In contrast to this expectation, which also formed part of our first hypothesis, we found that litter C:N decreased with increasing growth CO₂-concentration. A previous study from this site reported the expected positive correlation between C:N in plants harvested mid-season with growth CO₂ concentration (Gill et al., 2002), but this initial correlation did not persist after additional years of CO₂ treatment (Polley et al., 2003). Indeed, the most recent study from this site reported no significant CO₂ enrichment effects on the N concentration of grass tillers at mid-season on any of the three soil types (Polley et al., 2011). In this study, we measured C:N values in standing dead or senesced grass litter harvested at the end of the season. Litter C:N ratios at this time may have been influenced by N uptake late in the growing season and/or differences in end-of-season N translocation among CO₂ treatments. Specifically, CO₂ enrichment may have increased soil N availability to plants late in the season by reducing transpiration rates (Polley et al., 2008) and increasing soil water content (Jones and Donnelly, 2004; Dijkstra et al., 2008, 2010), resulting in greater litter N concentrations at higher CO₂. CO₂ enrichment also has been shown to reduce translocation of N from aboveground biomass between mid- and late-season (Dijkstra et al., 2008). A similar mechanism may have contributed to the significant decrease in litter C:N at high CO₂ that we observed.

4.2. Litter effects on C and N mineralization

Litter additions stimulated microbial activity, increasing total CO₂ production up to 17× more than no-litter controls (average 3×) and decreasing net N mineralization up to −0.5× less (average −0.5×) due to greater microbial N immobilization. Neither C nor N mineralization, however, was affected by differences in litter quality as observed by others (O'Neill and Norby, 1996; Cotrufo et al., 1998; Gorissen and Cotrufo, 2000; Torbert et al., 2000) but in contrast to other studies which found that total CO₂ production decreases with increasing litter C:N (Melillo et al., 1982; Moretto et al., 2001; Norby et al., 2001; Marhan et al., 2008). One potential source of error, which was not quantified here due to methodological constraints, is the role of soil-specific priming on respired CO₂. Although we were not able to quantitatively partition respired CO₂ into component sources of litter decomposition vs. soil-specific priming, multiple studies have shown that litter-associated soil priming can enhance SOM mineralization and limit soil C sequestration potential (Kuzakov et al., 2000; Xie et al., 2005; Niklaus and Falloon, 2006; Carney et al., 2007).

Differences in litter C:N also did not affect water-soluble soil organic C or organic N concentrations (Marhan et al., 2008) in pre-incubated or post-incubated soils. Litter C:N, however, did affect the change in water-soluble, or dissolved, organic C (Δ DOC; post-incubated concentrations – pre-incubated concentrations). A negative Δ DOC value suggests that more labile C was immobilized or consumed over the incubation period and potentially released as CO₂. The absence of litter C:N differences due to growth-CO₂ treatment effects on microbial activity suggests that the additional DOC in soils amended with litter grown under elevated CO₂ may have been immobilized in microbial biomass, transformed to a less bioavailable form of organic C, or sorbed to the soil matrix. Furthermore, changes in DOC concentrations were a magnitude smaller than soil respiration, so changes in Δ DOC due to litter quality may have been too low to detect as respired C. Although these litter C:N differences may have been expressed after a longer incubation period, shorter-term C mineralization and the formation/stabilization of SOC will likely be controlled by litter quantity and its interaction with the soil matrix more than litter quality (Liu et al., 2009; Gentile et al., 2011).

4.3. Soil type effects

Relationships between C mineralization and distinct soil properties such as mineralogy (Percival et al., 2000; Rasmussen et al., 2008), soil microsite (Billings et al., 2004; Schaeffer et al., 2007), and soil aggregate fractions (Williams et al., 2000; Gentile et al., 2011) suggest that variation in soil properties linked to landscape position will influence landscape-level ecosystem responses to global changes. In this study, soil type was a dominant factor affecting both litter and SOM decomposition in temperate grassland soils. In addition, litter additions interacted with soil type to affect total CO₂ production over the 112-day incubation period. In general, the C mineralized from litter decomposition contributed far less to the total CO₂ production in the fine-textured soils (Mollisol, Vertisol) than in coarse-textured soils (Alfisol). Decomposition of native SOM dominated CO₂ production in the Vertisol and less so in the Mollisol, whereas litter decomposition was the primary C source for mineralization in the low organic, coarse-textured Alfisol. Increased C inputs from added litter also could have further stimulated SOM decomposition in the Mollisol and Vertisol soils compared to the Alfisol soil, but partitioning a priming effect was beyond the scope of this experiment. In these Blackland Prairie soils, soil type was the primary controller determining the relative contribution of litter-C to total soil C mineralized.

Soil-specific differences in N mineralization and immobilization were likely due to different sized soil N pools in each soil type, with the Vertisol soils having a significantly larger total N pool compared to the other two soil types. Further, this large native N pool in the Vertisol soils may have allowed greater net N mineralization to occur at low WHC levels, which limited litter decomposition and microbial N immobilization.

4.4. Soil moisture effects

Average soil moisture also was a major factor controlling litter and SOM decomposition. In this study, total CO₂ production increased with increasing %WHC, and the %WHC effect became more pronounced as SOM content increased. Further, soil drying-rewetting enhanced CO₂ production compared to constantly moist soils, primarily because drying-rewetting stimulated SOC mineralization to a greater extent than litter decomposition in all soils. Similarly, net N mineralization increased with WHC, and was sensitive to D-RW frequency in the fine-textured Mollisol and Vertisol soils, but not in the coarse-textured Alfisol soils. Although the average annual soil CO₂ flux in tallgrass prairie soils increases with annual precipitation (Mielnick and Dugas, 2000), pulses in water and subsequent nutrient availability after rainfall trigger pulses in biological activity (Fierer and Schimel, 2002; Austin et al., 2004; Xiang et al., 2008). Some modeling studies estimate that pulse responses can account for up to 14% of annual CO₂ fluxes in some ecosystems (Yuste et al., 2005; Li et al., 2006). Other studies have also found increased cumulative CO₂ production from soils undergoing multiple drying-rewetting events compared to soils kept at constant optimum water content (Denef et al., 2001; Miller et al., 2005; Xiang et al., 2008), although some studies have shown decreases (Franzluebbers et al., 1994; Fierer and Schimel, 2002; Mikha et al., 2005). In one study, less frequent but larger rainfall events were linked to lower soil CO₂ emissions because substrate limitation exerted greater control on soil CO₂ efflux relative to water or temperature under this more extreme rainfall pattern (Harper et al., 2005).

When soils remained wetter for longer (i.e. decreased D-RW frequency), the contribution of litter-C to total CO₂ produced increased. This is consistent with a field study showing that lower soil moisture status due to reduced precipitation treatment in

tallgrass prairie can significantly reduce litter decomposition (Reed et al., 2009). Because these negative effects can persist even after precipitation inputs are returned to normal (Reed et al., 2009), interannual variability in soil moisture has both short- and long-term effects on soil C mineralization (Fierer and Schimel, 2002; Harper et al., 2005).

5. Conclusions

Nutrient and energy flows in grassland ecosystems under increasing atmospheric CO₂ ultimately will depend on how microbially-mediated soil processes respond to variability in current and future rainfall patterns (Weltzin et al., 2003; Mikha et al., 2005; Schimel et al., 2007; Gordon et al., 2008). Results from this study imply that grassland C mineralization will be controlled more in the short-term by changes in the amount of precipitation and its interactions with soil type than by CO₂-caused changes in litter C:N ratio. Thus, scaling-up controlled incubation results to complex landscape-level responses will be constrained by how landscape variation in soils interacts with this and other grassland species to affect how C and N cycling in grassland ecosystems will respond to altered precipitation patterns and increasing atmospheric CO₂ (Polley et al., 2003; Dijkstra et al., 2010; Fay et al., 2011, 2012; Polley et al., 2012a, 2012b).

Acknowledgments

The authors thank R. Gill for his constructive suggestions on the incubation design and D. Miller for his comments. We also thank A. Gibson, K. Jones, C. Kolodziejczyk, A. Naranjo, K. Tiner, and students M. Arredondo, P. Knutson, J. Roberts, and J. Roessler for technical support. Mention of trade names or commercial products in this publication does not imply recommendation or endorsement by the U.S. Department of Agriculture. USDA is an equal opportunity provider and employer.

References

- Anderson, J.P.E., 1982. Soil respiration. In: Page, A.L., Miller, R.H., Keeney, D.R. (Eds.), *Methods of Soil Analysis, Part 2. Chemical and Microbiological Properties*. Soil Science Society of America, Madison, WI, pp. 831–871.
- Austin, A.T., Yahdjian, L., Stark, J.M., Belnap, J., Porporato, A., Norton, U., Ravetta, D.A., Schaeffer, S.M., 2004. Water pulses and biogeochemical cycles in arid and semiarid ecosystems. *Oecologia* 141, 221–235.
- Billings, S.A., Schaeffer, S.M., Evans, R.D., 2004. Soil microbial activity and N availability with elevated CO₂ in Mojave Desert soils. *Global Biogeochemical Cycles* 18, GB1011. <http://dx.doi.org/10.1029/2003GB002137>.
- Carney, K.M., Hungate, B.A., Drake, B.G., Megegnal, J.P., 2007. Altered soil microbial community at elevated CO₂ leads to loss of soil carbon. *Proceedings of the National Academy of Sciences* 104, 4990–4995.
- Collins, H.P., Elliott, E.T., Paustian, K., et al., 2000. Soil carbon pools and fluxes in long-term corn belt agroecosystems. *Soil Biology and Biochemistry* 32, 157–168.
- Cotrufo, M.F., Briones, M.J.I., Ineson, P., 1998. Elevated CO₂ affects field decomposition rate and palatability of tree leaf litter: importance of changes in substrate quality. *Soil Biology and Biochemistry* 30, 1565–1571.
- Denef, K., Six, J., Bossuyt, H., Frey, S.D., Elliott, E.T., Merckx, R., Paustian, K., 2001. Influence of dry–wet cycles on the interrelationship between aggregate, particulate organic matter, and microbial community dynamics. *Soil Biology and Biochemistry* 33, 1599–1611.
- Dijkstra, F.A., West, J.B., Hobbie, S.E., Reich, P.B., Trost, J., 2007. Plant diversity, CO₂, and N influence inorganic and organic N leaching in grasslands. *Ecology* 88, 490–500.
- Dijkstra, F.A., Pendall, E., Mosier, A.R., King, J.Y., Milchunas, D.G., Morgan, J.A., 2008. Long-term enhancement of N availability and plant growth under elevated CO₂ in a semi-arid grassland. *Functional Ecology* 22, 975–982.
- Dijkstra, F.A., Blumenthal, D., Morgan, J.A., LeCain, D.R., Follett, R.F., 2010. Elevated CO₂ effects on semi-arid grassland plants in relation to water availability and competition. *Functional Ecology* 24, 1152–1161.
- Dukes, J.S., Chiariello, N.R., Cleland, E.E., Moore, L.A., Shaw, M.R., Thayer, S., Tobeck, T., Mooney, H.A., Field, C.B., 2005. Responses of grassland production to single and multiple global environmental changes. *PLoS Biology* 3, e319. <http://dx.doi.org/10.1371/journal.pbio.0030319>.
- Fay, P.A., Carlisle, J.D., Knapp, A.K., Blair, J.M., Collins, S.L., 2000. Altering rainfall timing and quantity in a mesic grassland ecosystem: design and performance of rainfall manipulation shelters. *Ecosystems* 3, 308–319.
- Fay, P.A., Kaufman, D.M., Nippert, J.B., Carlisle, J.D., Harper, C.W., 2008. Changes in grassland ecosystem function due to extreme rainfall events: implications for responses to climate change. *Global Change Biology* 14, 1600–1608.
- Fay, P.A., Kelley, A.M., Procter, A.C., Hui, D., Jin, V.L., Jackson, R.B., Johnson, H.B., Polley, H.W., 2009. Primary productivity and water balance of grassland vegetation on three soils in a continuous CO₂ gradient: initial results from the Lysimeter CO₂ gradient experiment. *Ecosystems* 12, 699–714.
- Fay, P.A., Blair, J.M., Smith, M.D., Nippert, J.B., Carlisle, J.D., Knapp, A.K., 2011. Relative effects of precipitation variability and warming on tallgrass prairie ecosystem function. *Biogeosciences* 8, 3053–3068.
- Fay, P.A., Jin, V.L., Way, D.A., Potter, K.N., Gill, R.A., Jackson, R.B., Polley, H.W., 2012. Soil-mediated effects of subambient to increased carbon dioxide on grassland productivity. *Nature-Climate Change*. <http://dx.doi.org/10.1038/NCLIMATE1573>.
- Fierer, N., Schimel, J.P., 2002. Effects of drying–rewetting frequency on soil carbon and nitrogen transformations. *Soil Biology and Biochemistry* 34, 777–787.
- Franck, W.M., Hungate, B.A., Chapin, F.S., Field, C.B., 1997. Decomposition of litter produced under elevated CO₂: dependence on plant species and nutrient supply. *Biogeochemistry* 36, 223–237.
- Franzleubbers, K., Weaver, R.W., Juo, A.S.R., Franzleubbers, A.J., 1994. Carbon and nitrogen mineralization from cowpea plant parts decomposing in moist and in repeatedly dried and wetted soil. *Soil Biology and Biochemistry* 26, 1379–1387.
- Gentile, R., Vanlauwe, B., Six, J., 2011. Litter quality impacts short- but not long-term soil carbon dynamics in soil aggregate fractions. *Ecological Applications* 21, 695–703.
- Gill, R.A., Polley, H.W., Johnson, H.B., Anderson, L.J., Maherali, H., Jackson, R.B., 2002. Nonlinear grassland responses to past and future atmospheric CO₂. *Nature* 417, 279–282.
- Gordon, H., Haygarth, P.M., Bardgett, R.D., 2008. Drying and rewetting effects on soil microbial community composition and nutrient leaching. *Soil Biology and Biochemistry* 40, 302–311.
- Gorissen, A., Cotrufo, M.F., 2000. Decomposition of leaf and root tissue of three perennial grass species grown at two levels of atmospheric CO₂ and N supply. *Plant and Soil* 224, 75–84.
- Guo, L.B., Gifford, R.M., 2002. Soil carbon stock and land use change: a meta analysis. *Global Change Biology* 8, 345–360.
- Haney, R.L., Franzleubbers, A.J., 2009. Soil CO₂ evolution: response from arginine additions. *Applied Soil Ecology* 42, 324–327.
- Haney, R.L., Franzleubbers, A.J., Hons, F.M., 2001. A rapid procedure for prediction of N mineralization. *Biology and Fertility of Soils* 33, 100–104.
- Harper, C.W., Blair, J.M., Fay, P.A., Knapp, A.K., Carlisle, J.D., 2005. Increased rainfall variability and reduced rainfall amount decreases soil CO₂ flux in a grassland ecosystem. *Global Change Biology* 11, 322–334.
- Jastrow, J.D., Miller, R.M., Owensby, C.E., 2000. Long-term effects of elevated atmospheric CO₂ on below-ground biomass and transformations to soil organic matter in grassland. *Plant and Soil* 22, 85–97.
- Jin, V.L., Evans, R.D., 2007. Elevated CO₂ increases microbial carbon substrate use and nitrogen cycling in Mojave Desert soils. *Global Change Biology* 13, 452–465.
- Jin, V.L., Schaeffer, S.A., Ziegler, S.E., Evans, R.D., 2011. Soil water availability and microsite mediate fungal and bacterial phospholipid fatty acid biomarker abundances in Mojave Desert soils exposed to elevated atmospheric CO₂. *Journal of Geophysical Research* 116, G02001. <http://dx.doi.org/10.1029/2010JG001564>.
- Jones, M.B., Donnelly, A., 2004. Carbon sequestration in temperate grassland ecosystems and the influence of management, climate and elevated CO₂. *New Phytologist* 164, 423–439.
- Körner, C., 2006. Plant CO₂ responses: an issue of definition, time and resource supply. *New Phytologist* 172, 393–411.
- Kuzyakov, Y., Friedel, J.K., Stahr, K., 2000. Review of mechanisms and quantification of priming effects. *Soil Biology and Biochemistry* 32, 1485–1498.
- Leakey, A.D.B., Ainsworth, E.A., Bernacchi, C.J., Rogers, A., Long, S.P., Ort, D.R., 2009. Elevated CO₂ effects on plant carbon, nitrogen, and water relations: six important lessons from FACE. *Journal of Experimental Botany* 60, 2859–2876.
- Li, X., Meixner, T., Sickman, J.O., Miller, A.E., Schimel, J.P., Melack, J.M., 2006. Decadal-scale dynamics of water, carbon and nitrogen in a California chaparral ecosystem: DAYCENT modeling results. *Biogeochemistry* 77, 217–245.
- Liu, L., King, J.S., Booker, F.L., Giardina, C.P., Allen, H.L., Hu, S., 2009. Enhanced litter input rather than changes in litter chemistry drive soil carbon and nitrogen cycles under elevated CO₂: a microcosm study. *Global Change Biology* 15, 441–453.
- Loladze, I., 2002. Rising atmospheric CO₂ and human nutrition: toward globally imbalanced plant stoichiometry? *Trends in Ecology and Evolution* 17, 457–461.
- Marhan, S., Demin, D., Erbs, M., Kuzyakov, Y., Fangmeier, A., Kandeler, E., 2008. Soil organic matter mineralization and residue decomposition of spring wheat grown under elevated CO₂ atmosphere. *Agriculture, Ecosystems and Environment* 123, 63–68.
- Melillo, J.M., Aber, J.D., Muratore, J.M., 1982. Nitrogen and lignin control of hardwood leaf litter decomposition dynamics. *Ecology* 63, 621–626.
- Mielnick, P.C., Dugas, W.A., 2000. Soil CO₂ flux in a tallgrass prairie. *Soil Biology and Biochemistry* 32, 221–228.

- Mikha, M.M., Rice, C.W., Milliken, G.A., 2005. Carbon and nitrogen mineralization as affected by drying and wetting cycles. *Soil Biology and Biochemistry* 37, 339–347.
- Millard, P., Midwood, A.J., Hunt, J.E., Whitehead, D., Boutton, T.W., 2008. Partitioning soil surface CO₂ efflux into autotrophic and heterotrophic components, using natural gradients in soil δ¹³C in an undisturbed savannah soil. *Soil Biology and Biochemistry* 40, 1575–1582.
- Miller, A.E., Schimel, J.P., Meixner, T., Sickman, J.O., Melack, J.M., 2005. Episodic rewetting enhances carbon and nitrogen release from chaparral soils. *Soil Biology and Biochemistry* 37, 2195–2204.
- Moretto, A.S., Distel, R.A., Didoné, N.G., 2001. Decomposition and nutrient dynamics of leaf litter and roots from palatable and unpalatable grasses in a semi-arid grassland. *Applied Soil Ecology* 18, 31–37.
- Morgan, J.A., Pataki, D.E., Körner, C., et al., 2004. Water relations in grassland and desert ecosystems exposed to elevated atmospheric CO₂. *Oecologia* 140, 11–25.
- Neitsch, S.L., Arnold, J.G., Kiniry, J.R., Srinivasan, R., Williams, J.R., 2011. Soil and Water Assessment Tool: Theoretical Documentation, Version 2009. Texas Water Resources Institute Technical Report No. 406, College Station, TX, pp. 40.
- Niklaus, P.A., Falloon, P., 2006. Estimating soil carbon sequestration under elevated CO₂ by combining carbon isotope labelling with soil carbon cycle modelling. *Global Change Biology* 12, 1–13.
- Norby, R.J., Cotrufo, M.F., Ineson, P., O'Neill, E.G., Canadell, J.G., 2001. Elevated CO₂, litter chemistry, and decomposition: a synthesis. *Oecologia* 127, 153–165.
- O'Neill, E.G., Norby, R.J., 1996. Litter quality and decomposition rates of foliar litter produced under CO₂ enrichment. In: Koch, G.W., Mooney, H.A. (Eds.), *Carbon Dioxide and Terrestrial Ecosystems*. Academic Press, San Diego, CA, pp. 87–103.
- Paul, E.A., Morris, S.J., Böhm, S., 2001. The determination of soil C pool sizes and turnover rates: biophysical fractionation and tracers. In: Lal, R., et al. (Eds.), *Assessment Methods for Soil Carbon*. Lewis Publications, Boca Raton, FL, pp. 193–205.
- Percival, H.J., Parfitt, R.L., Scott, N.A., 2000. Factors controlling soil carbon levels in New Zealand grasslands: is clay content important? *Soil Science Society of America Journal* 64, 1623–1630.
- Polley, H.W., Johnson, H.B., Derner, J.D., 2003. Increasing CO₂ from subambient to superambient concentrations alters species composition and increases above-ground biomass in a C3/C4 grassland. *New Phytologist* 160, 319–327.
- Polley, H.W., Johnson, H.B., Fay, P.A., Sanabria, J., 2008. Initial response of evapotranspiration from tallgrass prairie vegetation to CO₂ at subambient to elevated concentrations. *Functional Ecology* 22, 163–171.
- Polley, H.W., Fay, P.A., Jin, V.L., Combs Jr., G.F., 2011. CO₂ enrichment increases element concentrations in grass mixtures by changing species abundances. *Plant Ecology* 212, 945–957.
- Polley, H.W., Jin, V.L., Fay, P.A., 2012a. CO₂-caused changes in plant species composition rivals the shift in vegetation between mid-grass and tallgrass prairies. *Global Change Biology* 18, 700–710.
- Polley, H.W., Jin, V.L., Fay, P.A., 2012b. Feedback from plant species change amplifies CO₂ enhancement of grassland productivity. *Global Change Biology* 18, 2813–2823.
- Rasmussen, C., Southard, R.J., Horwath, W.R., 2008. Litter type and soil minerals control temperate forest soil carbon response to climate change. *Global Change Biology* 14, 2064–2080.
- Reed, H.E., Blair, J.M., Wall, D.H., Seastedt, T.R., 2009. Impacts of management legacies on litter decomposition in response to reduced precipitation in a tall-grass prairie. *Applied Soil Ecology* 42, 79–85.
- Schaeffer, S.M., Billings, S.A., Evans, R.D., 2007. Laboratory incubations reveal potential responses of soil nitrogen cycling to changes in soil C and N availability in Mojave Desert soils exposed to elevated atmospheric CO₂. *Global Change Biology* 13, 854–865.
- Schimel, J., Balsler, T.C., Wallenstein, M., 2007. Microbial stress-response physiology and its implications for ecosystem function. *Ecology* 88, 1386–1394.
- Soussana, J.-F., Lüscher, A., 2007. Temperate grasslands and global atmospheric change: a review. *Grass and Forage Science* 62, 127–134.
- Taub, D.R., Wang, X., 2008. Why are nitrogen concentrations in plant tissues lower under elevated CO₂? A critical examination of the hypotheses. *Journal of Integrative Plant Biology* 50, 1365–1374.
- Torbert, H.A., Prior, S.A., Rogers, H.H., Wood, C.W., 2000. Review of elevated atmospheric CO₂ effects on agro-ecosystems: residue decomposition processes and soil C storage. *Plant and Soil* 224, 59–73.
- Wan, S., Luo, Y., 2003. Substrate regulation of soil respiration in a tallgrass prairie: results of a clipping and shading experiment. *Global Biogeochemical Cycles* 17, 1054. <http://dx.doi.org/10.1029/2002GB001971>.
- Weltzin, J.F., Loik, M.E., Schwinning, et al., 2003. Assessing the response of terrestrial ecosystems to potential changes in precipitation. *Bioscience* 53, 941–952.
- Williams, M.A., Rice, C.W., Owensby, C.E., 2000. Carbon dynamics and microbial activity in tallgrass prairie exposed to elevated CO₂ for 8 years. *Plant and Soil* 227, 127–137.
- Xiang, S.-R., Doyle, A., Holden, P.A., Schimel, J.P., 2008. Drying and rewetting effects on C and N mineralization and microbial activity in surface and subsurface California grassland soils. *Soil Biology and Biochemistry* 40, 2281–2289.
- Xie, Z., Cadisch, G., Edwards, G., Baggs, E.M., Blum, H., 2005. Carbon dynamics in a temperate grassland soil after 9 years exposure to elevated CO₂ (Swiss FACE). *Soil Biology and Biochemistry* 37, 1387–1395.
- Yuste, J.C., Janssens, I.A., Ceulemans, R., 2005. Calibration and validation of an empirical approach to model soil CO₂ efflux in a deciduous forest. *Biogeochemistry* 73, 209–230.