

2013

Dynamics of Plant Root Growth Under Increased Atmospheric Carbon Dioxide

M. Madhu

Central Soil and Water Conservation Research and Training Institute

J. L. Hatfield

USDA-ARS, jerry.hatfield@ars.usda.gov

Follow this and additional works at: <http://digitalcommons.unl.edu/usdaarsfacpub>

Madhu, M. and Hatfield, J. L., "Dynamics of Plant Root Growth Under Increased Atmospheric Carbon Dioxide" (2013). *Publications from USDA-ARS / UNL Faculty*. 1352.

<http://digitalcommons.unl.edu/usdaarsfacpub/1352>

This Article is brought to you for free and open access by the U.S. Department of Agriculture: Agricultural Research Service, Lincoln, Nebraska at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Publications from USDA-ARS / UNL Faculty by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Dynamics of Plant Root Growth under Increased Atmospheric Carbon Dioxide

M. Madhu and J. L. Hatfield*

ABSTRACT

Plant growth is influenced by above- and belowground environmental conditions and increasing atmospheric carbon dioxide (CO₂) concentrations enhances growth and yield of most agricultural crops. This review covers current knowledge on the impact of increasing CO₂ concentration on root dynamics of plants in terms of growth, root/shoot (R/S) ratio, root biomass, root length, root longevity, root mortality, root distribution, root branching, root quality, and the response of these root parameters to management practices including soil water and nutrients. The effects of CO₂ concentration on R/S ratio are contradictory due to complexity in accurate underground biomass estimation under diverse crops and conditions. Roots become more numerous, longer, thicker, and faster growing in crops exposed to high CO₂ with increased root length in many plant species. Branching and extension of roots under elevated CO₂ may lead to altered root architecture and ability of roots to acquire water and nutrients from the soil profile with exploration of the soil volume. Root turnover is important to the global C budget as well as to nutrient cycling in ecosystems and individual plants. Agricultural management practices have a greater impact on root growth than rising atmospheric CO₂ since management practices influence soil physical, chemical, and biological properties of soil, consequently affects root growth dynamics in the belowground. Less understood are the interactive effects of elevated CO₂ and management practices including drought on root dynamics, fine-root production, and water-nutrient use efficiency, and the contribution of these processes to plant growth in water and nutrients limited environments.

GLOBAL CLIMATE CHANGE has emerged as an important environmental challenge due to its potential impact on biological systems on Earth. Atmospheric concentrations of CO₂ have steadily increased from approximately 315 μmol mol⁻¹ in 1959 to a current atmospheric concentration of approximately 385 μmol mol⁻¹ which converts to an average annual increase rate of nearly 2 μmol mol⁻¹ (Keeling and Whorf, 2005). At this rate of increase, concentrations are projected to reach levels between 500 and 1000 μmol mol⁻¹ by 2100 (IPCC, 2007). Carbon dioxide is not only a major greenhouse gas, but essential to plant growth (Kramer, 1981; Dahlman et al., 1985; Warrick, 1988; Kimball, 2011). The flow of C from photosynthesizing tissues of higher plants, through the roots and into the soil is one of the key processes in terrestrial ecosystems. Increases in atmospheric CO₂ concentration will have direct and indirect effects on crop plants and increases in CO₂ will generally increase plant productivity and water-use efficiency (Drake and González-Meier, 1997; IPCC, 2007). The long-term response to CO₂ remains uncertain and will depend on environmental constraints. Yields of most agricultural crops will increase under elevated CO₂ with productivity increases in the range 15 to 41% for C₃ crops and 5 to 10% for

C₄ crops (Cure, 1985; Kimball, 1983; IPCC, 2007; Lotze-Campen and Schellnhuber, 2009).

Analyses of plant responses to elevated atmospheric CO₂ have focused largely on aboveground processes; however, understanding the effects on photosynthesis are insufficient to answer questions about overall plant response to a changing atmosphere. A whole-plant perspective is required to understand the critical feedbacks and adjustments occurring within a plant and between plants and soil. An overlooked and under studied aspect of plant response to rising CO₂ is on belowground processes. The vital role of roots as an interface between the lithosphere and biosphere is necessary to understand plant response to elevated CO₂. Despite the important role roots play, they have been an understudied component of agricultural research since they exist underground. Root health in crop plants will play a major role in providing sustainable highly productive crops with the ability to cope with climate changes; however, the effect of increasing CO₂ on root growth and development is poorly understood. Climate change is expected to increase the incidence of extreme weather events, such as drought, heat waves, and heavy precipitation and floods, causing crop production to become more variable (IPCC, 2001, 2007; Hatfield et al., 2011). Under these conditions, many of the environmental factors, for example, water, temperature, light, nutrition, salinity, air pollutants, and competition have significant interactions with CO₂ concentration on root responses for numerous species as summarized by Rogers et al. (1994). Understanding of dynamics of crop roots is important from the point of view of management of available resources to increase the productivity of crops and resilience of crops to climate stress.

M. Madhu, Central Soil and Water Conservation Research and Training Institute, Research Centre, Sunabeda, Koraput, Odisha-763 002, India; J.L. Hatfield, Agriculture and the Environment (NLAE), USDA-ARS, Ames, IA 50011. Received 16 Jan. 2013. *Corresponding author (jerry.hatfield@ars.usda.gov).

Published in *Agron. J.* 105:657–669 (2013)

doi:10.2134/agronj2013.0018

Copyright © 2013 by the American Society of Agronomy, 5585 Guilford Road, Madison, WI 53711. All rights reserved. No part of this periodical may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording, or any information storage and retrieval system, without permission in writing from the publisher.

Abbreviations: ABA, abscisic acid; FACE, free air carbon dioxide enrichment; R/S, root/shoot ratio.

The objectives of this review are to express the importance of understanding the dynamic of crop species and to summarize the current knowledge on the effect of increasing CO₂ concentration on root growth, R/S ratio, root biomass, root length, root longevity, root mortality, root distribution, root branching, root quality, and the response of these root parameters to soil management practices, and interactions with soil water and nutrient status.

Root Biomass

Increasing atmospheric CO₂ significantly increased the final plant biomass, aboveground biomass, leaf area and belowground biomass (Obrist and Arnone, 2003). Increased root growth contributes to root biomass and root dry weight under elevated atmospheric CO₂ regardless of species or study conditions (Rogers et al., 1994, 1996). Roots often exhibit the greatest relative dry weight gain among plant organs (Imai and Murata, 1976; Wittwer, 1978; Rogers et al., 1983; Imai et al., 1985; Hocking and Meyer, 1991; Norby et al., 1992), even more than aboveground biomass or leaf area production (Norby, 1994; Rogers et al., 1999; Pritchard and Rogers, 2000; Bernacchi et al., 2000). However, increased allocation of C to plant-root systems in response to CO₂ levels may offset losses from increased activity of soil microorganisms (Suter et al., 2002). Increases in root mass may benefit mycorrhizal fungi and production of glomalin, a glycoprotein produced by endomycorrhizae contributing to soil aggregation (Rillig et al., 1999).

Wheat (*Triticum aestivum* L.) (Chaudhuri et al., 1990; McMaster et al., 1999; Wechsung et al., 1995), sorghum [*Sorghum bicolor* (L.) Moench] (Chaudhuri et al., 1986), Black gram (*Vigna mungo* var. *mungo*; Vanaja et al., 2007), and soybean [*Glycine max* (L.) Merr.] (Del Castillo et al., 1989; Rogers et al., 1992), have shown increases in root dry weight under CO₂ enrichment (Rogers et al., 1994). Laboratory studies on cotton (*Gossypium hirsutum* L.) plants in a free air carbon dioxide enrichment (FACE) project revealed dry weights, lengths, and volumes of taproots, lateral roots, and fine roots were higher for CO₂-enriched cotton plants with a short-term exposure to increased CO₂ for 6 wk (Prior, 1992; Prior et al., 1994a, 1994b; Rogers et al., 1993). Observations from a spring wheat FACE study revealed a 37% increase in total root dry mass in the CO₂-enriched plot during early vegetative growth (stem elongation) and sustained root growth rates until anthesis (Wechsung et al., 1999). The combined effect of temperature and elevated CO₂ concentration affect growth and development of roots, Pilumwong et al. (2007) observed in groundnut (*Arachis hypogaea* L.) that fibrous root dry weight increased with increased CO₂ when plants were grown at 25/15°C, but decreased with increasing CO₂ at 35/25°C. At 35/25°C, there was greater root dry weight by 34% at 400, 14% at 600, and 7% at 800 μmol mol⁻¹ compared with 25/15°C.

Narrow leaf plantain (*Plantago lanceolata*) seedlings grown under elevated CO₂ (800 μmol mol⁻¹) produced 180% greater root dry matter than at 400 μmol mol⁻¹ and amount of plant C recovered was 3.2 times greater from the elevated CO₂ treatment (Hodge and Millard, 1998). The same response was found for raspberry (*Rubus idaeus* L.) plantlets (Laforge et al., 1991). A meta-analysis of the effects of elevated CO₂ on woody plant species (Curtis and Wang, 1998) found biomass responses

were strongly affected by environmental stress factors and to a lesser degree by duration of CO₂ exposure. They reported belowground biomass responses to CO₂ were highly significant overall (+38%, k = 64) but were affected by stress and length of exposure to increased CO₂. About 36 studies reported underground biomass under unstressed trees increased by 41.5% under elevated CO₂. Some plant species namely, kidney vetch (*Anthyllis vulneraria*), hoary plantain (*Plantago media*) (Ferris and Taylor, 1993), spartina with shallow rooting depth (Curtis et al., 1990), and common bean (*Phaseolus vulgaris* L.) (Salsman et al., 1999) exhibited no response of root dry matter production to elevated CO₂.

Root growth of crop plants is often stimulated to a greater extent than other plant parts to increased CO₂ concentration (Rogers et al., 1994; Norby et al., 1992; Kimball et al., 2002; Pritchard and Amthor, 2005; Heinemann et al., 2006; Vanaja et al., 2007). Hodge and Millard (1998) observed belowground growth was stimulated in narrow leaf plantain seedlings with increased CO₂ concentration than aboveground components. In maize (*Zea mays* L.) grown under elevated CO₂ relative growth rate of maize roots was increased compared to the relative shoot growth rate (Whipps, 1985) because of increased C translocation to the roots. Pilumwong et al. (2007) reported increased root growth of groundnut due to increasing CO₂ concentration. With no limitations to water and nutrients, elevated CO₂ increased root and shoot growth of most plant species including tree species (Obrist and Arnone, 2003; Rogers et al., 1996; Pregitzer et al., 1995) but the response varied among species (Hanley et al., 2004). Salsman et al. (1999) found increased CO₂ affected root and shoot growth of *Phaseolus acutifolius* but not *P. vulgaris* due to increased concentration of starch in roots of *P. acutifolius* by 10-fold, while root concentrations of abscisic acid (ABA) doubled and caused more C to be allocated to root growth. Several laboratory studies have reported that roots have the capacity to produce significant amounts of ABA (Walton et al., 1976; Cornish and Zeevaart, 1985; Zhang and Davies, 1987) but applications of ABA have been shown to both increase (Mulkey et al., 1983; Pilet and Saugy, 1987) and decrease root growth (Watts et al., 1981; Jones et al., 1987).

Growth and morphology of four native chalk grassland herbs roots responded differently to increasing CO₂ (Ferris and Taylor, 1993). Root growth and biomass was stimulated by elevated CO₂ for *Sanguisorba minor* and *Lotus corniculatus* with no significant effect on growth and root biomass for *A. vulneraria* or *P. media*. They suggested these growth and root responses could potentially lead to changes in the structure of this plant community with continued increases in CO₂. Increasing CO₂ in greenhouse environments increased the percentage of cuttings which formed roots in a range of ornamental and floricultural species (Lin and Molnar, 1981; French and Alsbury, 1989), including raspberry plantlets (Laforge et al., 1991).

Agricultural management practices may have a greater impact on root growth and soil C storage than rising atmospheric CO₂ (Canadell et al., 1996b; Paustian et al., 1996) because tillage systems influence soil temperature (Dwyer et al., 1995), mechanical resistance (Cox et al., 1990), macropore continuity (Roseberg and McCoy, 1992), and available soil water (Cox et al., 1990) through the effects on rooting depth and root distribution with depth. Rooting depth varies with

season, soil texture, and tillage, and increased rooting depth is associated with increased tillage and decreased soil moisture in surface soil layers (Dwyer et al., 1996). Soil water content in the upper soil layers is confounded by reduced soil water holding capacity and thus plant growth and root development depends on rainfall distribution (Lampurlanés et al., 2001).

At the present time, the interactions of CO₂ enrichment and agricultural management practices on plant root growth are not well understood (Swinnen et al., 1995; van Noordwijk et al., 1994). Although a substantial quantity of data indicates that root spatial distributions are significantly affected by tillage practices, it remains unclear whether temporal or spatial distribution of roots differs among conventional compared with conservation tillage systems. Swinnen et al. (1995) found that decay and turnover in barley (*Hordeum vulgare* L.) roots were higher in conventional agricultural systems than in a production system characterized by reductions in nutrient input, soil tillage, and use of biocides, but were not different in wheat under the same management scenarios. In another study, sugar beet (*Beta vulgaris* L.) grown with minimum tillage had lower fine-root production than in the conventional cropping system, whereas fine-root production of wheat was unchanged (van Noordwijk et al., 1994).

Prior et al. (2005) studied the effects of management and CO₂ on aboveground biomass production with an emphasis on changes in soil C storage. They found a substantial increase in soil C concentration at the 0- to 5-cm depth increment in the conservation system compared to the conventional system, and in the CO₂-enriched compared to ambient plots, after two cropping cycles (4 yr). Pritchard et al. (2006) concluded that CO₂ enrichment enhanced cumulative seasonal root length production (+58%) and mortality (+59%) in conventionally managed sorghum, but had no effect in plots with conservation systems. Sorghum under CO₂ enrichment produced more roots at shallow soil depths compared to ambient plots and this additional root production occurred early in the growing season. Conventionally managed sorghum produced less root length in the top few centimeters of soil compared to conservation plots due to increased soil water availability in the conservation tillage plots. Seasonal root length mortality and production were higher at the shallower soil depths in the conservation plots than in the conventional plots. Increases in root mortality in CO₂-enriched plots were proportional to production increases and no effect on root longevity which associated with depth in the soil profile (Pritchard et al., 2006). This was probably caused by the effects of tillage on soil water content due the presence of surface residues coupled with no-till management enhance soil moisture absorption and retention compared to conventional tillage with no residue (Newell and Wilhelm, 1987).

A particular limitation to understanding root dynamics is that most studies do not include measurements below a meter (Nepstad et al., 1994; Batjes, 1996; Canadell et al., 1996b; Jobbagy and Jackson, 2000; Guo and Gifford, 2002; Schenk and Jackson, 2002a, 2002b; Robinson, 2004; Bradley et al., 2005; Lorenz and Lal, 2005; Mokany et al., 2006; Qin and Huang, 2010; Wang et al., 2010). Baker et al. (2007) argued depth of root sampling confounded understanding the impact of tillage and management factors on root growth, root turnover, distribution, etc. and sequestration of C at different depths in the soil.

Roots of drought-stressed plants grown with elevated CO₂ produced significantly higher root dry mass than roots of well-watered plants grown with ambient CO₂ (Chaudhuri et al., 1990). These responses in root growth and morphology led to elevated CO₂ and reduced soil water supports the hypothesis that plants grown in a high CO₂ environment may better compensate to soil-water-stress conditions (Wechsung et al., 1999; Schenk and Jackson, 2002b). An adoption mechanism of plants to water stress condition occurs by physiological and biochemical changes in the plant. Eggplant (*Solanum melongena* L.) roots showed low hydraulic conductance at repetitive stress due to synthesis and accumulation of suberin in the root cells under elevated CO₂ and this change in root structure may occur from synthesizing suberin lipids in endo- and exo-dermis layers regardless of soil water availability or stress intensity. Lowering conductance is highly desirable for agricultural plants when water supply is limited in the growth and development stage (Sarker and Hara, 2009). These observations suggest increasing CO₂ concentration in the global atmosphere might be beneficial for using soil water by minimizing transpiration loss and lowering hydraulic conductance under higher CO₂. In a comprehensive review of FACE experiments on agricultural crops, Kimball (2011) and Kimball et al. (2002) determined that for a 300 μmol mol⁻¹ increase in atmospheric CO₂ concentration, the root biomass of wheat, ryegrass (*Lolium multiflorum* Lam.), and rice (*Oryza sativa* L.) increased 70% with ample water and N, 58% at low N, and 34% at low water, while clover experienced a 38% increase at ample water and N, plus a 32% increase at low N. The changes were greatest in cotton with a 96% increase in root biomass at ample water and N. Another FACE study on cotton and sorghum in monoculture and two plant densities under dry and wet soil water condition was conducted by Derner et al. (2003) and showed total root biomass in the upper 1.2 m of the soil was not influenced by CO₂ or soil moisture in monoculture or in mixture; however, under dry conditions there were significantly more roots at soil depths below 45 cm. Sorghum roots comprised 81 to 85% of the total roots in the low density mixture and 58 to 73% in the high density mixture. Carbon dioxide enrichment partly offsets the negative effects of interspecific competition on cotton in both low and high density mixture by increasing aboveground biomass (Derner et al., 2003). The interacting effects of CO₂ and watering frequency on wheat root growth and nutrient uptake pattern increased water use per plant 1 to 25 times at 350 than at 700 μmol mol⁻¹ CO₂ and 1 to 4 times in the “wet” compared to the “dry” treatment (Van Vuuren et al., 1997). Root biomass increased with CO₂ and with watering frequency. Nutrient uptake was affected and there were interactions with water availability but interactions were different for N, K, and P, due to differences in nutrient availability and mobility in the soil (Van Vuuren et al., 1997).

Root growth is positively affected by increases in CO₂ concentration across almost all species and is often stimulated more than shoot biomass. There is no simple assessment of the interacting effects of soil physical conditions and nutrient availability and the interaction with increasing CO₂ on root and shoot growth. A summary of the combined effects of increased CO₂ for the different root parameters is shown in Table 1.

Table 1. Overall effects of increased CO₂ on various root parameters.

Root parameter	Species	Type of CO ₂ enrichment	Overall effect
Root biomass/growth	Agricultural plants, groundnut, grassland herbs	Laboratory studies, greenhouse studies	Increase in root growth was larger than shoot growth and root biomass increased as part of overall growth response to increased CO ₂ .
Root/shoot (R/S) ratio	Agricultural plants, groundnut, grassland herbs, forest species	Laboratory, greenhouse, controlled chamber studies	Results vary among studies; however, 60% of studies show a positive response on R/S, 37% show a negative response, and 3% show no response (Rogers et al., 1996). Under limited soil water or nutrient availability, responses to increased CO ₂ are more variable.
Root length/rooting depth	Agricultural plants, groundnut, grassland herbs, forest species	Laboratory, greenhouse, controlled chamber studies	Increased root length across species is the most common response observed. Root branching also increased with CO ₂ enrichment (Rogers et al., 1994).
Root branching	Agricultural plants, groundnut, grassland herbs, horticultural species	Laboratory, greenhouse, controlled chamber studies, free air carbon dioxide enrichment (FACE) studies	Increased root branching is commonly observed in most species and more evident in the upper soil profile.
Root thickness, quality, diameter	Agricultural plants, groundnut, grassland herbs, horticultural species, forests	Laboratory, greenhouse, controlled chamber studies	Increased root thickness and diameter are most commonly observed. Diameter changes are more difficult to quantify.
Root mortality/root turnover	Agricultural plants, groundnut, grassland herbs, horticultural species, forests	Laboratory, greenhouse, controlled chamber studies, FACE studies	Increases in root mortality and turnover rates are observed across species with increased CO ₂ . There is an interaction with temperature, soil moisture, and nutrient supply which confounds observations on root turnover rates.

Root/Shoot Ratio

Exposure of plant canopies to high CO₂ concentration often stimulates growth of shoots and roots, but the question remains whether elevated atmospheric CO₂ concentration affects root and shoot growth proportionally (Rogers et al., 1996). The general consensus is that photosynthesis and C allocation to plant roots increases as atmospheric CO₂ rises which leads to an increase in above and below biomass (Del Castillo et al., 1989; Norby et al., 1992; Rogers et al., 1994). Root/shoot ratio is the simple calculation of the ratio of root dry mass to shoot (or stem) dry mass and should serve as a measure of the preferential allocation of C to roots or shoots. Allen et al. (1988) reported that soybean grown under increased CO₂ maintained a similar partitioning of C into their respective components. Similar results were reported by Baxter et al. (1994) in grasses, but roots often exhibit the greatest relative dry weight gain under high CO₂ (Hocking and Meyer, 1991; Imai and Murata, 1976; Imai et al., 1985; Norby et al., 1992; Rogers et al., 1983). The partitioning pattern of photosynthates depends on plant development stage, plant species, and plant growth conditions along with physiological factors (Van Veen et al., 1991). Various studies have shown between 10 and 40% of the total net C assimilation of crops was transferred to underground biomass in the form of roots, microbial and soil biomass, and respired CO₂ and organic compounds (Barber and Martin, 1976; Whipps, 1984; Helal and Sauerbeck, 1986; Liljeroth et al., 1990; Keith et al., 1986).

Observations on R/S response to CO₂ are contradictory as evidenced by the wide range of results obtained from different studies. Oechel and Strain (1985) found root growth under increasing atmospheric CO₂, plants allocated proportionately more C belowground, increasing R/S ratios. The range of response in R/S among crop plants to CO₂ doubling ranged from a 8.5% decrease to a 6.4% increase, except in sweet potato [*Ipomoea batatas* (L.) Lam. var. batatas], in which a 34.9% increase was observed (Cure, 1985). Results showed R/S was

not significantly affected by higher CO₂ concentration in *Larrea tridentate*, a desert herb (Obrist and Arnone, 2003), tall grasses (Mo et al., 1992), four native chalk grassland herbs (Ferris and Taylor, 1993), and winter wheat (Chaudhuri et al., 1990; Havelka et al., 1984). Many studies showed increased R/S (Rogers et al., 1983; Biswas and Hileman, 1985; Idso et al., 1988; Baker et al., 1990; Rogers et al., 1992; Ceulemans and Mousseau, 1994; Pritchard et al., 1999; Rogers et al., 1999), while other studies showed decreased R/S (Salsman et al., 1999; McMaster et al., 1999). In field studies, an increase in the proportion of total dry mass in tubers after exposure to increased CO₂ was observed for sweet potato (Biswas and Hileman, 1985; Bhattacharya et al., 1990), carrot (*Daucus carota* L.), and radish (*Raphanus sativus* L.) (Idso et al., 1988). Idso et al. (1988) suggested that plants whose primary yield component is produced aboveground, atmospheric CO₂ enrichment has a significant effect on R/S. Increases in R/S occurred more frequently for agronomic crops, particularly root and tuber crops and natural community species. Substantial CO₂ research conducted with several forest tree species namely, shortleaf pine (*Pinus echinata* Mill.) (Norby et al., 1987; O'Neill et al., 1987b), virginia pine (*P. virginiana* Mill.) (Luxmoore et al., 1986), white oak (*Quercus alba* L.) (Norby et al. (1986a, 1986b; O'Neill et al., 1987b; Norby and O'Neill, 1989), yellow poplar (*Liriodendron tulipifera* L.) (O'Neill et al., 1987a; Norby et al., 1992), and N-fixing woody plants (Norby, 1987) have revealed an increased R/S, C allocation to roots, root exudation, nutrient uptake, and mycorrhizal colonization under elevated CO₂.

Rogers et al. (1996) summarized a substantial variation in R/S response in crop plants across a range of experimental conditions. He reported crops showed positive responses in R/S to elevated CO₂, 59.5% of 264 observations from 62 reports; negative responses appeared 37.5% of the studies and 3.0% of the studies showed no response (Table 1). There is no general consensus among researchers on the degree of positive response on R/S of

different species; however, these results are tempered by the interactions with water and nutrient management in the soil profile.

Under non-limiting conditions of water and nutrients for growth, dry matter partitioning to the root is moderately changed by CO₂ enrichment. However, increases in R/S, frequently observed under limiting conditions of water and/or nutrients, allows the plant to explore a greater soil volume and extract more water and nutrients (Stulen and den Hertog, 1993). Bazzaz (1990) stated that there would be an increase in C allocation to roots when nutrients and water were limiting. Biomass allocation models such as functional balance developed by Davidson (1969) and C/nutrient substrate ratio developed by Reynolds and Thornley (1982) predicted increases in tissue C/N ratios as N becomes more limiting relative to C leading to increased biomass allocation to roots. Eamus and Jarvis (1989) summarized that under low nutrient conditions there would be an increase in R/S, and this conclusion was confirmed by Ceulemans and Mousseau (1994). There are confounding effects of increasing CO₂ on root allocation and developmental rate and Norby (1994) observed a small (+6%) increase in R/S at high CO₂, with no significant response under nutrient stress. McGuire et al. (1995) and Wullschleger et al. (1995) did not find a significant CO₂ effect on biomass allocation between roots and shoots. The interactions between nutrient status and CO₂ are not clearly defined because the conclusions by Norby et al. (1987) were that limiting nutrients may promote partitioning to the roots and assimilate partitioning under high CO₂ may depend on nutrient status (Lawlor and Mitchell, 1991) and requires that we quantify the nutrient levels in the soil when interpreting R/S responses.

The direct effect of increasing CO₂ on R/S has no clearly defined conclusion (Stulen and den Hertog, 1993; Rogers et al., 1994). The underlying assumption that a larger proportion of dry matter produced under CO₂ enrichment is preferentially allocated to roots and increased under limiting water and nutrient supply needs to be evaluated. Differences among experiments can be attributed to crop type, genotypes, resource supply (water and nutrients), and other experimental factors, that is, duration and method of exposure, and interacting treatment variables (Idso et al., 1988; Rogers et al., 1996). This is further complicated by poor estimate of C returned to the soil via the death of fine roots at harvest leading to an underestimation of C allocation (Stulen and den Hertog, 1993). Recent observations of in situ root dynamics from nondestructive monitoring methods reveals estimates of net belowground production significantly underestimates gross production (Fahey and Hughes, 1994; Hendrick and Pregitzer, 1992, 1993a, 1993b). Berntson and Bazzaz (1997a) argued that elevated CO₂ increases the magnitude of cumulative gross production and cumulative gross root loss with a large seasonally dependence on these responses. Elevated CO₂ enhances gross production of roots early in the growing season, before maximum root length and enhance cumulative gross loss of roots in the latter portion of the growing season after maximum root length had been achieved. They concluded that static assessments of belowground productivity may underestimate gross fine root productivity and turnover, and this bias may be exaggerated with elevated CO₂. Stulen and den Hertog (1993) list several reasons why determination of R/S is subject to experimental error; including problems in defining the R/S boundary, quantitative

recovery of roots (especially fine roots), root decomposition, nutrient and water regime in the soil, ontogenetic effects, and accumulation of non-structural carbohydrates.

Root Length and Rooting Depth

When exposed to increased CO₂, roots have been observed to become more numerous, longer, thicker, and faster growing in crops (Chaudhuri et al., 1990; Prior et al., 1994a, 1994b, 1995) with increased root length in many plant species (Norby, 1994; Rogers et al., 1999; Pritchard and Rogers, 2000; Bernacchi et al., 2000). Root distributions with depth of sorghum with elevated CO₂ was studied by Pritchard et al. (2006) and they concluded the greatest root length was observed between 4- and 15-cm soil depths. At physiological maturity, a larger percentage of total root length had died at shallow soil depths compared to deeper depths; for example, 87% of roots produced at the shallow soil depths died before harvest compared to 48% at the lower depths. Daily production of new roots occurred primarily in the early part of the growing season and the maximum root length production per day occurred when mortality and production rates were nearly equivalent at approximately 59 d after planting.

Lengths and volumes of taproots, lateral roots and fine roots were higher for CO₂-enriched cotton plants (Prior, 1992; Prior et al., 1994a, 1994b; Rogers et al., 1993). A 110% increase in root length of soybean was observed as CO₂ concentration increased from 350 to 700 μmol mol⁻¹ (Rogers et al., 1992). Increased root length and number was found in sweet potato (Bhattacharya et al., 1990), and in *P. acutifolius* and *P. vulgaris* (Salsman et al., 1999). Increased stele and cortex diameter, root diameter in the root hair zone, length of unbranched first order lateral roots, total root length, and root volume was increased in cotton with increased CO₂ (Rogers et al., 1992).

Temperature and CO₂ effects were studied in groundnut and total root length increased with increasing CO₂ from 400 to 800 μmol mol⁻¹ at both low and high temperatures (Pilmwong et al., 2007). Although the difference was not significant, there was an observed increase of 13% when plants were grown at 600 μmol mol⁻¹ CO₂, and 20% at 800 μmol mol⁻¹ CO₂. However, visible root length, root number, and root length density increased with CO₂ increases from 400 to 800 μmol mol⁻¹ at 25/15°C, but decreased as CO₂ increased when temperatures increased to 35/25°C (maximum/minimum temperature). Root branching was the primary factor affecting total length and volume of roots. Berntson and Woodward (1992) concluded CO₂ enrichment resulted in more branched and longer root system of *Senecio vulgaris*. Many studies have found elevated CO₂ resulted in more and/or longer plant roots or faster root growth, possibly leading to increased penetration of the soil profile (Baker et al., 1990; Chaudhuri et al., 1990; Rogers et al., 1992) and/or horizontal spread of the roots (Idso and Kimball, 1991)

Root length and number have been the most frequently evaluated in horticultural species (45 and 74% of studies, respectively) compared with plants in other categories (2–12%). A majority (61–100%) of studies found that increased CO₂ resulted in more and/or longer plant roots (Rogers et al., 1994) which led to increased depth (Baker et al., 1990; Rogers et al., 1992) and/or horizontal spread (Idso and Kimball, 1991, 1992) (Table 1).

Rooting depth has important implications for hydrology, biogeochemistry, and primary productivity of terrestrial

ecosystems (Jackson et al., 1997) and since the depth of plant roots depends on a variety of conditions, including soil properties, vegetation type, nutrient availability, and climate there are a number of potential implications from rooting depth response to CO₂ (Guswa, 2010). A number of papers and reviews describe the genetic control of root architecture and the variation between plant types and individual plant cultivars about the maximum rooting depth (Kell, 2011). Plant root depths vary greatly in similar soils for different organisms (e.g., Burch and Johns, 1978; Jackson et al., 1996; Jobbagy and Jackson, 2000), and varies substantially within the same soils or growth media for different cultivars of the same plant, and between different cultivars of the same parent.

Canadell et al. (1996a) reviewed rooting depth from 290 observations of maximum rooting depth for 253 woody and herbaceous species and found maximum rooting depth ranged from 0.3 m for some tundra species to 68 m for *Boscia albitrunca* in the central Kalahari. In this analysis, 194 species had roots at least 2-m deep, 50 species had roots with a depth of 5 m or more and 22 species with rooting depth to 10 m or deeper (Canadell et al., 1996a). The average rooting depth from this comparison was 4.6 ± 0.5 m. When the maximum rooting depth was segregated by biome the results showed 2.0 ± 0.3 m for boreal forest, 2.1 ± 0.2 m for cropland, 9.5 ± 2.4 m for desert, 5.2 ± 0.8 m for sclerophyllous shrub land and forest, 3.9 ± 0.4 m for temperate coniferous forest, 2.9 ± 0.2 m for temperate deciduous forest, 2.6 ± 0.2 m for temperate grassland, 3.7 ± 0.5 m for tropical deciduous forest, 7.3 ± 2.8 m for tropical evergreen forest, 15.0 ± 5.4 m for tropical grassland/savanna, and 0.5 ± 0.1 m for tundra. If the aggregation was conducted across species within biomes (except croplands) by three basic functional groups maximum rooting depth was 7.0 ± 1.2 m for trees, 5.1 ± 0.8 m for shrubs, and 2.6 ± 0.1 m for herbaceous plants. These analyses data show that deep root habits are quite common in woody and herbaceous species across terrestrial biomes, deeper than previously assumed. These results argue for an enhanced understanding of ecosystem function and its application in developing ecosystem models.

Carbon dioxide enrichment resulted in enhanced root proliferation at shallower soil depths in wheat (Chaudhuri et al., 1990; Fitter et al., 1996; Van Vuuren et al., 1997; Van Vuuren et al., 1997), cotton (Rogers et al., 1992), sorghum (Chaudhuri et al., 1986), and grassland (Fitter et al., 1997). Chaudhuri et al. (1990) found winter wheat grown under elevated CO₂ achieved maximum rooting penetration significantly faster than plants grown in ambient air. They also found that root growth differences between ambient and elevated CO₂ occurred in the upper 10 cm of soil and concluded that high levels of CO₂ compensate for restriction in growth of wheat roots by drought. In contrast to wheat, Chaudhuri et al. (1986) found root numbers and dry weights of sorghum roots were higher at all depths to 150 cm under increased CO₂. Controlled studies on rooting pattern of cotton plants under FACE (Prior, 1992; Prior et al., 1994a, 2003; Rogers et al., 1993) demonstrated that root architecture, defined as the distribution of fine root density per unit volume of soil and expressed as length or dry weight per m³, increased both vertically and horizontally with increased CO₂. Fine root density increased under

CO₂ enrichment to 90 cm with the most significant increase in the upper 45 cm. Root length and dry weight densities exhibited greater difference between ambient and elevated CO₂ as distance from row center increased, indicating faster and more prolific spread of cotton roots (Rogers et al., 1994).

Experimental evidence from forested ecosystems shows CO₂ enrichment may lead to deeper rooting distributions (Rogers et al., 1994). Altered rooting distributions will affect ecosystem processes and the depth to which fine roots are produced influences root chemistry, physiological function, and mycorrhizal infection, leading to altered N uptake rates and slower root turnover. Changes in rooting patterns influence nutrient dynamics, thus influencing crop performance when nutrient demand is high. Whole plant nutrient uptake is often higher, while tissue nutrient concentration is reduced for CO₂-enriched plants (Prior et al., 1998, 2003; Rogers et al., 1994, 1997). Shifts in rooting patterns alter their competitive effectiveness for edaphic resources. Root proliferation into nutrient-rich zones can be an important mechanism in the exploitation of soil resources (Borkert and Barber, 1985; Jackson and Caldwell, 1989).

Root Branching

The branching patterns of root systems have been described as “developmentally” (Barley, 1970; Rose, 1983) and “topologically” (Fitter, 1985, 1986, 1987). Previous studies show the primary response of crop plants to elevated CO₂ on branching and extension of first-order and higher-order laterals root over the extension of the root system deeper into the soil (Rogers et al., 1999; Pritchard and Rogers, 2000). They suggested that, with higher atmospheric CO₂ concentrations, crop roots will be larger, more highly branched (especially at shallow depths), but less efficient in nutrient and water uptake (decreased specific root activity). This hypothesis is supported from observations by Berntson and Bazzaz (1995), demonstrating the potential of root systems to acquire resources may increase because of greater total root length densities; however, there may be a decline in the efficiency of resource capture. Root systems of crops grown in CO₂-enriched environments are often more branched, especially at shallower soil depths, compared to roots of crops grown in ambient air (sorghum, Chaudhuri et al., 1986; wheat, Chaudhuri et al., 1990; Fitter et al., 1996; *Ceratonia siliqua*, Cruz et al., 1997; soybean, Del Castillo et al., 1989; cotton, Rogers et al., 1992; Pilumwong et al., 2007; *Senecio vulgaris*, Berntson and Woodward, 1992). These observations indicate altered belowground plant function suggesting higher level soil ecological processes may change as atmospheric CO₂ continue to increase (Pritchard and Rogers, 2000; Rogers et al., 1997, 1999).

In soybean, Del Castillo et al. (1989) reported no effect on rates of elongation; however, there was an increase in the number of more highly branched roots. Cruz et al. (1997) reported that carob (*Ceratonia siliqua* L.) seedlings grown in elevated CO₂ produced more lateral roots and shorter and thicker roots. In a FACE study with cotton, Rogers et al. (1992) showed that the number, length, and mass of lateral roots were enhanced by CO₂ enrichment. Although taproot diameter, mass, and volume were enhanced, length was not affected. In contrast, Rogers et al. (1992) reported an increase in soybean root length but no change in the number of lateral roots. Under FACE studies, the lengths and volumes of taproots, lateral roots, and fine roots were higher

for CO₂-enriched cotton plants but the number of lateral roots per unit length of taproot was not significantly increased, the greater taproot lengths with CO₂ enrichment increased total number of laterals (Prior, 1992; Prior et al., 1994a, 1994b; Rogers et al., 1993). Another FACE study on spring wheat by Wechsung et al. (1999) reported a 37% increase in total root dry mass in the FACE vs. control plots during stem elongation. During early vegetative growth, root dry mass from the inter-row space was significantly higher for FACE compared to control treatments suggesting that elevated CO₂ promoted the production of first-order lateral roots per main axis. During the reproductive period of growth, more branching of lateral roots was found in the FACE treatment along with significantly higher root dry mass when soil water supply was limiting. These responses in root growth and morphology to elevated CO₂ and reduced soil water supports the hypothesis that plants grown in a high CO₂ environment may compensate for soil water stress because of the increased root density ability to extract more water per unit soil volume.

Differential effects of CO₂ concentration on root branching may lead to altered root architecture and ability of roots to extract water and nutrients. For example, Berntson and Woodward (1992) observed for *Senecio vulgaris* grown under elevated CO₂ more horizontal branching angles of roots leading to greater horizontal root spread. Rogers et al. (1992) observed longer second-order laterals in soybean, which could lead to deeper root penetration and greater exploration of soil for nutrients and water but could ultimately lead to competition among roots in resource-limited environments. Conversely, Del Castillo et al. (1989) found more roots attributed to increased branching rather than longer roots in soybean grown under enriched CO₂. The positive effect of increased root numbers leads to more efficient exploration of the same volume of soil compared to deeper root penetration (Pritchard et al., 1999). Increasing CO₂ caused increased root branching across species suggesting that greater root exploration of the profile would occur as a result of the positive effect of CO₂ on root growth (Table 1).

Root Thickness, Quality, and Diameter

The additional root biomass production from elevated CO₂ induced increased thickness or diameter or quality (root length density) of plant roots and increased total root length and volume along with root diameter in the root hair zone and length of unbranched first-order laterals (Rogers et al., 1992). Roots become more numerous, longer, thicker, and faster growing in crops grown under high CO₂ (Chaudhuri et al., 1990; Prior et al., 1994a, 1994b, 1995) and the diameters of dead roots did not differ from the diameter of roots that remained alive during the crop growth period of sorghum (Pritchard et al., 2006). Changes in root dry weight density caused by CO₂ enrichment suggests possible changes in root quality. The results of a FACE experiment on cotton by Prior et al. (1994a, 1994b) showed large shifts in root dry weight density. These results suggest possible changes in root quality from changes in root dry weight density through changes in tissue density of roots (fine and taproots) resulting in shifts in carbohydrate storage, cell number, cell size, and percentage of intercellular space or other structural modifications. Soybean plants grown under CO₂ enrichment exhibited a 27% increase in root diameter, stele diameter (23%), and cortex width (28%) in the root hair zone (Rogers et al., 1992). Studies on

structural aspects of plant roots (i.e., diameter, volume, branching, relative growth rate, etc.) have usually shown positive effects of high CO₂. Tubers (number, dry weight, diameter, etc.) and nodulation (number, dry weight, activity, etc.) benefit from elevated CO₂ in most crops (Rogers et al., 1994).

Zhao et al. (2000) germinated pea (*Pisum sativum* L.) seeds and exposed the young plants to various atmospheric CO₂ concentrations in controlled environment chambers to determine if elevated CO₂ impacted root border cells, which are major contributors of root exudates in most agronomic plants. They found elevated CO₂ increased the production of root border cells in pea seedlings. Changing the ambient air to air enriched with 3000 and 6000 μmol mol⁻¹ CO₂, border-cell numbers increased by more than 50 and 100%, respectively. It is likely that continued increases of CO₂ will likely produce greater numbers of root border cells in crop plants and lead to increased amounts of root exudates into the rhizosphere, suggesting that associated soil microbial and fungal activities would be stimulated from increases in plant-derived C inputs. These changes should make the soil environment more favorable for plant growth and development in a high CO₂ environment. Few studies have addressed CO₂-induced changes in root quality/thickness/diameter and structure and further research is needed, especially under field environments (Table 1).

Root Mortality/Root Turnover

Root turnover is important to the global C budget as well as to nutrient cycling in ecosystems and to the success of individual plants in obtaining water and nutrients. Replacement of older roots with newly formed roots is referred to as root turnover. Root turnover is estimated from the difference between cumulative births and deaths, which represents either a net accumulation or disappearance of roots. If roots are not accumulating even though production is enhanced, then turnover must be equally enhanced (i.e., birth rate equals death rate). The difference in accumulation rate between elevated and ambient CO₂ levels can be used to determine whether turnover is affected by CO₂ and a positive difference means that roots are accumulating faster under elevated CO₂. There are several papers discussing approaches for estimating annual root turnover (Milchunas et al., 1985; Caldwell and Eissenstat, 1987; Hendrick and Pregitzer, 1992; Milchunas and Lauenroth, 1992; Hendricks et al., 1993; Fahey and Hughes, 1994), which reveal no consensus on the best approach for estimating root turnover. Del Castillo et al. (1989) observed a significant linear increase in the number of actively growing roots of soybean with increased CO₂ compared to elongation rate. Eissenstat et al. (2000) applied an efficiency model to hypothesize that rising CO₂ would increase average root lifespan. In contrast, other investigators have suggested root lifespan is unlikely to change with increasing CO₂ (Pritchard and Rogers, 2000; van Noordwijk et al., 1998).

Increased photosynthesis under elevated CO₂ stimulates belowground C input, fine-root growth (Curtis, 1996; Curtis and Wang, 1998; Pendall et al., 2005) and root turnover rates and biomass (Berntson and Bazzaz, 1997b; Fitter et al., 1999; Allen et al., 2000; Pregitzer et al., 2000; Wan et al., 2004). Increased fine-root production and mortality are associated with higher temperatures (Gill and Jackson, 2000) and turnover rates (Hendrick and Pregitzer, 1993b, 1997; Forbes et al.,

1997; Fitter et al., 1999; King et al., 1999; Wan et al., 2004). Turnover rates increase with increased C allocation to root systems when plants are grown under elevated CO₂, at least in grasslands and forest (Fitter et al., 1996). Root life span is highly variable, from a few weeks in some plants to a few years in others (Eissenstat and Yanai, 1997). Root longevity varies widely within an individual species.

Fine roots are essential for plant water and nutrient uptake, soil C input, and soil microbial activity (Eissenstat, 1992; Norby, 1994). Turnover of fine roots (<2.0-mm diam.) play a critical role in regulating ecosystem C balance and accurate estimates of belowground net primary productivity (NPP) are required for estimating net ecosystem productivity (NEP) (Pendall et al., 2005). As much as 33% of global annual NPP has been estimated for the production of fine roots (Jackson et al., 1997). Caldwell (1977) suggested for perennial systems, at least 25% of root systems die and replaced annually, accounting for as much as 50 to 80% of net production. Death of older roots and the growth of new ones are a significant energetic cost to the plant (Canadell et al., 1996b). Understanding the trade-offs between the cost of maintaining old roots compared with that of growing new ones is an important step in understanding how increased C assimilation, associated with rising global CO₂ affects root turnover. van Noordwijk et al. (1998) developed a quantitative model for this system by estimating for a root with a growth respiration of 2 g CH₂O g⁻¹ root dry weight and a maintenance respiration of 0.03 g Cg⁻¹ root dry weight, approximately 60 d of root maintenance in unfavorable conditions would cost as much as one cycle of root death followed by regrowth. van der Werf et al. (1988) showed, for a different species, that the adenosine-5'-triphosphate (ATP) costs of producing 1 g fresh wt. of root is approximately equal to the cost of maintaining 1 g of root for 10 d. The implications of this energy balance for roots suggest it may be more energetically expensive to maintain one root when the costs of maintenance during periods of stress exceed the energy requirements of growing new roots on the return of favorable soil conditions.

The availability of underground resources (water and nutrients) influences the C allocation to roots and turnover rates. If we follow the argument of Stulen and den Hertog (1993) that plants grown under non-limiting conditions do not alter their allocation of C between roots and shoots under elevated atmospheric CO₂ then the hypothesis of van Noordwijk et al. (1998) that increasing atmospheric CO₂ is not likely to have a significant direct effect caused by a greater C supply on root turnover. However, indirect effects, mediated by shifts in plant water availability and nutrient relations, are likely to affect root turnover. Experimental evidence shows that root turnover increases more often than not when soil conditions are limiting root uptake (Pregitzer et al., 1995; Canadell et al., 1996b). Canadell et al. (1996b) found from four experiments with grass systems there was increased root turnover in three with no change in one experiment. In four experiments on trees, increased turnover was observed in three, while one showed a decrease (Canadell et al., 1996b). Finally, no difference in root turnover was observed in wheat grown in elevated CO₂ (Fitter et al., 1996). It is likely that root turnover will be different between annual and perennial plants.

Temporal pattern of root growth is consistent with C allocation to roots being highest during early vegetative growth

and decreasing as plants mature and transition to reproductive growth (Swinnen et al., 1994). Pritchard et al. (2006) concluded daily production of new roots in sorghum occurred primarily in the early part of the growing season and daily mortality of roots increased steadily over the growing season in a linear fashion under elevated CO₂. The death of roots and growth of new roots corresponds to the availability of soil water and nutrients (Huck et al., 1987; Smucker, 1993). Huck et al. (1987) found that birth and death rates of roots were mediated in time by rainfall events and in space by moist soil microsites. They observed that the number of active soybean roots decreased after each rain, increased rapidly during dry periods, and declined after subsequent rainfall events. They observed spatial peaks in root growth during dry conditions occurred in the soil profile where soil moisture was favorable (Huck et al., 1987). Resource availability (nutrients, moisture, and temperature) in the top layer of soil plays a key role on the rate of fine root mortality. Pritchard et al. (1999) found a negative linear relationship between soil depth and the proportion of roots that died in sorghum with a similar conclusion made on alfalfa roots (Goins and Russelle, 1996; Watson et al., 2000), groundnut (Krauss and Deacon, 1994), and sugar beet (van Noordwijk et al., 1994). Changes in nutrient availability, temperature, and moisture throughout a day with depth a possible explanation for these observations because of the greater change in the upper soil layers compared to deeper layers (Watson et al., 2000). In tree species, rates of fine root mortality was roughly twice as in trees grown at elevated CO₂ compared to ambient CO₂ in both high and low soil N (Pregitzer et al., 1995).

The projected increases in air temperatures, changing precipitation patterns, and potential decreases in soil nutrients availability could be expected to have impacts on root mortality (Rastetter et al., 1997). Root mortality might be accelerated by decreases in soil moisture (Klepper et al., 1973; Hayes and Seastedt, 1987; Huck et al., 1987) or by increases in soil temperature (Fitter et al., 1999). However, root mortality might be reduced by decreased availability of nutrients (Eissenstat and Yanai, 1997). Fitter et al. (1997) observed in peat and limestone grassland soils, cumulative root birth and death totals were greater under elevated CO₂. In the peat soil, cumulative births were 20% greater (36 as compared to 30 per tube) and deaths 21% greater (34 and 28 per tube) after two seasons under elevated CO₂. In the limestone soil, cumulative births were increased by 30% (35 vs. 27) and deaths by 29% (31 vs. 24). Turnover was faster under elevated CO₂ in the peat soil, but there was only a small effect on turnover in the limestone soil (Fitter et al., 1997).

Finally, it's suggested that more long-term studies are needed, to detect both existing patterns of root mortality and lifespan with resource availability and plant strategy, and to progress beyond short-term studies, which may be misleading. Factors that should be included in studies of root lifespan, in addition to resource availability, include root age, plant C status, and pathogen pressure. The effects of increasing CO₂ on root mortality are less clear and the linkage to changes in temperature and soil water has been difficult to quantify.

Conclusions and Recommendations

Crops grown under elevated atmospheric CO₂ exhibit changes in above- and belowground growth responses which

could potentially lead to alteration in plant communities. Increased allocation of C to roots may stimulate lateral rooting with extensive root branching. Rooting depth of roots will be influenced by availability of water, nutrients, and other resources in the soil profile. The greater proliferation of roots under elevated atmospheric CO₂ will lead to higher extraction of soil resources with less efficient rooting systems. Carbon dioxide often suppresses plant water use, increasing water use efficiency with relatively less water flow within the plant and consequently in the soil matrix. The enhanced proliferation of roots grown under elevated atmospheric CO₂ concentration may be a strategy which permits adequate nutrient acquisition in the absence of normal water absorption rates. Deeper roots could penetrate into deeper soils to create more resilience to droughts in natural plant communities and agronomic crops. This deeper rooting would be an advantage if climates became drier and altered belowground plant competition.

Variability in response of plant root to CO₂ have been observed in response to the interacting effects of CO₂ with other climatic factors and soil resources coupled with species and even genotypic variability along with levels of the treatment variables. In addition, environmental changes concomitant with the rise of atmospheric CO₂ levels might substantially modify plant and ecosystem level responses. For example, some studies suggest that plants subjected to water stress will respond more favorably to elevated CO₂ while other studies report the opposite response.

Enhanced root growth and biomass production from exposure to increased CO₂ concentrations could deliver more C to the soil profile potentially altering rhizosphere microbiology (populations and dynamics); and inducing possible physicochemical changes in soil by increasing root activity, including rhizodeposition due to root turnover. It is evident from several studies that increasing levels of CO₂ will have virtually no adverse effects on plant root growth except where decreases in efficiency of root systems (specific root activity) and inability of roots to acquire nutrients due to reduced water use/uptake to keep pace with the increased C assimilation would have negative consequences on crop productivity and nutrient content in aboveground organs.

Under favorable conditions of water and nutrients for growth, dry matter partitioning to the root is not altered by CO₂ enrichment. The increase in R/S, frequently observed under limiting conditions of water and/or nutrients, enables the plant to explore a greater soil volume and extract more water and nutrients. However, there are many debates on whether R/S is valuable as a standard root variable due to our inability to accurately estimate belowground biomass, particularly fine roots, root turnover biomass at later stages of the crop and at deeper soil depth apart from microbial decomposition of roots. As observed from many studies, plants allocate more C toward the root growth in the early vegetative stage of development and then decrease. Static assessments of belowground productivity at different stages of crop growth may greatly underestimate the root biomass leading to an exaggeration of R/S values.

Root lifespan has important consequences for plant growth and productivity, plant competition, and C and nutrient cycling. Very limited information is available to quantify the effects of elevated CO₂ on the trade-offs associated with maintaining old roots in comparison with growing new roots. Long-term studies

are needed to detect existing patterns of root lifespan with resource availability and plant strategy and to progress beyond short-term responses to treatments. Factors that should be included in studies of root lifespan in addition to resource availability include root age, plant C status, and pathogen pressure. Because root lifespan differs from leaf lifespan in many species, better knowledge of root lifespan may result in revision of current theories regarding plant adaptation and growth strategies.

Agricultural management practices may have a greater impact on root growth than rising atmospheric CO₂ since management practices influence soil physical, chemical, and biological properties of soil, and consequently root growth dynamics. Less understood are the interactive effects of elevated CO₂ and management practices (tillage, soil moisture, and nutrients) including drought on root dynamics, fine-root production and water-nutrient use efficiency, and the contribution of these processes to plant growth in water and nutrient-limited environments. Increases in water use efficiency and reductions in water use can contribute to enhanced soil water content under elevated CO₂.

Roots are difficult to study because they are hidden within the soil profile and extraction methods destroy the sample. The advent of nondestructive techniques, that is, minirhizotrons or rhizotrons, provide a method of visual and qualitative assessments but do not permit quantitative evaluations of the changes in root biomass or detailed examinations of the branching or root turnover rates. The current state of knowledge regarding root systems is rather simplistic because of the methodological limitations; however, these studies have begun to show that there will be effects of increasing CO₂ on the roots along with the shoots and these responses will have impacts on the utilization of the soil resources, water, nutrients, gases, and temperature. We can continue to strive for more detailed understanding of root responses to changing CO₂; however, these should not be done without considering the interactions with the soil environment. An aspect which is evident from the combined literature is that root function under increased CO₂ does increase when there is no limitation from water, nutrients, or temperature which does suggest that under climate change adoption of management practices which reduces the occurrence of these stresses will have a positive benefit on plant growth and productivity. Understanding these processes will provide the foundation for design of future adaptation and mitigation strategies linking improved soil management practices with crop and root response to increasing CO₂ and will provide a structure for improving the stability of agriculture under climate change.

REFERENCES

- Allen, A.S., J.A. Andrews, A.C. Finzi, R. Matamala, D.D. Richter, and W.H. Schlesinger. 2000. Effects of free-air CO₂ enrichment (FACE) on below-ground processes in *Pinus taeda* forest. *Ecol. Appl.* 10:437–448.
- Allen, L.H., J.C.V. Vu, R.R. Valle, K.J. Boote, and P.H. Jones. 1988. Nonstructural carbohydrates and nitrogen of soybean grown under carbon dioxide enrichment. *Crop Sci.* 28:84–94. doi:10.2135/cropsci1988.0011183X002800010020x
- Baker, J.T., L.H. Allen, and K.J. Boote. 1990. Growth and yield responses of rice to carbon dioxide concentrations. *J. Agric. Sci.* 115:313–320. doi:10.1017/S0021859600075729
- Baker, J.M., T.E. Ochsner, R.T. Venterea, and T.J. Griffis. 2007. Tillage and soil carbon sequestration—What do we really know? *Agric. Ecosyst. Environ.* 118:1–5. doi:10.1016/j.agee.2006.05.014
- Barber, D.A., and J.K. Martin. 1976. The release of organic substances by cereal roots in soil. *New Phytol.* 76:69–80. doi:10.1111/j.1469-8137.1976.tb01439.x

- Barley, K.P. 1970. The configuration of the root system in relation to nutrient uptake. *Adv. Agron.* 22:159–201. doi:10.1016/S0065-2113(08)60268-0
- Batjes, N.H. 1996. Total carbon and nitrogen in the soils of the world. *Eur. J. Soil Sci.* 47:151–163. doi:10.1111/j.1365-2389.1996.tb01386.x
- Baxter, B., T.W. Ashenden, T.H. Sparks, and J.F. Farrar. 1994. Effects of elevated carbon dioxide on three montane grass species. I. Growth and dry matter partitioning. *J. Exp. Bot.* 45(3):305–315. doi:10.1093/jxb/45.3.305
- Bazzaz, F.A. 1990. Response of natural ecosystems to the rising global CO₂ levels. *Annu. Rev. Ecol. Syst.* 21:167–196. doi:10.1146/annurev.es.21.110190.001123
- Bernacchi, C.J., J.S. Coleman, F.A. Bazzaz, and K.D.M. McCaughay. 2000. Biomass allocation in old-field annual species grown in elevated CO₂ environments: No evidence for optimal partitioning. *Glob. Change Biol.* 6:855–863. doi:10.1046/j.1365-2486.2000.00370.x
- Berntson, G.M., and F.A. Bazzaz. 1995. Belowground positive and negative feedbacks on CO₂ growth enhancement. *Plant Soil* 187:119–131. doi:10.1007/BF00017085
- Berntson, G.M., and F.A. Bazzaz. 1997a. Nitrogen cycling in microcosms of yellow birch exposed to elevated CO₂: Simultaneous positive and negative below-ground feedbacks. *Glob. Change Biol.* 3:247–258. doi:10.1046/j.1365-2486.1997.00070.x
- Berntson, G.M., and F.A. Bazzaz. 1997b. Elevated CO₂ and the magnitude and seasonal dynamics of root production and loss in *Betula papyrifera*. *Plant Soil* 190:211–216. doi:10.1023/A:1004222131516
- Berntson, G.M., and F.I. Woodward. 1992. The root system architecture and development of *Senecio vulgaris* in elevated CO₂ and drought. *Funct. Ecol.* 6:324–333. doi:10.2307/2389524
- Bhattacharya, N.C., D.R. Hileman, P.P. Ghosh, R.L. Musser, S. Bhattacharya, and P.K. Biswas. 1990. Interaction of enriched CO₂ and water stress on the physiology and production in sweet potato grown in open-top chambers. *Plant Cell Environ.* 13:933–940. doi:10.1111/j.1365-3040.1990.tb01983.x
- Biswas, P.K., and D.R. Hileman. 1985. Response of vegetation to carbon dioxide: Field studies of sweet potatoes and cowpeas in response to elevated carbon dioxide. Rep. 022. U.S. Dep. of Energy. Carbon Dioxide Res. Div., Office of the Energy Res., Washington, DC.
- Borkert, C.M., and S.A. Barber. 1985. Soybean shoot and root growth and phosphorus concentration as affected by phosphorus placement. *Soil Sci. Soc. Am. J.* 49:152–155. doi:10.2136/sssaj1985.03615995004900010031x
- Bradley, R.I., R. Milne, J. Bell, A. Lilly, C. Jordan, and A. Higgins. 2005. A soil carbon and land use database for the United Kingdom. *Soil Use Manage.* 21:363–369. doi:10.1079/SUM2005351
- Burch, G.J., and G.G. Johns. 1978. Root absorption of water and physiological responses to water deficits by *Festuca arundinacea* Shrub and *Trifolium repens* (L.). *Aust. J. Plant Physiol.* 5:859–871. doi:10.1071/PP9780859
- Caldwell, M.M. 1977. Root structure: The considerable cost of belowground function. In: O.T. Solbrig, editor, *Topics in plant population biology*. Columbia Univ. Press, New York. p. 408–427.
- Caldwell, M.M., and D.M. Eissenstat. 1987. Coping with variability: Examples of tracer use in root function studies. In: J.D. Tenhunen et al., editors, *Plant response to stress*. NATO ASI Series 15G. Springer-Verlag, Berlin. p. 95–106.
- Canadell, J.R., B. Jackson, J.B. Ehleringer, H.A. Mooney, O.E. Sala, and E.D. Schulze. 1996a. Maximum rooting depth of vegetation types at the global scale. *Oecologia* 108:583–595. doi:10.1007/BF00329030
- Canadell, J.G., L.F. Pitelka, and J.S.I. Ingram. 1996b. The effects of elevated CO₂ on plant-soil carbon below ground: Summary and conclusions. *Plant Soil* 187:391–400. doi:10.1007/BF00017102
- Ceulemans, R., and M. Mousseau. 1994. Effects of elevated atmospheric CO₂ on woody plants. *New Phytol.* 127:425–446. doi:10.1111/j.1469-8137.1994.tb03961.x
- Chaudhuri, U.N., R.B. Burnett, M.B. Kirkham, and E.T. Kanemasu. 1986. Effect of carbon dioxide on sorghum yield, root growth, and water use. *Agric. For. Meteorol.* 37:109–122. doi:10.1016/0168-1923(86)90002-X
- Chaudhuri, U.N., M.B. Kirkham, and E.T. Kanemasu. 1990. Root growth of winter wheat under elevated carbon dioxide and drought. *Crop Sci.* 30:853–857. doi:10.2135/cropsci1990.0011183X003000040017x
- Cornish, K., and J.A.D. Zeevaart. 1985. Abscisic acid accumulation in roots of *Xanthium strumarium* (L.) and *Lycopersicon esculentum* (Mill.). In relation to water stress. *Plant Physiol.* 79:653–658. doi:10.1104/pp.79.3.653
- Cox, W.J., R.W. Zobel, H.M. Van Es, and D.J. Otis. 1990. Tillage effects on some soil physical and corn physiological characteristics. *Agron. J.* 82:806–812. doi:10.2134/agronj1990.00021962008200040030x
- Cruz, C., S.H. Lips, and M.A. Martins-Loucao. 1997. Changes in the morphology of roots and leaves of carob seedlings induced by nitrogen source and atmospheric carbon dioxide. *Ann. Bot. (London)* 80:817–823. doi:10.1006/anbo.1997.0524
- Cure, J.D. 1985. Carbon dioxide doubling responses: A crop survey. In: B.R. Strain and J.D. Cure, editors, *Direct effects of increasing carbon dioxide on vegetation*. DOE/ER-0238. U.S. Dep. of Energy, Washington, DC. p. 99–116.
- Curtis, P.S. 1996. A meta-analysis of leaf gas exchange and nitrogen in trees grown under elevated carbon dioxide. *Plant Cell Environ.* 19:127–137. doi:10.1111/j.1365-3040.1996.tb00234.x
- Curtis, P.S., L.M. Balduman, B.G. Drake, and D.F. Whigham. 1990. Elevated atmospheric CO₂ effects on belowground processes in C₃ and C₄ estuarine marsh communities. *Ecology* 71:2001–2006. doi:10.2307/1937608
- Curtis, P.S., and X. Wang. 1998. A meta-analysis of elevated CO₂ effects on woody plant mass, form and physiology. *Oecologia* 113:299–313. doi:10.1007/s004420050381
- Dahlman, R.C., B.R. Strain, and H.H. Rogers. 1985. Research on the response of vegetation to elevated atmospheric carbon dioxide. *J. Environ. Qual.* 14:1–8. doi:10.2134/jeq1985.00472425001400010001x
- Davidson, R.L. 1969. Effect of root: Leaf temperature differentials on root:shoot ratios in some pasture grasses and clover. *Ann. Bot. (London)* 33:561–569.
- Del Castillo, D., B. Acock, V.R. Reddy, and M.C. Acock. 1989. Elongation and branching of roots on soybean plants in a carbon dioxide-enriched aerial environment. *Agron. J.* 81:692–695. doi:10.2134/agronj1989.00021962008100040026x
- Derner, J.D., H.B. Johnson, B.A.K. Iimball, P.J. Pinter, Jr., H.W. Polley, C.R. Tischler et al. 2003. Above and below ground responses of C₃ and C₄ species mixtures in elevated CO₂ and soil water availability. *Glob. Change Biol.* 9:452–460. doi:10.1046/j.1365-2486.2003.00579.x
- Drake, B.G., and M.A. González-Meier. 1997. More efficient plants: A consequence of rising atmospheric CO₂? *Annu. Rev. Plant Physiol.* 48:609–639. doi:10.1146/annurev.arplant.48.1.609
- Dwyer, L.M., B.L. Ma, H.N. Hayhoe, and J.L.B. Culley. 1995. Tillage effects on soil temperature shoot dry matter accumulation and corn grain yield. *J. Sustain. Agric.* 5:85–99. doi:10.1300/J064v05n01_07
- Dwyer, L.M., B.L. Ma, D.W. Stewart, H.N. Hayhoe, D. Balchin, J.L.B. Culley, and M. McGovern. 1996. Root mass distribution under conventional and conservation tillage. *Can. J. Soil Sci.* 76:23–28. doi:10.4141/cjss96-004
- Eamus, D., and P.G. Jarvis. 1989. Direct effects of CO₂ increases on trees and forests (natural and commercial) in the UK. *Adv. Ecol. Res.* 19:1–55. doi:10.1016/S0065-2504(08)60156-7
- Eissenstat, D.M. 1992. Costs and benefits of constructing roots of small diameter. *J. Plant Nutr.* 15:763–782. doi:10.1080/01904169209364361
- Eissenstat, D.M., and R.D. Yanai. 1997. The ecology of root life span. *Adv. Ecol. Res.* 27:1–62. doi:10.1016/S0065-2504(08)60005-7
- Eissenstat, D.M., C.E. Wells, R.D. Yanai, and J.L. Whitbeck. 2000. Building roots in a changing environment: Implications for root longevity. *New Phytol.* 147:33–42. doi:10.1046/j.1469-8137.2000.00686.x
- Fahey, T.J., and J.W. Hughes. 1994. Fine root dynamics in a northern hardwood forest ecosystem Hubbard Brook Experimental Forest, NH. *J. Ecol.* 82:533–548. doi:10.2307/2261262
- Ferris, R., and G. Taylor. 1993. Contrasting effects of elevated on the root and shoot growth of four native herbs commonly found in chalk grassland. *New Phytol.* 125:855–866. doi:10.1111/j.1469-8137.1993.tb03934.x
- Fitter, A.H. 1985. Functional significance of root morphology and root system architecture. In: A.H. Fitter et al., editors, *Ecological interactions in soil*. Blackwell Scientific Publ., Oxford. p. 87–106.
- Fitter, A.H. 1986. The topology and geometry of plant root systems: Influence of watering rate on root system topology in *Trifolium pratense*. *Ann. Bot. (London)* 58:91–101.
- Fitter, A.H. 1987. An architectural approach to the comparative ecology of plant root systems. *New Phytol.* 106:61–77. doi:10.1111/j.1469-8137.1987.tb04683.x
- Fitter, A.H., J.D. Graves, J. Wolfenden, G.K. Self, T.K. Brown, D. Bogie, and T.A. Mansfield. 1997. Root production and turnover and carbon budgets of two contrasting grasslands under ambient and elevated atmospheric carbon dioxide concentrations. *New Phytol.* 137:247–255. doi:10.1046/j.1469-8137.1997.00804.x
- Fitter, A.H., G.K. Self, T.K. Brown, D. Bogie, J.D. Graves, D. Benham, and P. Ineson. 1999. Root production and turnover in an upland grassland subjected to artificial soil warming respond to radiation flux and nutrients, not temperature. *Oecologia* 120:575–581. doi:10.1007/s004420050892

- Fitter, A.H., G.K. Self, J. Wolfenden, M.M.I. Van Vuuren, T.K. Brown, L. Williamson et al. 1996. Root production and mortality under elevated atmospheric carbon dioxide. *Plant Soil* 187:299–306. doi:10.1007/BF00017095
- Forbes, P.J., K.E. Black, and J.E. Hooker. 1997. Temperature-induced alteration to root longevity in *Lolium perenne*. *Plant Soil* 190:87–90. doi:10.1023/A:1004298804353
- French, C.J., and J. Alsbury. 1989. Supplementary lighting and CO₂ mist influence rooting of *Camellia japonica*. *HortScience* 24:452–454.
- Gill, R.A., and R.B. Jackson. 2000. Global patterns of root turnover for terrestrial ecosystems. *New Phytol.* 147:13–31. doi:10.1046/j.1469-8137.2000.00681.x
- Goins, G.D., and M.P. Russelle. 1996. Fine root demography in alfalfa (*Medicago sativa* (L.)). *Plant Soil* 185:281–291. doi:10.1007/BF02257534
- Guo, L.B., and R.M. Gifford. 2002. Soil carbon stocks and land use change: A meta-analysis. *Glob. Change Biol.* 8:345–360. doi:10.1046/j.1354-1013.2002.00486.x
- Guswa, A.J. 2010. Effect of plant-uptake strategy on the water optimal root depth. *Water Resour. Res.* 46:W09601. doi:10.1029/2010WR009122. doi:10.1029/2010WR009122
- Hanley, M., S. Trošimov, and G. Taylor. 2004. Species level effects more important than functional group level responses to elevated CO₂: Evidence from simulated turves. *Funct. Ecol.* 18:304–313. doi:10.1111/j.0269-8463.2004.00845.x
- Hatfield, J.L., K.J. Boote, B.A. Kimball, L.H. Ziska, R.C. Izaurralde, D. Ort et al. 2011. Climate impacts on agriculture: Implications for crop production. *Agron. J.* 103:351–370.
- Havelka, U.D., V.A. Wittenbach, and M.G. Boyle. 1984. CO₂ enrichment effects on wheat yield and physiology. *Crop Sci.* 24:1163–1168. doi:10.2135/cropsci1984.0011183X002400060037x
- Hayes, D.C., and T.R. Seastedt. 1987. Root dynamics of tall grass prairie in wet and dry years. *Can. J. Bot.* 65:787–791. doi:10.1139/b87-105
- Heinemann, A.B., H.N. Maia Ad, D. Dourado-Neto, K.T. Ingram, and G. Hoogenboom. 2006. Soybean [*Glycine max* (L.) Merr.] growth and development response to CO₂ enrichment under different temperature regimes. *Eur. J. Agron.* 24:52–61. doi:10.1016/j.eja.2005.04.005
- Helal, H.M., and D. Sauerbeck. 1986. Effect of plant roots on carbon metabolism of soil microbial biomass. *Z. Pflanzenernaehr. Bodenkd.* 149:181–188. doi:10.1002/jpln.19861490205
- Hendrick, R.L., and K.S. Pregitzer. 1992. The demography of fine roots in a northern hardwood forest. *Ecology* 73:1094–1104. doi:10.2307/1940183
- Hendrick, R.L., and K.S. Pregitzer. 1993a. The dynamics of fine root length, biomass, and nitrogen content in two northern hardwood ecosystems. *Can. J. For. Res.* 23:2507–2520. doi:10.1139/x93-312
- Hendrick, R.L., and K.S. Pregitzer. 1993b. Patterns of fine root mortality in two sugar maple forests. *Nature (London)* 361:59–61. doi:10.1038/361059a0
- Hendrick, R.L., and K.S. Pregitzer. 1997. The relationship between fine root demography and the soil environment in northern hardwood forests. *Ecoscience* 4:99–105.
- Hendricks, J.J., K.J. Nadelhoffer, and J.D. Aber. 1993. Assessing the role of fine roots in carbon and nutrient cycling. *Trends Ecol. Evol.* 8:174–178. doi:10.1016/0169-5347(93)90143-D
- Hocking, P.J., and C.P. Meyer. 1991. Effects of CO₂ enrichment and nitrogen stress on growth, and partitioning of dry matter and nitrogen in wheat and maize. *Aust. J. Plant Physiol.* 18:339–356. doi:10.1071/PP9910339
- Hodge, A., and P. Millard. 1998. Effect of elevated CO₂ on carbon partitioning and exudates release from *Plantago lanceolata* seedlings. *Physiol. Plant.* 103:280–286. doi:10.1034/j.1399-3054.1998.1030216.x
- Huck, M.G., G. Hoogenboom, and C.M. Peterson. 1987. Soybean root senescence under drought stress. In: H.M. Taylor, editor, *Minirhizotron observation tubes: Methods and applications for measuring rhizosphere dynamics*. ASA Spec. Publ. 50. ASA, Madison, WI.
- Idso, S.B., and B.A. Kimball. 1991. Effects of two and a half years of atmospheric CO₂ enrichment on the root density distribution of three-year-old sour orange trees. *Agric. For. Meteorol.* 55:345–349. doi:10.1016/0168-1923(91)90070-7
- Idso, S.B., and B.A. Kimball. 1992. Seasonal root growth of sour orange trees maintained in atmospheres of ambient and elevated CO₂ concentrations. *Plant Cell Environ.* 15:337–341. doi:10.1111/j.1365-3040.1992.tb00982.x
- Idso, S.B., B.A. Kimball, and J.R. Mauney. 1988. Effects of atmospheric CO₂ enrichment on root:shoot ratios of carrot, radish, cotton and soybean. *Agric. Ecosyst. Environ.* 21:293–299. doi:10.1016/0167-8809(88)90095-3
- Imai, K., D.F. Coleman, and T. Yanagisawa. 1985. Increase in atmospheric partial pressure of carbon dioxide and growth and yield of rice (*Oryza sativa* (L.)). *Jpn. J. Crop. Sci.* 54:413–418. doi:10.1626/jcs.54.413
- Imai, K., and Y. Murata. 1976. Effect of carbon dioxide concentration on growth and dry matter production in crop plants. *Japanese J. of Crop Sci.* 45:598–606. doi:10.1626/jcs.45.598
- IPCC. 2001. *Climate change 2001: The scientific basis*. Cambridge Univ. Press, Cambridge, UK.
- IPCC. 2007. *Climate change 2007: The physical science basis*. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge Univ. Press, Cambridge, UK.
- Jackson, R.B., and M.M. Caldwell. 1989. The timing and degree of root proliferation in fertile-soil microsites for three cold-desert perennials. *Oecologia* 81:149–153.
- Jackson, R.B., J. Canadell, J.R. Ehleringer, H.A. Mooney, O.E. Sala, and E.D. Schulze. 1996. A global analysis of root distributions for terrestrial biomes. *Oecologia* 108:389–411. doi:10.1007/BF00333714
- Jackson, R.B., H.A. Mooney, and E.D. Schulze. 1997. A global budget for fine-root biomass, surface area, and nutrient contents. *Proc. Natl. Acad. Sci. USA* 94:7362–7366. doi:10.1073/pnas.94.14.7362
- Jobbagy, E.G., and R.B. Jackson. 2000. The vertical distribution of soil organic carbon and its relation to climate and vegetation. *Ecol. Appl.* 10:423–436. doi:10.1890/1051-0761(2000)010[0423:TVDOS0]2.0.CO;2
- Jones, H., R.A. Leigh, A.D. Tomos, and R.G. Wyn Jones. 1987. The effect of abscisic acid on cell turgor pressures, solute content and growth of wheat roots. *Planta* 170:257–262. doi:10.1007/BF00397896
- Keeling, C.D., and T.P. Whorf. 2005. Atmospheric CO₂ records from sites in the SIO air sampling network. In: *Trends: A compendium of data on global change*. Carbon Dioxide Information Analysis Ctr., Oak Ridge Natl. Lab., U.S. Dep. of Energy, Oak Ridge, TN.
- Keith, H., J.M. Oades, and J.K. Martin. 1986. Input of carbon to soil from wheat plants. *Soil Biol. Biochem.* 18:445–449. doi:10.1016/0038-0717(86)90051-9
- Kell, D.B. 2011. Breeding crop plants with deep roots: Their role in sustainable carbon, nutrient and water sequestration. *Ann. Bot. (London)* 10.1093/aob/mcr175.
- Kimball, B.A. 1983. Carbon dioxide and agricultural yield: An assemblage and analysis of 430 prior observations. *Agron. J.* 75:779–786. doi:10.2134/agronj1983.00021962007500050014x
- Kimball, B.A. 2011. Lessons from FACE: CO₂ effects and interactions with water, nitrogen and temperature. In: D. Hillel and C. Rosenzweig, editors, *Handbook of climate change and agroecosystems: Impacts, adaptation, and mitigation*. Imperial College Press, Hackensack, NJ, p. 87–107.
- Kimball, B.A., K. Kobayashi, and M. Bindi. 2002. Responses of agricultural crops to free-air CO₂ enrichment. *Adv. Agron.* 77:293–368. doi:10.1016/S0065-2113(02)77017-X
- King, J.S., K.S. Pregitzer, and D.R. Zak. 1999. Clonal variation in above- and belowground growth responses of *Populus tremuloides* (Michaux): Influence of soil warming and nutrient availability. *Plant Soil* 217:119–130. doi:10.1023/A:1004560311563
- Klepper, B., H.M. Taylor, M.G. Huck, and E.L. Fiscus. 1973. Water relations and growth of cotton in drying soils. *Agron. J.* 65:307–310. doi:10.2134/agronj1973.00021962006500020036x
- Kramer, P.J. 1981. Carbon dioxide concentration, photosynthesis, and dry matter production. *Bioscience* 31:29–33. doi:10.2307/1308175
- Krauss, U., and J.W. Deacon. 1994. Root turnover of groundnut (*Arachis hypogaea* (L.)) in soil tubes. *Plant Soil* 166:259–270. doi:10.1007/BF00008339
- Laforge, F., C. Lussier, Y. Desjardins, and A. Gosselin. 1991. Effect of light intensity and enrichment during in-vitro rooting on subsequent growth of plantlets of strawberry, raspberry and asparagus in acclimatization. *Sci. Hortic. (Amsterdam)* 47:259–269. doi:10.1016/0304-4238(91)90009-N
- Lampurlanés, J.P., C. Angás, and C. Cantero-Martínez. 2001. Root growth, soil water content and yield of barley under different tillage systems on two soils in semiarid conditions. *Field Crops Res.* 69:27–40. doi:10.1016/S0378-4290(00)00130-1
- Lawlor, D.W., and R.A.C. Mitchell. 1991. The effect of increasing CO₂ on crop photosynthesis and production: A review of field studies. *Plant Cell Environ.* 14:807–818. doi:10.1111/j.1365-3040.1991.tb01444.x
- Liljeroth, E., J.A. Van Veen, and H.J. Miller. 1990. Assimilate translocation to the rhizosphere of two wheat lines and subsequent utilization by rhizosphere microorganisms at two soil nitrogen concentrations. *Soil Biol. Biochem.* 22:1015–1021. doi:10.1016/0038-0717(90)90026-V
- Lin, W.C., and J.M. Molnar. 1981. Effects of mist and high intensity supplementary lighting on propagation of selected woody ornamentals. *Can. J. Plant Sci.* 61:965–969. doi:10.4141/cjps81-142

- Lorenz, K., and R. Lal. 2005. The depth distribution of soil organic carbon in relation to land use and management and the potential of carbon sequestration in subsoil horizons. *Adv. Agron.* 88:35–66. doi:10.1016/S0065-2113(05)88002-2
- Lotze-Campen, H., and H.J. Schellnhuber. 2009. Climate impacts and adaptation options in agriculture: What we know and what we don't know. *J. Verbraucherschutz und Lebensmittelsicherheit.* 4:145–150. doi:10.1007/s00003-009-0473-6
- Luxmoore, R.J., E.G. O'Neill, J.M. Ells, and H.H. Rogers. 1986. Nutrient uptake and growth responses of Virginia pine to elevated atmospheric carbon dioxide. *J. Environ. Qual.* 15:244–251. doi:10.2134/jeq1986.00472425001500030008x
- McGuire, D., J.M. Melillo, and L.A. Joyce. 1995. The role of nitrogen in the response of forest net primary production to elevated atmospheric carbon dioxide. *Annu. Rev. Ecol. Syst.* 26:473–503. doi:10.1146/annurev.es.26.110195.002353
- McMaster, G.S., D.R. LeCain, J.A. Morgan, L. Aiguo, and D.L. Hendrix. 1999. Elevated CO₂ Increases wheat CER, leaf and tiller development, and shoot and root growth. *J. Agron. Crop Sci.* 183:119–128. doi:10.1046/j.1439-037x.1999.00325.x
- Milchunas, D.G., and W.K. Lauenroth. 1992. Carbon dynamics and estimates of primary production by harvest, (14)C dilution, and (14)C turnover. *Ecology* 73:593–607. doi:10.2307/1940765
- Milchunas, D.G., W.K. Lauenroth, J.S. Singh, C.V. Dole, and H.W. Hunt. 1985. Root turnover and production by 14C dilution: Implications of carbon partitioning in plants. *Plant Soil* 88:353–365. doi:10.1007/BF02197492
- Mo, G.D., M.B. Nie, H. Kirkham He, L.K. Ballou, F.W. Caldwell, and E.T. Kanemasu. 1992. Root and shoot weight in a tall grass prairie under elevated carbon dioxide. *Environ. Exp. Bot.* 32(3):193–201. doi:10.1016/0098-8472(92)90002-J
- Mokany, K., R.J. Reason, and A.S. Prokushkin. 2006. Critical analysis of root: Shoot ratios in terrestrial biomes. *Glob. Change Biol.* 12:84–96. doi:10.1111/j.1365-2486.2005.001043.x
- Mulkey, T.J., M.L. Evans, and K.M. Kuzmanoff. 1983. The kinetics of abscisic acid action on root growth and gravitropism. *Planta* 157:150–157. doi:10.1007/BF00393649
- Nepstad, D.C., C.R. Decarvalho, E.A. Davidson, P.H. Jipp, P.A. Lefebvre, G.H. Negrelros et al. 1994. The role of deep roots in the hydrological and carbon cycles of Amazonian forests and pastures. *Nature (London)* 372:666–669. doi:10.1038/372666a0
- Newell, R.L., and W.W. Wilhelm. 1987. Conservation tillage and irrigation effects on corn root development. *Agron. J.* 79:160–165. doi:10.2134/agronj1987.00021962007900010033x
- Norby, R.J. 1987. Nodulation and nitrogenase activity in nitrogen-fixing woody plants stimulated by CO₂ enrichment of the atmosphere. *Physiol. Plant.* 71:77–82. doi:10.1111/j.1399-3054.1987.tb04620.x
- Norby, R.J. 1994. Issues and perspectives for investigating responses to elevated atmospheric carbon dioxide. *Plant Soil* 165:9–20. doi:10.1007/BF00009958
- Norby, R.J., C.A. Gunderson, S.D. Wullschleger, E.G. O'Neill, and M.K. McCracken. 1992. Productivity and compensatory responses of yellow-poplar trees in elevated. *Nature (London)* 357:322–332. doi:10.1038/357322a0
- Norby, R.J., and E.G. O'Neill. 1989. Growth dynamics and water use of seedlings of *Quercus alba* (L.) in CO₂ enriched atmospheres. *New Phytol.* 111:491–500. doi:10.1111/j.1469-8137.1989.tb00712.x
- Norby, R.J., E.G. O'Neill, W.G. Hood, and R.J. Luxmoore. 1987. Carbon allocation, root exudation and mycorrhizal colonization of *Pinus echinata* seedlings grown under CO₂ enrichment. *Tree Physiol.* 3:203–210. doi:10.1093/treephys/3.3.203
- Norby, R.J., E.G. O'Neill, and R.J. Luxmoore. 1986a. Effects of atmospheric CO₂ enrichment on the growth and mineral nutrition of *Quercus alba* seedlings in a nutrient poor soil. *Plant Physiol.* 82:83–89. doi:10.1104/pp.82.1.83
- Norby, R.J., J. Pastor, and J.M. Melillo. 1986b. Carbon nitrogen interactions in CO₂-enriched white oak: Physiological and long-term perspectives. *Tree Physiol.* 2:233–241. doi:10.1093/treephys/2.1-2-3.233
- Obrist, D., and J.A. Arnone. 2003. Increasing CO₂ accelerates root growth and enhances water acquisition during early stages of development in *Larrea tridentate*. *New Phytol.* 159:175–184. doi:10.1046/j.1469-8137.2003.00791.x
- Oechel, W.C., and B.R. Strain. 1985. Native species responses to increased atmospheric carbon dioxide concentration. In: B.R. Strain, and J.D. Cure, editors, *Direct effects of increasing carbon dioxide on vegetation.* DOE/ER-0238. U.S. Dep. of Energy, Washington, DC. p. 117–154.
- O'Neill, E.G., R.J. Luxmoore, and R.J. Norby. 1987a. Elevated atmospheric CO₂ effects on seedling growth, nutrient uptake, and rhizosphere bacterial populations of *Liriodendron tulipifera* (L.). *Plant Soil* 104:3–11. doi:10.1007/BF02370618
- O'Neill, E.G., R.J. Luxmoore, and R.J. Norby. 1987b. Increases in mycorrhizal colonization and seedling growth in *Pinus echinata* and *Quercus alba* in an enriched CO₂ atmosphere. *Can. J. For. Res.* 17:878–883. doi:10.1139/x87-139
- Paustian, K., E.T. Elliot, G.A. Peterson, and K. Killian. 1996. Modeling climate, CO₂ and management impacts on soil carbon in semi-arid agro ecosystems. *Plant Soil* 187:351–365. doi:10.1007/BF00017100
- Pendall, E., J.Y. King, A.R. Mosier, J.A. Morgan, and D.S. Milchunas. 2005. Stable isotope constraints on net ecosystem production in elevated CO₂ experiments. In: L.B. Flanagan, J.R. Ehleringer, and D.E. Pataki, editors, *Stable isotopes and biosphere atmosphere interactions.* Elsevier, New York. p. 182–198.
- Pilet, P.E., and M. Saugy. 1987. Effect on root growth of endogenous and applied IAA and ABA: A critical reexamination. *Plant Physiol.* 83:33–38. doi:10.1104/pp.83.1.33
- Pilumwong, J., C. Senthong, S. Srichuwong, and K.T. Ingram. 2007. Effects of temperature and elevated CO₂ on shoot and root growth of Peanut (*Arachis hypogaea* (L.)) grown in controlled environment chambers. *Sci. Asia* 33:79–87. doi:10.2306/scienceasia1513-1874.2007.33.079
- Pregitzer, K.S., D.R. Zak, P.S. Curtis, M.E. Kubiske, J.A. Teeri, and C.S. Vogel. 1995. Atmospheric CO₂, soil nitrogen and turnover of fine roots. *New Phytol.* 129:579–585. doi:10.1111/j.1469-8137.1995.tb03025.x
- Pregitzer, K.S., D.R. Zak, J. Maziasz, J. DeForest, P.S. Curtis, and J. Lussenhop. 2000. Fine-root growth, mortality, and morphology in a factorial elevated atmospheric CO₂ soil N availability experiment. *Ecol. Appl.* 10:18–33.
- Prior, S.A. 1992. Cotton root response to free-air CO₂ enrichment. PhD. diss. Auburn Univ., Auburn, AL.
- Prior, S.A., H.H. Rogers, G.L. Mullins, and G.B. Runion. 2003. The effects of elevated atmospheric CO₂ and soil P placement on cotton root deployment. *Plant Soil* 255:179–187. doi:10.1023/A:1026143410238
- Prior, S.A., H.H. Rogers, G.B. Runion, and G.R. Hendrey. 1994a. Free-air CO₂ enrichment of cotton: Vertical and lateral root distribution patterns. *Plant Soil* 165:33–44. doi:10.1007/BF00009960
- Prior, S.A., H.H. Rogers, G.B. Runion, B.A. Kimball, J.R. Mauney, K.F. Lewin, J. Nagy, and G.R. Hendrey. 1995. Free-air CO₂ enrichment of cotton: Root morphological characteristics. *J. Environ. Qual.* 24:678–683. doi:10.2134/jeq1995.00472425002400040019x
- Prior, S.A., H.H. Rogers, G.B. Runion, and J.R. Mauney. 1994b. Effects of free-air CO₂ enrichment on cotton root growth. *Agric. For. Meteorol.* 70:69–86. doi:10.1016/0168-1923(94)90048-5
- Prior, S.A., G.B. Runion, H.H. Rogers, H.A. Torbert, and D.W. Reeves. 2005. Elevated atmospheric CO₂ effects on biomass production and soil carbon in conventional and conservation cropping systems. *Glob. Change Biol.* 11:657–665. doi:10.1111/j.1365-2486.2005.00935.x
- Prior, S.A., H.A. Torbert, G.B. Runion, G.L. Mullins, H.H. Rogers, and J.R. Mauney. 1998. Effects of CO₂ enrichment on cotton nutrient dynamics. *J. Plant Nutr.* 21:1407–1426. doi:10.1080/01904169809365492
- Pritchard, S.G., and J.S. Amthor. 2005. Crops and environmental change: An introduction to effects of global warming, increasing atmospheric and O₃ concentrations and soil salinization on crop physiology and yield. The Haworth Press, Binghamton, NY.
- Pritchard, S.G., S.A. Prior, H.H. Rogers, A. Micheal, G. Davis, B. Runion, and T.W. Popham. 2006. Effects of elevated atmospheric CO₂ on root dynamics and productivity of sorghum grown under conventional and conservation agricultural Management practices. *Agric. Ecosyst. Environ.* 113:175–183. doi:10.1016/j.agee.2005.09.010
- Pritchard, S.G., and H.H. Rogers. 2000. Spatial and temporal deployment of crop roots in CO₂-enriched environments. *New Phytol.* 147:55–71. doi:10.1046/j.1469-8137.2000.00678.x
- Pritchard, S.G., H.H. Rogers, S.A. Prior, and C.M. Peterson. 1999. Elevated CO₂ and plant structure: A review. *Glob. Change Biol.* 5:807–837. doi:10.1046/j.1365-2486.1999.00268.x
- Qin, Z., and Y. Huang. 2010. Quantification of soil organic carbon sequestration potential in cropland: A model approach. *Sci. China-Life Sci.* 53:868–884. doi:10.1007/s11427-010-4023-3
- Rastetter, E.B., G.I. Agren, and G.R. Shaver. 1997. Responses of N limited ecosystems to increased CO₂: A balanced-nutrition, coupled-element-cycles model. *Ecol. Appl.* 7:444–460.
- Reynolds, J.F., and J.H.M. Thornley. 1982. A shoot: Root partitioning model. *Ann. Bot. (London)* 49:585–597.

- Rillig, M.C., S.F. Wright, M.F. Allen, and C.B. Field. 1999. Rise in carbon dioxide changes soil structure. *Nature (London)* 400:628. doi:10.1038/23168
- Robinson, D. 2004. Scaling the depths: Below-ground allocation in plants, forests and biomes. *Funct. Ecol.* 18:290–295. doi:10.1111/j.0269-8463.2004.00849.x
- Rogers, H.H., G.E. Bingham, J.D. Cure, J.M. Smith, and K.A. Surano. 1983. Responses of selected plant species to elevated carbon dioxide in the field. *J. Environ. Qual.* 12:569–574. doi:10.2134/jeq1983.00472425001200040028x
- Rogers, H.H., C.M. Peterson, J.M. McCrimmon, and J.D. Cure. 1992. Response of soybean roots to elevated atmospheric carbon dioxide. *Plant Cell Environ.* 15:749–752. doi:10.1111/j.1365-3040.1992.tb01018.x
- Rogers, H.H., S.A. Prior, and E.G. O'Neill. 1993. Cotton root and rhizosphere responses to free-air CO₂ enrichment. In: G.R. Hendrey, editor, Free-air CO₂ enrichment for plant research in the field. CRC Press, Boca Raton, FL.
- Rogers, H.H., S.A. Prior, G.B. Runion, and R.J. Mitchell. 1996. Root to shoot ratio of crops as influenced by CO₂. *Plant Soil* 187:229–248. doi:10.1007/BF00017090
- Rogers, H.H., G.B. Runion, and S.V. Krupa. 1994. Plants' responses to atmospheric enrichment with emphasis on roots and the rhizosphere. *Environ. Pollut.* 83:155–189. doi:10.1016/0269-7491(94)90034-5
- Rogers, H.H., G.B. Runion, S.V. Krupa, and S.A. Prior. 1997. Plant responses to atmospheric carbon dioxide enrichment: Implications in root–soil–microbe interactions. In: L.H. Allen et al., editors, *Advances in carbon dioxide effects research*. ASA, Madison, WI. p. 1–34.
- Rogers, H.H., G.B. Runion, S.A. Prior, and H.A. Torbert. 1999. Response of plants to elevated atmospheric CO₂: Root growth, mineral nutrition, and soil carbon. In: Y. Luo and H.A. Mooney, editors, *Carbon dioxide and environmental stress*. Academic Press, San Diego, CA. p. 215–244.
- Rose, D.A. 1983. The description of the growth of root systems. *Plant Soil* 75:405–415. doi:10.1007/BF02369974
- Roseberg, R.J., and E.L. McCoy. 1992. Tillage and traffic induced changes in macro porosity and macropore continuity: Air permeability assessment. *Soil Sci. Soc. Am. J.* 56:1261–1267. doi:10.2136/sssaj1992.03615995005600040042x
- Salsman, K.J., D.N. Jordan, S.D. Smith, and D.S. Neuman. 1999. Effect of atmospheric enrichment on root growth and carbohydrate allocation of *Phaseolus* spp. *Int. J. Plant Sci.* 160:1075–1081.
- Sarker, B.C., and M. Hara. 2009. Effects of elevated CO₂ and water stress on root structure and hydraulic conductance of *Solanum melongena* L. *Bangladesh J. Bot.* 38:55–63.
- Schenk, H.J., and R.B. Jackson. 2002a. The global biogeography of roots. *Ecol. Monogr.* 72:311–328. doi:10.1890/0012-9615(2002)072[0311:TGBOR]2.0.CO;2
- Schenk, H.J., and R.B. Jackson. 2002b. Rooting depths, lateral root spreads and below-ground/above-ground allometries of plants in water-limited ecosystems. *J. Ecol.* 90:480–494. doi:10.1046/j.1365-2745.2002.00682.x
- Smucker, A.J.M. 1993. Soil environmental modifications of root dynamics and measurement. *Annu. Rev. Phytopathol.* 31:191–216. doi:10.1146/annurev.py.31.090193.001203
- Stulen, I., and J. den Hertog. 1993. Root growth and functioning under atmospheric CO₂ enrichment. *Vegetatio* 104-105:99–115. doi:10.1007/BF00048147
- Suter, D., M. Frehner, B.U. Fischer, J. Nösberger, and A. Lüscher. 2002. Elevated CO₂ increases carbon allocation to the roots of *Lolium perenne* under free-air CO₂ enrichment but not in a controlled environment. *New Phytol.* 154:65–75. doi:10.1046/j.1469-8137.2002.00368.x
- Swinnen, J., J.A. Van Veen, and R. Merckx. 1994. C pulse-labeling of field grown spring wheat: An evaluation of its use in rhizosphere carbon budget estimations. *Soil Biol. Biochem.* 26:161–170. doi:10.1016/0038-0717(94)90159-7
- Swinnen, J., J.A. Van Veen, and R. Merckx. 1995. Root decay and turnover of rhizodeposits in field-grown winter wheat and spring barley estimated by ¹⁴C pulse-labeling. *Soil Biol. Biochem.* 27:211–217. doi:10.1016/0038-0717(94)00161-S
- Vanaja, M., P. Raghuram Reddy, N. Jyothi Lakshmi, M. Maheswari, P. Vagheera, P. Ratnakumar et al. 2007. Effect of elevated atmospheric CO₂ concentrations on growth and yield of black gram (*Vigna mungo* (L.) Hepper) – A rainfed pulse crop. *Plant Soil Environ.* 53(2):81–88.
- van der Werf, A., A. Kooijman, R. Welschen, and H. Lambers. 1988. Respiratory energy costs for the maintenance of biomass, for growth and for ion uptake in roots of *Carex diandra* and *Carex acutiformis*. *Physiol. Plant.* 72:483–491. doi:10.1111/j.1399-3054.1988.tb09155.x
- van Noordwijk, M., G. Brouwer, H. Koning, F.W. Meijboom, and W. Grzebisz. 1994. Production and decay of structural root material of winter wheat and sugar beet in conventional and integrated cropping systems. *Agric. Ecosyst. Environ.* 51:99–113. doi:10.1016/0167-8809(94)90037-X
- van Noordwijk, M., P. Martikainen, P. Bottner, E. Cuevas, C. Rouland, and S.S. Dhillon. 1998. Global change and root function. *Glob. Change Biol.* 4:759–772. doi:10.1046/j.1365-2486.1998.00192.x
- Van Veen, J.A., E. Liljeroth, L.J.A. Lekkerkerk, and S.C. van de Geijn. 1991. Carbon fluxes in plant-soil systems at elevated atmospheric CO₂ levels. *Ecol. Appl.* 1(2):175–181. doi:10.2307/1941810
- Van Vuuren, M.M.I., D. Robinson, A.H. Fitter, S.D. Chasalow, L. Williamson, and J.A. Raven. 1997. Effects of elevated atmospheric CO₂ and soil water availability on root biomass, root length and N, P, and K uptake by wheat. *New Phytol.* 135:455–466. doi:10.1046/j.1469-8137.1997.00682.x
- Walton, D.C., M.A. Harrison, and P. Cote. 1976. The effects of water stress on abscisic acid levels and metabolism in roots of *Phaseolus vulgaris* and other plants. *Planta* 131:141–144. doi:10.1007/BF00389985
- Wan, S., R.J. Norby, K.S. Pregitzer, J. Ledford, and E.G. O'Neill. 2004. CO₂ enrichment and warming of the atmosphere enhance both productivity and mortality of maple tree fine roots. *New Phytol.* 162:437–446. doi:10.1111/j.1469-8137.2004.01034.x
- Wang, Y.G., Y. Li, X.H. Ye, Y. Chu, and X.P. Wang. 2010. Profile storage of organic inorganic carbon in soil: From forest to desert. *Sci. Total Environ.* 408:1925–1931. doi:10.1016/j.scitotenv.2010.01.015
- Warrick, R.A. 1988. Carbon dioxide, climatic change and agriculture. *Geogr. J.* 154:221–233. doi:10.2307/633848
- Watson, C.A., J.M. Ross, U. Bangnaresi, G.F. Minotta, F. Roffi, D. Atkinson et al. 2000. Environmental induced modifications to root longevity in *Lolium perenne* and *Trifolium repens*. *Ann. Bot. (London)* 85:397–401. doi:10.1006/anbo.1999.1048
- Watts, S., J.L. Rodriguez, S. Evans, and W.J. Davies. 1981. Root and shoot growth of plants treated with abscisic acid. *Ann. Bot. (London)* 47:595–602.
- Wechsung, G. 1999. The effects of free-air CO₂ enrichment and soil water availability on spatial and seasonal patterns of wheat root growth. *Global Change Biol.* 5:519–529.
- Wechsung, G., F. Wechsung, G.W. Wall, F.J. Adamsen, B.A. Kimball, R.L. Garcia, P.J. Pinter, Jr., and T. Kartschall. 1995. Biomass and growth rate of a spring wheat root system grown in free-air CO₂ enrichment (FACE) and ample moisture. *J. Biogeogr.* 22:623–634. doi:10.2307/2845963
- Wechsung, G., F. Wechsung, G.W. Wall, F.J. Adamsen, B.A. Kimball, P.J. Pinter, Jr., T. Kartschall, R.L. Garcia, and R.L. LaMorte. 1999. The effects of free-air CO₂ enrichment and soil water availability on spatial and seasonal patterns of wheat root growth. *Glob. Change Biol.* 5:519–529. doi:10.1046/j.1365-2486.1999.00243.x
- Whipps, J.M. 1984. Environmental factors affecting the loss of carbon from the roots of wheat and barley seedlings. *J. Exp. Bot.* 35:767–773. doi:10.1093/jxb/35.6.767
- Whipps, J.M. 1985. Effect of CO₂ concentration on growth, carbon distribution and loss of carbon from the roots of maize. *J. Exp. Bot.* 36:644–651. doi:10.1093/jxb/36.4.644
- Wittwer, S.H. 1978. Carbon dioxide fertilization of crop plants. In: U.S. Gupta, editor, *Problems in crop physiology*. Haryana Agric. Univ., Hissar, India. p. 310–333.
- Wullschlegel, S.D., W.M. Post, and A.W. King. 1995. On the potential for a CO₂ fertilization effect in forests: Estimates of the biotic growth factor based on 58 controlled-exposure studies. In: A.W. Woodwell and F.T. Mackenzie, editors, *Biospheric feedbacks in the global climate system: Will warming feed the warming?* Oxford Univ. Press, Cambridge, UK. p. 85–107.
- Zhang, J., and W.J. Davies. 1987. Increased synthesis of ABA in partially dehydrated root tips and ABA transport from roots to leaves. *J. Exp. Bot.* 38:2015–2023. doi:10.1093/jxb/38.12.2015
- Zhao, X., I.J. Misaghi, and M.C. Hawes. 2000. Stimulation of border cell production in response to increased carbon dioxide levels. *Plant Physiol.* 122:181–188. doi:10.1104/pp.122.1.181