University of Nebraska - Lincoln DigitalCommons@University of Nebraska - Lincoln

Historical Materials from University of Nebraska-Lincoln Extension

Extension

1-1-2005

G05-1557 Planning Your Riparian Buffer: Design and Plant Selection

Amanda Fox University of Nebraska - Lincoln

Thomas G. Franti University of Nebraska - Lincoln, tfranti1@unl.edu

Scott J. Josiah University of Nebraska - Lincoln, sjosiah2@unl.edu

Mike Kucera Natural Resources Conservation Service

Follow this and additional works at: http://digitalcommons.unl.edu/extensionhist



Part of the Agriculture Commons, and the Curriculum and Instruction Commons

Fox, Amanda; Franti, Thomas G.; Josiah, Scott J.; and Kucera, Mike, "G05-1557 Planning Your Riparian Buffer: Design and Plant Selection" (2005). Historical Materials from University of Nebraska-Lincoln Extension. Paper 1433. http://digitalcommons.unl.edu/extensionhist/1433

This Article is brought to you for free and open access by the Extension at Digital Commons@University of Nebraska - Lincoln. It has been accepted for inclusion in Historical Materials from University of Nebraska-Lincoln Extension by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.



Planning Your Riparian Buffer: Design and Plant Selection

Learn how to plan and design a riparian buffer and select appropriate tree and grass species. A companion NebGuide, *Installing Your Riparian Buffer: Tree and Grass Planting, Postplanting Care and Maintenance* (G05-1558), addresses buffer installation, planting trees and grasses, postplanting care and long-term maintenance.

Amanda Fox, Graduate Student in Biological Systems Engineering; Tom Franti, Extension Surface Water Management Specialist; Scott Josiah, Nebraska State Forester; Mike Kucera, State Resource Conservationist for the Natural Resources Conservation Service

- Goal Identification
- Selecting the Appropriate Buffer Design
- Selecting Appropriate Plant Materials
 - o Trees and Shrubs, Seedlings or Seeds?
 - o Grasses
- Resources
- References
- Funding

Conservation buffers are planted for environmental, aesthetic, recreational, and economic reasons. Grass filter strips, grassed waterways, field borders, and field windbreaks are examples of conservation buffers. A conservation buffer also may be a streamside or riparian forest buffer and include trees, shrubs, and grasses. Riparian buffers are a *best management practice* to protect stream water quality, reduce streambank erosion, and provide wildlife habitat. Buffers also can provide income through payments from federal, state and local cost-share programs or through production and sale of specialty crops. This NebGuide provides instructions on riparian buffer planning, design and selection of tree and grass species appropriate for riparian buffers. It compliments the instructional video *Streamside Conservation: Installing and Maintaining Your Riparian Buffer*, available from University of Nebraska Cooperative Extension. (To order the video, contact the University of Nebraska CIT, P.O. Box 830918, Lincoln, NE 68583-0918 or call 800-755- 7765.)

Planning is the first step in buffer installation. First, identify your goals for the buffer, then select a design and plant materials to achieve your goals.

Goal Identification

Consider what you want to accomplish with your buffer. Do you want to protect surface water quality, enhance wildlife habitat, stabilize streambanks, and/or generate income? Buffers can be used to meet one or more of these goals.

Buffers protect surface water by intercepting runoff and irrigation water flowing from crop fields. Vegetation in the buffer slows the water, increasing infiltration and allowing sediment deposition. This allows nutrients, chemicals, and other pollutants to be removed. Riparian forest buffers also stabilize streambanks and provide shaded areas for aquatic habitat. Plant roots anchor the stream bank and help prevent erosion.

Income can be generated through land rental and maintenance payments by enrolling the buffer in the Continuous Conservation Reserve Program (CCRP) or by selling specialty crops such as nuts, fruits, and woody florals grown in the buffer. Typically, specialty products grown in buffers under a CCRP contract cannot be harvested for sale during the contract period; however, these products can be harvested for personal use. Contact your local Natural Resources Conservation Service (NRCS) or Farm Service Agency (FSA) office for more information on regulations affecting the harvest and sale of specialty crops planted in buffers.

Selecting the Appropriate Buffer Design

Select a buffer design based on your goals. In the eastern United States, riparian forest buffers provide streambank stabilization, shade streams, and absorb nutrients from shallow groundwater. These buffers typically consist of three zones: trees near the stream, then shrubs, and then 30 feet or more of grasses adjacent to the cropland.

In the Great Plains producers as well as government officials, have been reluctant to have trees planted next to streams because fallen trunks and limbs may block streams, ditches and culverts. An alternative design, with shrubs and small trees planted next to the stream followed by taller trees and then grasses next to the cropland, may be more suitable in this region (*Figure 1*).

If your main goal is to provide wildlife habitat, design your buffer accordingly. Some birds, including game species such as pheasant and prairie grouse and nongame species such as songbirds, prefer open grassland to woody cover. On the other hand, deer and sharptail grouse prefer woody edges (along narrow riparian forest buffers). Western meadowlark and mink prefer narrow (less than 35 feet wide) grass buffers. White-tailed deer, beaver, and red fox are content with just about any vegetation and width, whereas wild turkey and wood duck prefer forested buffers over




Figure 1. Cross section of an alternative buffer design.

35 feet wide. For more information on wildlife needs refer to NRCS Conservation Practice Standard *Riparian Forest Buffer* Code 391.

A buffer also can be entirely grass (also called a grass filter or filter strip). Grass buffers are narrow strips of land between the crop and stream. The NRCS recommends a minimum width of 20-30 feet for grass buffers; however, widths may vary according to your goals, cropping history, crop field area, and the program in which your buffer is enrolled (*Table 1*). For more information, refer to NRCS Conservation Practice Standard *Filter Strip* Code 393.

| Table I. NRCS conservation buffer and Continuous Conservation Reserve Program (CCRP) width guidelines. 1 | | | | | |
|---|-----------------------|---|---------------------------------|----------------------------------|--|
| | Previous Land Use | Minimum NRCS Width Requirements (feet) | CCRP Maximum Width (feet) | CCRP Grass Cover (feet) | CCRP Trees or Shrubs Cover (feet) |
| Grass Filter Strip | Cropland only | 20-30 ¹ | 120 | 120 | N.A. |
| Riparian Forest Buffer | Cropland | 35 | 180 | 20-120 ² | 35-180 |
| Riparian Forest Buffer | Marginal pasture | 35 | 180 | 20 | 35-180 |
| Riparian Herbaceous Buffer | Marginal pasture only | 20-50 ³ | 120 | 20-120 ³ | Shrub clumps may be planted. |

¹Greater minimum may be required if the ratio of minimum filter strip to drainage area is more than 1:30.

³50-foot minimum width required for wildlife purposes.

| Table II. Commonly used shrub and tree species in riparian forest buffers ¹ | | | |
|--|---|--|---|
| | Common Examples ² | Desirable Characteristics | Best Planting Location |
| Small Shrubs | Sand cherry Peking cotoneaster Elderberry Sandbar willow Streamco willow Chokecherry Gray dogwood Redosier dogwood Hansen rose Snowberry Golden currant | White flowers, black cherries Berry-like fruit Dark purple berries Flood tolerant, fast growing Flood tolerant, fast growing Edible fruit High wildlife value Floral stems, winter color Flowers, fruit High wildlife value Yellow flowers, purple fruit | Statewide Statewide All but Panhandle Statewide |
| Large Shrubs | Chokecherry American hazelnut Lilac Juneberry American plum Caragena | White flowers and cherries Hazelnuts Flowers Edible fruit White flowers, fruit, fall color Nitrogen fixing | Statewide All but Panhandle Statewide Statewide Statewide Statewide Statewide |
| Small Trees | Black cherry | Cherry wood, fruit | East |

²Grass width is considered along the outside edge of the buffer only.

| | Chickasaw plum | Drought tolerant | Statewide |
|-------------|--|---|--|
| | Amur maple | Drought tolerant, fall color | Statewide |
| Large Trees | Red mulberry Green ash Cottonwood Hackberry Black walnut Black willow Silver maple Bur oak | Edible fruit Yellow foliage in fall State tree, major wood supply Tolerant to adverse weather Wood, nuts Flood tolerant Flood tolerant Drought tolerant, acorns | Statewide Statewide Statewide Statewide East Statewide Statewide Statewide Statewide Statewide |

¹Additional species are approved by NRCS for buffer plantings. ²Species listed in bold are well suited for direct seeding methods.

| Table IV. Common grass species used in grass buffers. ¹ | | | | |
|--|---|----------------------------|---|--------------------------------|
| Recommended Mixtures | Species in the Mixture | Percent of Mixture (%) | Seeding Rate ² (PLS/ft ²) | Upland Bird Wildlife Value |
| Cool Season | Virginia wild rye Canada wild rye Western wheatgrass | 30 26 44 | 7.0 4.0 7.0 | Nesting cover |
| Warm Season | Big bluestem Switchgrass Indiangrass | 40 35 25 | 4.2 1.6 2.5 | Winter cover |
| Warm/Cool Season Mix | Switchgrass Big bluestem Intermediate wheatgrass | 60 20 20 | 2.7 2.1 4.0 | Winter cover and nesting cover |
| Warm Season | Big bluestem Switchgrass Indiangrass Sideoats grama Little bluestem | 30 25 20 15 10 | 3.2 1.1 2.0 1.4 0.7 | Winter cover and nesting cover |
| Warm/Cool Season Mix | Pubescent wheatgrass Western wheatgrass Switchgrass | 30 30 40 | 5.2 4.8 1.8 | Winter cover and nesting cover |

 $^{^{1}\}mbox{Additional species}$ are approved by NRCS for buffer plantings.

²Species listed in bold are well suited for direct seeding methods.




Figure 2. Types of seedlings.

Table III. Common specialty crop species. 1

| | Common Examples | Preferred Planting Location | |
|---------------|--|---|--|
| Nut Producing | Black walnut Chinese chestnut Northern pecan Hybrid hazelnut | East Southeast East East | |
| Woody Florals | Redstem dogwood Pussy willow Curly willow Forsythia | Statewide with irrigation Statewide with irrigation Statewide with irrigation Statewide with irrigation | |
| Fruit | Sand cherry Elderberry Nanking cherry Corneliancherry dogwood Chokecherry American plum Mulberry | Statewide East All but Panhandle East East Statewide Statewide | |

¹When these species are used to produce a marketable product, planting location is limited to sites with superior growing conditions for maximum production. Planting locations are more restrictive than those listed in *Table II*.

Selecting Appropriate Plant Materials

Trees and Shrubs, Seedlings or Seeds?

Trees and shrubs can be planted as seeds or seedlings. Tree seeds can be purchased or collected free from trees in the fall. Seeds from native trees and shrubs may be best adapted to your area's climate and soil conditions. Direct planting seeds can be cheaper than planting seedlings and usually results in a denser stand of trees and shrubs. This stand will be more like a natural forest and better able to withstand

wildlife damage, especially in early growth stages. Trees grown directly from seeds develop strong root systems that remain undisturbed throughout establishment. Seeds also can be planted in the fall. Some species suitable for Nebraska and appropriate for direct seeding are noted in *Table II*. For more information on direct seeding trees and shrubs, refer to the University of Nebraska Cooperative Extension NebGuide, *Establishing Conservation Plantings of Nut Trees and Shrubs by Direct Seeding Methods* (G03-1512).

Tree and shrub seedlings can be purchased as unrooted cuttings, bareroot plants, or in a container. Unrooted cuttings look like sticks and are cut from dormant tree branches during the winter (*Figure 2*). When planted, the underground portion forms roots and the above ground portion forms branches and leaves. Establishment of unrooted cuttings is more difficult than other seedlings and only a limited number of species, such as willows and some poplars, are suitable for this method in Nebraska. Bareroot seedlings are the most common type of seedlings used (*Figure 2*). They are grown from seed in a nursery until healthy root systems form. They are lifted without soil from the nursery beds and are transported for planting. Containerized seedlings are produced in containers. They may be shipped in the containers in which they were originally grown or removed from the containers at the nursery and packed in plastic bags (*Figure 2*). Although they are more expensive, their roots are better protected and they may be better suited for planting in dry and hard-to-establish sites.

Commonly used tree and shrub species are listed in *Table II*. Many tree and shrub species produce commercially valuable products such as decorative stems for the floral industry, fruit and nuts for the food industry, and other products for the pharmaceutical and herbal industries (*Table III*). Personnel at a local Cooperative Extension, Natural Resources Conservation Service, or Nebraska Forest Service office or commercial nurseries can help you select plants suited to your climate and soil type.

Grasses

The selection of appropriate grass species will depend on your goals. Densely planted, stiff-stemmed species should be selected to trap sediment and protect water quality. Other species may be used if wildlife habitat is a goal. Buffers planted under a CCRP contract must follow NRCS specifications for grass selection. To meet NRCS specifications, grass seed must be a mixture of at least three species adapted to the site. Warm season grass mixtures must contain at least 60 percent sod-forming stiff stem species such as big bluestem or switchgrass, and cool season grass mixtures must contain at least 40 percent sodforming stiff stem species such as western wheatgrass or Virginia wild rye. Grass seed should be planted at a minimum rate of 40 pure live seeds (PLS) per square foot. Several common warm and cool season grass mixtures appropriate for Nebraska are presented in Table IV. Wildflowers, such as coneflower and blanket flower, may be added to grass mixtures for appearance and wildlife enhancement.

Resources

For more information on installing and maintaining riparian buffers, contact: 1) your local University of Nebraska Cooperative Extension office or visit the University's Conservation Buffer Web site at http://www.conservationbuffers.unl.edu/; 2) your local Natural Resources Conservation Service office or visit the Natural Resources Conservation Service Web site, Buffer Strips: Common Sense Conservation at http://www.nrcs.usda.gov/feature/buffers; or 3) the USDA Agroforestry Center or its Riparian Forest Buffers Web site at http://www.unl.edu/nac/riparian.html.

References

Establishing Conservation Plantings of Nut Trees and Shrubs by Direct Seeding Methods, University of Nebraska Cooperative Extension NebGuide G03-1512, 2003.

Riparian Forest Buffer, Natural Resources Conservation Service Conservation Practice Standard CODE 391. Filter Strip, Natural Resources Conservation Service Conservation Practice Standard CODE 393.

Funding

Partial funding for this publication was provided by an Education Grant from the Environmental Quality Incentives Program (EQIP) through the U.S. Department of Agriculture Natural Resources Conservation Service.

File G1557 under: WATER RESOURCE MANAGEMENT

A-39, Water Quality

Issued February 2005; 2,000 printed.

Issued in furtherance of Cooperative Extension work, Acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture. Elbert C. Dickey, Director of Cooperative Extension, University of Nebraska, Institute of Agriculture and Natural Resources.

University of Nebraska Cooperative Extension educational programs abide with the non-discrimination policies of the University of Nebraska-Lincoln and the United States Department of Agriculture.