

10-25-2006

Phase stability analysis using interval Newton method with NRTL model

Hatice Gecegomez
University of Mersin

Yasar Demirel
University of Nebraska-Lincoln, ydemirel2@unl.edu

Follow this and additional works at: <https://digitalcommons.unl.edu/chemengfluidmech>



Part of the [Chemical Engineering Commons](#)

Gecegomez, Hatice and Demirel, Yasar, "Phase stability analysis using interval Newton method with NRTL model" (2006). *Papers in Fluid Mechanics*. 2.

<https://digitalcommons.unl.edu/chemengfluidmech/2>

This Article is brought to you for free and open access by the Chemical and Biomolecular Engineering Research and Publications at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Papers in Fluid Mechanics by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Phase stability analysis using interval Newton method with NRTL model

Abstract

The Gibbs energy minimization using activity coefficient models and nonlinear equation solution techniques are commonly applied for phase stability problems. However, dependence on the initial estimates and multiple solutions for these highly nonlinear equations are common drawbacks for some of the conventional approaches. We have used interval Newton method with the local composition model of NRTL for the phase stability analysis of 10 binary systems and 2 ternary systems at various feed compositions to locate all the stationary points. Results indicate that the interval Newton method is reliable and efficient.

Keywords: Phase stability; Interval Newton method; NRTL model

Comment:- This paper was published in the Journal of “Fluid Phase Equilibria” Volume 232. and Issue1- 2 on 25 October 2005 ,pages 48-45. And all the copy rights © of this paper belong to Elsevier B.V.

1. Introduction

Phase splitting due to thermodynamic instability of liquid mixtures plays important role in simulation and design problems of separation by distillation and extraction. In three-phase distillation, for example, feed may become unstable and splits into two liquid phases at some stages in the column. To know the exact number of phases on a stage contributes considerably towards the mathematical stability of phase equilibria predictions [1] and [2]. Location of a feed point with respect to the binodal curve may be used to predict the number of exact phases in a ternary mixture of types I and II; only the feed points within the binodal curve will split into two liquid phases, and require three-phase equilibrium calculations [3].

For some time, it has been a challenge to find a reliable stability analysis for fluid mixtures in any number of component and phases. Most commonly, stability analysis is based on the distance of tangent plane with respect to the Gibbs energy of mixing surface [4], [5], [6] and [7]. When the distance is negative then a mixture at specified temperature and pressure becomes unstable and splits into two liquid phases. Tangent plane distance function is the difference between the Gibbs free energy of a phase with composition x and the tangent plane to the Gibbs free energy surface estimated from a candidate phase composition z . However, the tangent plane distance approach requires the solution of nonlinear equations obtained with the activity coefficient models or the equations of state. Such solutions may lead to multiple stationary points depending of the initial values or require the number of phases known [8], and sometimes may not be reliable [7].

Interval Newton method proposed about 10 years ago is another approach for the phase stability analysis [9], [10] and [11]. The method is a generalized bisection algorithm with some modifications so that it is relatively less sensitive to initial values, and should provide all the roots including global optimum [11]. The method has been tested with equation of states and activity coefficient models [9], [12] and [13], and for process design calculations, such as mixed flow reactor and a reaction kinetics model [14]. This study further tests the reliability of phase stability analysis for 10 binary mixtures and 2

ternary mixtures using the interval Newton method and the activity coefficient model of NRTL, which might predict multiple roots in phase equilibrium calculations.

2. Theoretical framework

2.1. Phase stability analysis

The tangent plane distance D for a mixture with n number of components is expressed by:

$$D(x) = g_m(x) - g_m(z) - \sum_{i=1}^n \left(\frac{\partial g_m}{\partial x_i} \right)_z (x_i - z_i) \quad (1)$$

Where g_m is the reduced Gibbs energy of mixing, and given by:

$$g_m(x) = \sum_{i=1}^n x_i \ln x_i + g^E(x) \quad (2)$$

$$g^E = \frac{G^E}{RT} \quad (3)$$

where g^E is the reduced molar excess Gibbs energy. When the tangent plane distance D for a composition x is negative, a phase with feed mole fractions z is unstable, and the molar Gibbs energy of mixing surface $g_m = \Delta G_{\text{mix}}/RT$ falls below a plane tangent to the surface at z . The partial derivatives in Eq. (1) are calculated at $x = z$. The tangent plane distance analysis minimizes Eq. (1), and the solutions of the following system of nonlinear equations identify the stationary points [13]:

$$\left[\left(\frac{\partial g_m}{\partial x_i} \right) - \left(\frac{\partial g_m}{\partial x_n} \right) \right] - \left[\left(\frac{\partial g_m}{\partial x_i} \right) - \left(\frac{\partial g_m}{\partial x_n} \right) \right]_z = 0, \quad (4)$$

$i = 1, \dots, n - 1$

$$\sum_{i=1}^n x_i = 1 \quad (5)$$

The interval Newton/generalized bisection method requires no initial guess, and finds all the stationary points of the tangent plane distance D by solving Eqs. (4) and (5) with some easy modifications, which are explained in detail by Hua et al. [9] and [10] and Gau and Stadtherr [14].

2.2. NRTL model

For an n component system, the NRTL equation for the reduced molar excess Gibbs energy is given by:

$$g^E(x) = \sum_{k=1}^n x_k \frac{\sum_{i=1}^n G_{ik} \tau_{ik} x_i}{\sum_{i=1}^n G_{ik} x_i} \quad (6)$$

where x_k is the mole fraction for species k , and τ_{ik} and G_{ik} are the NRTL binary interaction parameters. Eq. (6) uses the mole fraction weighted averages of the interaction parameters to improve the efficiency of the interval method [10] and [14]. The parameter G_{ik} is a function of τ_{ik} and the parameter $\alpha_{ik} = \alpha_{ki}$, and is given by $G_{ik} = \exp(-\alpha_{ik}\tau_{ik})$. With the NRTL model, Eq. (4) becomes [13]

$$\ln \left(\frac{x_i}{x_n} \right) - \ln \left(\frac{z_i}{z_n} \right) + b_i(x) - b_i(z) = 0, \quad i = 1, \dots, n-1 \quad (7)$$

where

$$b_i(x) = \frac{\sum_{i=1}^{n-1} G_{ik} \tau_{ik} x_i}{\sum_{i=1}^{n-1} G_{ik} x_i} + \sum_{k=1}^n x_k \times \left[\frac{G_{ik} \tau_{ik} - G_{nk} \tau_{nk}}{\sum_{i=1}^n G_{ik} x_i} + \frac{\sum_{i=1}^n G_{ik} \tau_{ik} x_i (G_{nk} - G_{ik})}{(\sum_{i=1}^n G_{ik} x_i)^2} \right] \quad (8)$$

Eqs. (7) and (5) represent a set of $n \times n$ equation system whose solutions are the roots of D .

2.3. Interval Newton method

The interval method is a general-purpose computational method to solve nonlinear equations to find all the solutions lying within the variable bounds [12]. It uses interval vectors and matrices starting with a specified initial box of intervals, and search all the roots by solving the linear interval equation system for a new interval $N^{(k)}$:

$$F'(X^{(k)})(N^{(k)}-x^{(k)})=-f(x^{(k)}) \quad (9)$$

Where k is the iteration number, $F'(X^{(k)})$ is an interval extension of the real Jacobian of $f(x)$ over the current interval $(X^{(k)})$ and $(x^{(k)})$ is a midpoint of that interval. Tightness of the interval $(N^{(k)})$ containing the solution set mostly depends on how to solve $(N^{(k)})$. Usually, iterative methods with Gauss elimination is performed using preconditioned Eq. (9) for a tighter intervals [13] and [14]. Details of preconditioning of Eq. (9) and enhancing interval calculations are given by Schnepfer and Stadtherr [12], Tessier et al. [13] and Gau and Stadtherr [14]. A reasonable initial box should be wide enough so that the interval Newton method provides all the solutions of local minima and maxima, saddle points and global minimum.

3. Results and discussion

We used 10 binary and 2 ternary systems for the stability analysis with the reduced Gibbs energy g^E obtained from the NRTL model. The local composition model of NRTL is known to have multiple roots in phase equilibrium calculations [2]. The systems include mostly the polar–polar mixtures. In order to test the method, we have obtained all the stationary points and the values of tangent plane distance D for those roots from a MATLAB program. We verified the results of our MATLAB program with one of the binary systems used by Stadtherr and Schnepfer [15] and a ternary system used by Tessier et al. [13]. The results of point approximations [13] presented in [Table 1](#), [Table 2](#), [Table 3](#), [Table 4](#), [Table 5](#), [Table 6](#), [Table 7](#), [Table 8](#), [Table 9](#), [Table 10](#), [Table 11](#) and [Table 12](#) represent the verified enclosures which contain a unique root x . The values of the tangent plane distance D at each root have been obtained after satisfying that the deviations between the successive roots are as low as 1.00×10^{-10} . [Table 1](#), [Table 2](#),

[Table 3](#), [Table 4](#), [Table 5](#), [Table 6](#), [Table 7](#), [Table 8](#), [Table 9](#) and [Table 10](#) display the number of root inclusion tests and the solution times in seconds on a PC Genuine Intel computer, which is rather slow. The systems and the input data for the NRTL model are described in the following problems:

Problem 1

The binary parameters of polar–polar mixture *n*-pentanol(1)–2,2-dimethylbutane(2) are $\tau_{12} = 1.0114044$, $\tau_{21} = 2.9413053$, $G_{12} = 0.5527024$ and $G_{21} = 0.1782906$ [16]. [Table 1](#) shows the values of D , which are negative at the stationary points $z_1 = 0.05$ and 0.10 , and the phases at these feed points are unstable. At the feed compositions $z_1 = 0.15$ and 0.20 , the phases are stable as $D > 0$.

Problem 2

The binary parameters of polar–polar mixture *n*-pentanol(1)–2-methylpentane(2) are $\tau_{12} = 0.9598699$, $\tau_{21} = 2.9389421$, $G_{12} = 0.5746116$ and $G_{21} = 0.1833367$ [16]. As the values of D in [Table 2](#) show, the feed composition of $z_1 = 0.05$, 0.20 and 0.25 are stable, while the feeds at $z_1 = 0.10$, 0.11 and 0.12 are unstable.

Problem 3

The binary parameters of polar–apolar mixture ethanol(1)–cyclohexane(2) are $\tau_{12} = 1.7252352$, $\tau_{21} = 3.1963108$, $G_{12} = 0.4541860$ and $G_{21} = 0.2317199$ [16]. The values of D in [Table 3](#) show that the feeds at $z_1 = 0.10$, 0.15 , 0.20 and 0.25 are unstable, while the feed compositions of $z_1 = 0.45$, 0.50 , 0.60 and 0.65 are stable.

Problem 4

Water (1)–butyl glycol (2) mixture is a polar–polar mixture with the binary parameters of $\tau_{12} = 1.2005955$, $\tau_{21} = 1.4859846$, $G_{12} = 0.8644289$ and $G_{21} = 0.8350056$ [16]. The values of D in [Table 4](#) show that the feed points at $z_1 = 0.05$ and 0.10 are stable, while the feed points at $z_1 = 0.30$, 0.35 , 0.40 , 0.45 , 0.50 , 0.60 and 0.65 are unstable.

Problem 5

The binary parameters of polar–polar mixture water(1)–citric acid(2) are $\tau_{12} = 0.9889317$, $\tau_{21} = 13.7521382$, $G_{12} = 0.6887706$ and $G_{21} = 5.6008823 \times 10^{-3}$ [17]. As seen in [Table 5](#), the values of D at the stationary points are positive and the feeds are stable.

Problem 6

The binary parameters of polar–polar mixture citric acid(1)–1-butanol(2) are $\tau_{12} = 2.5479457 \times 10^{-2}$, $\tau_{21} = 11.2949857$, $G_{12} = 0.9948420$ and $G_{21} = 0.1010207$ [17]. As [Table 6](#) shows, the values of D at the feed points $z_1 = 0.05, 0.10, 0.15, 0.20$ and 0.25 are negative leading to phase instability. At the feed compositions $z_1 = 0.65$ and 0.75 , the system is stable with positive values of D .

Problem 7

The binary parameters of polar–polar mixture citric acid(1)–2-butanol(2) are $\tau_{12} = -1.3581754$, $\tau_{21} = 7.4341774$, $G_{12} = 1.3199010$ and $G_{21} = 0.2188763$ [17]. As [Table 7](#) shows, the feed points of $z_1 = 0.05, 0.10, 0.15, 0.20$ and 0.25 are unstable. At the feed compositions $z_1 = 0.50, 0.60, 0.65$ and 0.75 the system is stable with positive values of D .

Problem 8

Water(1)–1,4-dicyanobutane(2) mixture is a polar–polar system, with the binary parameters of $\tau_{12} = 4.6707725$, $\tau_{21} = 0.6918617$, $G_{12} = 0.2462934$ and $G_{21} = 0.8125657$ [18]. The interval Newton method is successfully obtained 38 stationary points, which are shown in [Table 8](#). At the feed compositions $z_1 = 0.05, 0.10, 0.15, 0.20$ and 0.25 the system is stable with positive values of D . At the feed compositions $z_1 = 0.60, 0.65$ and 0.75 the system is unstable with negative values of D .

Problem 9

The binary parameters of polar–polar mixture water(1)–butanenitrile(2) are $\tau_{12} = 4.9011241$, $\tau_{21} = 1.8395856$, $G_{12} = 0.2298480$ and $G_{21} = 0.5758687$ [18]. [Table 9](#) shows that the feed point at $z_1 = 0.05$ is stable, while the feed points at $z_1 = 0.40, 0.45, 0.50$ and 0.75 are unstable.

Problem 10

The binary parameters of polar–polar mixture water(1)–benzonitrile(2) are $\tau_{12} = 6.7197320$, $\tau_{21} = 2.7710776$, $G_{12} = 0.1331979$ and $G_{21} = 0.4354727$ [18]. [Table 10](#) shows that most of the feed points are unstable.

Problem 11

The ternary parameters for acetonitrile(1)–benzene(2)–*n*-heptane(3) mixture are $\tau_{12} = 0.5661821$, $\tau_{21} = 0.4472257$, $\tau_{23} = 1.4918912$, $\tau_{32} = -0.5982783$, $\tau_{13} = 2.3187177$, $\tau_{31} = 0.6964173$, $G_{12} = 0.9175650$, $G_{21} = 0.9343016$, $G_{23} = 0.6290071$, $G_{32} = 1.2043230$, $G_{13} = 0.7589032$ and $G_{31} = 0.9204803$ [16]. [Table 11](#) shows that the feed compositions are unstable.

Problem 12

The ternary parameters for water(1)–citric acid(2)–2-butanol(3) mixture are $\tau_{13} = 2.9732685$, $\tau_{31} = 0.5249036$, $G_{13} = 0.2472300$ and $G_{31} = 0.7813714$ [17]. The values of τ_{12} , τ_{21} , G_{12} and G_{21} are the same with those given in [Problem 5](#), while the values of τ_{23} , τ_{32} , G_{23} and G_{32} are given in [Problem 7](#). [Table 12](#) shows the values of D for stable and unstable feed compositions.

Table 1.

Stationary points for *n*-pentanol(1)–2,2-dimethylbutane(2) system of [Problem 1](#) at various feed points at 25 °C and 1 atm

Feed (z_1, z_2)	Stationary points (x_1, x_2)	D	Root inclusion tests	Solution time (s)
(0.05, 0.95)	(0.0500, 0.9500)	0.0000	399	38
	(0.1294, 0.8706)	-1.8495×10^{-3}		
(0.10, 0.90)	(6.9187×10^{-2} , 0.9308)	-4.7473×10^{-5}	217	26
	(0.1000, 0.9000)	0.0000		
	(0.1500, 0.8500)	-9.8214×10^{-5}		
(0.15, 0.85)	(6.9168×10^{-2} , 0.9308)	5.0013×10^{-5}	217	26
	(0.1000, 0.9000)	9.7765×10^{-5}		
	(0.1500, 0.8500)	0.0000		
(0.20, 0.80)	(6.5086×10^{-2} , 0.9349)	2.9150×10^{-3}	209	25
	(0.2000, 0.8000)	0.0000		

Table 2.

Stationary points for *n*-pentanol(1)–2-methylpentane(2) system of [Problem 2](#) at various feed points at 25 °C and 1 atm

Feed (z_1, z_2)	Stationary points (x_1, x_2)	D	Root inclusion tests	Solution time (s)
(0.05, 0.95)	(0.0500, 0.9500)	0.0000	394	38
	(0.1365, 0.8635)	2.1266×10^{-3}		
(0.10, 0.90)	(6.8845×10^{-2} , 0.9311)	-5.7673×10^{-5}	212	26
	(0.1000, 0.9000)	0.0000		
	(0.1657, 0.8343)	-2.2122×10^{-4}		
(0.11, 0.89)	(6.5423×10^{-2} , 0.9346)	-1.4238×10^{-4}	222	26
	(0.1100, 0.8900)	0.0000		
	(0.1582, 0.8418)	-9.3395×10^{-5}		
(0.12, 0.88)	(6.3345×10^{-2} , 0.9367)	-2.3299×10^{-4}	238	27
	(0.1200, 0.8800)	0.0000		
	(0.1495, 0.8505)	-2.2301×10^{-5}		
(0.20, 0.80)	(7.0986×10^{-2} , 0.9290)	2.0998×10^{-3}	222	26
	(0.2000, 0.8000)	0.0000		
(0.25, 0.75)	(5.5296×10^{-2} , 0.9447)	9.3157×10^{-3}	218	26
	(0.2500, 0.7500)	0.0000		

Table 3.

Stationary points for ethanol(1)–cyclohexane(2) system of [Problem 3](#) at various feed points at 5 °C and 1 atm

Feed (z_1, z_2)	Stationary points (x_1, x_2)	D	Root inclusion tests	Solution time (s)
(0.05, 0.95)	(0.0500, 0.9500)	0.0000	202	25
	(9.4213×10^{-2} , 0.9058)	6.2615×10^{-4}		
	(0.5979, 0.4021)	-5.1329×10^{-2}		
(0.10, 0.90)	(4.7514×10^{-2} , 0.9525)	-1.0107×10^{-3}	187	23
	(0.1000, 0.9000)	0.0000		
	(0.5878, 0.4122)	-4.8019×10^{-2}		
(0.15, 0.85)	(3.5205×10^{-2} , 0.9648)	-7.0829×10^{-3}	173	22
	(0.1500, 0.8500)	0.0000		
	(0.4879, 0.5121)	-1.8081×10^{-2}		
	(0.9815, 1.8550×10^{-2})	-2.2408×10^{-3}		
(0.20, 0.80)	(3.0362×10^{-2} , 0.9696)	-1.4273×10^{-2}	170	22
	(0.2000, 0.8000)	0.0000		
	(0.4000, 0.6000)	-4.0523×10^{-3}		
(0.25, 0.75)	(2.8470×10^{-2} , 0.9715)	-1.9142×10^{-2}	191	24
	(0.2500, 0.7500)	0.0000		
	(0.3314, 0.6686)	-2.8350×10^{-4}		
(0.35, 0.60)	(2.8832×10^{-2} , 0.9712)	-1.7255×10^{-2}	197	24
	(0.2351, 0.7649)	7.9166×10^{-4}		

Feed (z_1, z_2)	Stationary points (x_1, x_2)	D	Root inclusion tests	Solution time (s)
	(0.3500, 0.6500)	0.0000		
(0.40, 0.55)	$(3.0363 \times 10^{-2}, 0.9696)$	-1.0218×10^{-2}	160	21
	(0.2000, 0.8000)	4.0543×10^{-3}		
	(0.4000, 0.6000)	0.0000		
(0.45, 0.60)	$(3.2732 \times 10^{-2}, 0.9673)$	5.4846×10^{-4}	179	22
	(0.1702, 0.8298)	1.0652×10^{-2}		
	(0.4500, 0.5500)	0.0000		
	$(0.9875, 1.2543 \times 10^{-2})$	2.8205×10^{-2}		
(0.50, 0.50)	$(3.6146 \times 10^{-2}, 0.9638)$	1.4701×10^{-2}	174	22
	(0.1439, 0.8561)	2.0885×10^{-2}		
	(0.5000, 0.5000)	0.0000		
	$(0.9794, 2.0600 \times 10^{-2})$	1.1910×10^{-2}		
(0.60, 0.40)	$(5.0600 \times 10^{-2}, 0.9495)$	5.2247×10^{-2}	196	24
	$(9.2933 \times 10^{-2}, 0.9071)$	5.2801×10^{-2}		
	(0.6000, 0.4000)	0.0000		
(0.65, 0.35)	$(6.4067 \times 10^{-2}, 0.9359)$	7.5606×10^{-2}	285	32
	(0.6500, 0.3500)	0.0000		
(0.75, 0.25)	$(4.9309 \times 10^{-2}, 0.9507)$	0.1409	204	25
	(0.7500, 0.2500)	0.0000		
	$(0.9339, 6.6045 \times 10^{-2})$	-5.6707×10^{-2}		

Table 4.Stationary points for water(1)–butyl glycol(2) system of [Problem 4](#) at various feed points at 5 °C and 1 atm

Feed (z_1, z_2)	Stationary points (x_1, x_2)	D	Root inclusion tests	Solution time (s)
(0.05, 0.95)	(0.0500, 0.9500)	0.0000	265	30
	(0.8403, 0.1597)	0.4708		
(0.10, 0.90)	(0.1000, 0.9000)	0.0000	404	39
	(0.7234, 0.2766)	0.1255		
(0.15, 0.85)	(0.1500, 0.8500)	0.0000	178	22
	(0.4768, 0.5232)	2.6938×10^{-2}		
	(0.8439, 0.1561)	-7.7671×10^{-3}		
(0.20, 0.80)	(0.2000, 0.8000)	0.0000	183	23
	(0.3656, 0.6344)	3.9322×10^{-3}		
	(0.8709, 0.1291)	-7.0380×10^{-2}		
(0.25, 0.75)	(0.2500, 0.7500)	0.0000	213	26
	(0.2956, 0.7044)	8.4999×10^{-5}		
	(0.8786, 0.1214)	-9.2025×10^{-2}		
(0.30, 0.70)	(0.2463, 0.7537)	-1.3869×10^{-4}	199	24
	(0.3000, 0.7000)	0.0000		
	(0.8783, 0.1217)	-9.1484×10^{-2}		
(0.35, 0.65)	(0.2096, 0.7904)	-2.4208×10^{-3}	186	23

Feed (z_1, z_2)	Stationary points (x_1, x_2)	D	Root inclusion tests	Solution time (s)
	(0.3500, 0.6500)	0.0000		
	(0.8733, 0.1267)	-7.9379×10^{-2}		
(0.40, 0.60)	(0.1815, 0.8185)	-8.7794×10^{-3}	182	23
	(0.4000, 0.6000)	0.0000		
	(0.8644, 0.1356)	-6.2151×10^{-2}		
(0.45, 0.55)	(0.1597, 0.8403)	-1.9546×10^{-2}	190	24
	(0.4500, 0.5500)	0.0000		
	(0.8520, 0.1480)	-4.3912×10^{-2}		
(0.50, 0.50)	(0.1426, 0.8574)	-3.4091×10^{-2}	173	22
	(0.5000, 0.5000)	0.0000		
	(0.8358, 0.1642)	-2.7355×10^{-2}		
(0.60, 0.40)	(0.1198, 0.8802)	-6.8384×10^{-2}	184	23
	(0.6000, 0.4000)	0.0000		
	(0.7895, 0.2105)	-5.3761×10^{-3}		
(0.65, 0.35)	(0.1134, 0.8866)	-8.3043×10^{-2}	192	24
	(0.6500, 0.3500)	0.0000		
	(0.7568, 0.2432)	-9.8759×10^{-4}		
(0.75, 0.25)	(0.1126, 0.8874)	-8.4479×10^{-2}	192	24
	(0.6590, 0.3410)	6.1269×10^{-4}		
	(0.7500, 0.2500)	0.0000		

Table 5.Stationary points for water(1)–citric acid(2) system of [Problem 5](#) at various feed points at 25 °C and 1 atm

Feed (z_1, z_2)	Stationary points (x_1, x_2)	D	Root inclusion tests	Solution time (s)
(0.05, 0.95)	$(1.0335 \times 10^{-2}, 0.9897)$	9.4569×10^{-3}	333	35
	(0.0500, 0.9500)	0.0000		
(0.10, 0.90)	$(7.1546 \times 10^{-3}, 0.9929)$	4.7783×10^{-2}	275	31
	(0.1000, 0.9000)	0.0000		
(0.15, 0.85)	$(5.9859 \times 10^{-3}, 0.9940)$	9.3235×10^{-2}	218	26
	(0.1500, 0.8500)	0.0000		
(0.20, 0.80)	$(5.3287 \times 10^{-3}, 0.9947)$	0.1412	180	22
	(0.2000, 0.8000)	0.0000		
(0.25, 0.75)	$(4.8886 \times 10^{-3}, 0.9951)$	0.1906	175	22
	(0.2500, 0.7500)	0.0000		
(0.30, 0.70)	$(4.5635 \times 10^{-3}, 0.9954)$	0.2413	172	22
	(0.3000, 0.7000)	0.0000		
(0.35, 0.65)	$(3.6100 \times 10^{-7}, 0.9999)$	0.2829	163	21
	$(4.3076 \times 10^{-3}, 0.9957)$	0.2933		
	(0.3500, 0.6500)	0.0000		
(0.40, 0.60)	$(4.1700 \times 10^{-7}, 0.9999)$	0.3371	160	21
	$(4.0963 \times 10^{-3}, 0.9959)$	0.3469		

Feed (z_1, z_2)	Stationary points (x_1, x_2)	D	Root inclusion tests	Solution time (s)
	(0.4000, 0.6000)	0.0000		
(0.45, 0.55)	(4.7600×10^{-7} , 0.9999)	0.3934	161	21
	(3.9150×10^{-3} , 0.9961)	0.4026		
	(0.4500, 0.5500)	0.0000		
(0.50, 0.50)	(5.4000×10^{-7} , 0.9999)	0.4525	153	21
	(3.7541×10^{-3} , 0.9963)	0.4612		
	(0.5000, 0.5000)	0.0000		
(0.60, 0.40)	(6.8700×10^{-7} , 0.9999)	0.5845	150	21
	(3.4665×10^{-3} , 0.9965)	0.5923		
	(0.6000, 0.4000)	0.0000		
(0.65, 0.35)	(7.7700×10^{-7} , 0.9999)	0.6618	159	21
	(3.3288×10^{-3} , 0.9967)	0.6692		
	(0.6500, 0.3500)	0.0000		
(0.75, 0.25)	(1.0330×10^{-6} , 0.9999)	0.8605	159	21
	(3.0361×10^{-3} , 0.9970)	0.8670		
	(0.7500, 0.2500)	0.0000		

Table 6.

Stationary points for citric acid(1)–1-butanol(2) system of [Problem 6](#) at various feed points at 25 °C and 1 atm

Feed (z_1, z_2)	Stationary points (x_1, x_2)	D	Root inclusion tests	Solution time (s)
(0.05, 0.95)	$(7.6637 \times 10^{-5}, 0.9999)$	-8.2724×10^{-2}	237	27
	(0.0500, 0.9500)	0.0000		
	$(0.9516, 4.8386 \times 10^{-2})$	-1.9587		
(0.10, 0.90)	$(1.3061 \times 10^{-5}, 0.9999)$	-0.2079	181	23
	(0.1000, 0.9000)	0.0000		
	(0.7585, 0.2415)	-0.5434		
(0.15, 0.85)	$(6.1420 \times 10^{-6}, 0.9999)$	-0.2995	168	22
	(0.1500, 0.8500)	0.0000		
	(0.5596, 0.4404)	-0.1325		
(0.20, 0.80)	$(4.4010 \times 10^{-6}, 0.9999)$	-0.3567	173	22
	(0.2000, 0.8000)	0.0000		
	(0.4270, 0.5730)	-2.4001×10^{-2}		
(0.25, 0.75)	$(3.8540 \times 10^{-6}, 0.9999)$	-0.3859	190	24
	(0.2500, 0.7500)	0.0000		
	(0.3422, 0.6578)	-1.6667×10^{-3}		
(0.30, 0.70)	$(3.7530 \times 10^{-6}, 0.9999)$	-0.3928	237	27
	(0.2852, 0.7148)	7.0149×10^{-6}		
	(0.3000, 0.7000)	0.0000		

Feed (z_1, z_2)	Stationary points (x_1, x_2)	D	Root inclusion tests	Solution time (s)
(0.35, 0.65)	(3.8870×10^{-6} , 0.9999)	-0.3813	188	23
	(0.2444, 0.7556)	2.4989×10^{-3}		
	(0.3500, 0.6500)	0.0000		
(0.40, 0.60)	(4.1820×10^{-6} , 0.9999)	-0.3537	175	22
	(0.2137, 0.7863)	1.3448×10^{-2}		
	(0.40000, 0.6000)	0.0000		
(0.45, 0.55)	(4.6210×10^{-6} , 0.9999)	-0.3113	172	22
	(0.1895, 0.8105)	3.5874×10^{-2}		
	(0.4500, 0.5500)	0.0000		
(0.50, 0.50)	(5.2080×10^{-6} , 0.9999)	-0.2544	170	22
	(0.1697, 0.8303)	7.1332×10^{-2}		
	(0.5000, 0.5000)	0.0000		
(0.60, 0.40)	(6.9620×10^{-6} , 0.9999)	-9.4428×10^{-2}	173	22
	(0.1384, 0.8616)	0.1871		
	(0.6000, 0.4000)	0.0000		
(0.65, 0.35)	(8.2660×10^{-6} , 0.9999)	1.2754×10^{-2}	174	22
	(0.1253, 0.8747)	0.2717		
	(0.6500, 0.3500)	0.0000		
(0.75, 0.25)	(1.2528×10^{-5} , 0.9999)	0.3041	182	23

Feed (z_1, z_2)	Stationary points (x_1, x_2)	D	Root inclusion tests	Solution time (s)
	(0.1019, 0.8981)	0.5162		
	(0.7500, 0.2500)	0.0000		

Table 7.

Stationary points for citric acid(1)–2-butanol(2) system of [Problem 7](#) at various feed points at 25 °C and 1 atm

Feed (z_1, z_2)	Stationary points (x_1, x_2)	D	Root inclusion tests	Solution time (s)
(0.05, 0.95)	(7.0315×10^{-3} , 0.9930)	-9.6957×10^{-3}	201	25
	(0.0500, 0.9500)	0.0000		
	(0.7096, 0.2904)	-0.6614		
(0.10, 0.90)	(2.6555×10^{-3} , 0.9973)	-5.9344×10^{-2}	181	23
	(0.1000, 0.9000)	0.0000		
	(0.5705, 0.4295)	-0.2535		
(0.15, 0.85)	(1.5716×10^{-3} , 0.9984)	-0.1147	168	22
	(0.1500, 0.8500)	0.0000		
	(0.4558, 0.5442)	-7.3246×10^{-2}		
(0.20, 0.80)	(1.1991×10^{-3} , 0.9988)	-0.1570	175	22
	(0.2000, 0.8000)	0.0000		
	(0.3695, 0.6305)	-1.2904×10^{-2}		
(0.25, 0.75)	(1.0739×10^{-3} , 0.9989)	-0.1795	188	23
	(0.2500, 0.7500)	0.0000		

Feed (z_1, z_2)	Stationary points (x_1, x_2)	D	Root inclusion tests	Solution time (s)
	(0.3046, 0.6956)	-4.3191×10^{-4}		
(0.30, 0.70)	(1.0701×10^{-3} , 0.9989)	-0.1800	197	24
	(0.2539, 0.7461)	2.6409×10^{-4}		
	(0.3000, 0.7000)	0.0000		
(0.35, 0.65)	(1.1485×10^{-3} , 0.9989)	-0.1585	175	22
	(0.2136, 0.7864)	6.7593×10^{-3}		
	(0.3500, 0.6500)	0.0000		
(0.40, 0.60)	(1.3013×10^{-3} , 0.9987)	-0.1151	168	22
	(0.1806, 0.8194)	2.7671×10^{-2}		
	(0.4000, 0.6000)	0.0000		
(0.45, 0.55)	(1.5380×10^{-3} , 0.9985)	-5.0293×10^{-2}	167	22
	(0.1529, 0.8471)	6.7348×10^{-2}		
	(0.4500, 0.5500)	0.0000		
(0.50, 0.50)	(1.8835×10^{-3} , 0.9981)	3.5767×10^{-2}	194	24
	(0.1291, 0.8709)	0.1282		
	(0.5000, 0.5000)	0.0000		
(0.60, 0.40)	(3.1335×10^{-3} , 0.9969)	0.2739	185	23
	(8.8965×10^{-2} , 0.9110)	0.3207		
	(0.6000, 0.4000)	0.0000		

Feed (z_1, z_2)	Stationary points (x_1, x_2)	D	Root inclusion tests	Solution time (s)
(0.65, 0.35)	$(4.3155 \times 10^{-3}, 0.9957)$	0.4291	178	22
	$(7.1117 \times 10^{-2}, 0.9289)$	0.4569		
	(0.6500, 0.3500)	0.0000		
(0.75, 0.25)	$(1.1696 \times 10^{-2}, 0.9883)$	0.8308	229	26
	$(3.3412 \times 10^{-2}, 0.9666)$	0.8323		
	(0.7500, 0.2500)	0.0000		

Table 8.

Stationary points for water(1)–1,4-dicyanobutane(2) system of [Problem 8](#) at various feed points at 25 °C and 1 atm

Feed (z_1, z_2)	Stationary points (x_1, x_2)	D	Root inclusion tests	Solution time (s)
(0.05, 0.95)	(0.0500, 0.9500)	0.0000	299	33
	$(0.9828, 1.7225 \times 10^{-2})$	1.2358		
(0.10, 0.90)	(0.1000, 0.9000)	0.0000	254	29
	$(0.9294, 7.0586 \times 10^{-2})$	0.6703		
	$(0.9822, 1.7844 \times 10^{-2})$	0.6648		
(0.15, 0.85)	(0.1500, 0.8500)	0.0000	223	26
	(0.8787, 0.1213)	0.3997		
	$(0.9904, 9.5702 \times 10^{-3})$	0.3657		
(0.20, 0.80)	(0.2000, 0.8000)	0.0000	205	25
	(0.8381, 0.1619)	0.2425		
	$(0.9931, 6.9371 \times 10^{-3})$	0.1780		

Feed (z_1, z_2)	Stationary points (x_1, x_2)	D	Root inclusion tests	Solution time (s)
(0.25, 0.75)	(0.2500, 0.7500)	0.0000	195	24
	(0.8005, 0.1995)	0.1441		
	(0.9944, 5.6094×10^{-3})	5.0924×10^{-2}		
(0.30, 0.70)	(0.3000, 0.7000)	0.0000	188	23
	(0.7638, 0.2362)	8.1133×10^{-2}		
	(0.9952, 4.8177×10^{-3})	-3.8166×10^{-2}		
(0.35, 0.65)	(0.3500, 0.6500)	0.0000	183	23
	(0.7267, 0.2733)	4.1516×10^{-2}		
	(0.9957, 4.3056×10^{-3})	-0.1012		
(0.40, 0.60)	(0.4000, 0.6000)	0.0000	183	23
	(0.6882, 0.3118)	1.7966×10^{-2}		
	(0.9960, 3.9635×10^{-3})	-0.1449		
(0.45, 0.55)	(0.4500, 0.5500)	0.0000	205	25
	(0.6475, 0.3525)	5.6431×10^{-3}		
	(0.9963, 3.7390×10^{-3})	-1.7361		
(0.50, 0.50)	(0.5000, 0.5000)	0.0000	217	26
	(0.6037, 0.3963)	8.0606×10^{-4}		
	(0.9964, 3.6072×10^{-3})	-0.1899		
(0.60, 0.40)	(0.5041, 0.4959)	-6.3609×10^{-4}	219	26

Feed (z_1, z_2)	Stationary points (x_1, x_2)	D	Root inclusion tests	Solution time (s)
	(0.6000, 0.4000)	0.0000		
	(0.9964, 3.6001×10^{-3})	-0.1914		
(0.65, 0.35)	(0.4470, 0.5530)	-6.1343×10^{-3}	205	25
	(0.6500, 0.3500)	0.0000		
	(0.9963, 3.7496×10^{-3})	-0.1784		
(0.75, 0.25)	(0.3187, 0.6813)	-6.4009×10^{-2}	188	23
	(0.7500, 0.2500)	0.0000		
	(0.9954, 4.6007×10^{-3})	-0.1284		

Table 9.

Stationary points for water(1)–butanenitrile(2) system of [Problem 9](#) at various feed points at 25 °C and 1 atm

Feed (z_1, z_2)	Stationary points (x_1, x_2)	D	Root inclusion tests	Solution time (s)
(0.05, 0.95)	(0.0500, 0.9500)	0.0000	255	29
	(0.2423, 0.7577)	0.2398		
	(0.8199, 0.1801)	0.4452		
	(0.9958, 4.1846×10^{-3})	0.3478		
(0.10, 0.90)	(0.1000, 0.9000)	0.0000	227	26
	(0.3612, 0.6388)	0.1374		
	(0.6923, 0.3077)	0.1369		
	(0.9975, 2.5029×10^{-3})	-6.3535×10^{-2}		
(0.15, 0.85)	(0.1500, 0.8500)	0.0000	213	26
	(0.5810, 0.4190)	4.2080×10^{-2}		

Feed (z_1, z_2)	Stationary points (x_1, x_2)	D	Root inclusion tests	Solution time (s)
	(0.9979, 2.0467×10^{-3})	-0.2230		
(0.20, 0.80)	(0.2000, 0.8000)	0.0000	199	24
	(0.4835, 0.5165)	1.0652×10^{-2}		
	(0.9981, 1.8634×10^{-3})	-0.2942		
(0.25, 0.75)	(0.2500, 0.7500)	0.0000	207	25
	(0.4034, 0.5966)	1.6082×10^{-3}		
	(0.9982, 1.7843×10^{-3})	-0.3253		
(0.30, 0.70)	(0.3000, 0.7000)	0.0000	239	27
	(0.3407, 0.6593)	2.9584×10^{-5}		
	(0.9982, 1.7564×10^{-3})	-0.3359		
(0.40, 0.60)	(0.2524, 0.7476)	-1.4309×10^{-3}	212	26
	(0.4000, 0.6000)	0.0000		
	(0.9982, 1.7821×10^{-3})	-0.3276		
(0.45, 0.55)	(0.2195, 0.7805)	-5.5825×10^{-3}	209	25
	(0.4500, 0.5500)	0.0000		
	(0.9982, 1.8242×10^{-3})	-0.3153		
(0.50, 0.50)	(0.1909, 0.8091)	-1.4010×10^{-2}	204	25
	(0.5000, 0.5000)	0.0000		
	(0.9981, 1.8865×10^{-3})	-0.2991		

Feed (z_1, z_2)	Stationary points (x_1, x_2)	D	Root inclusion tests	Solution time (s)
(0.60, 0.40)	(0.1411, 0.8589)	-5.2405×10^{-2}	209	25
	(0.4099, 0.5901)	4.2647×10^{-2}		
	(0.6000, 0.4000)	0.0000		
	(0.9979, 2.0994×10^{-3})	-0.2557		
(0.65, 0.35)	(0.1184, 0.8816)	-8.9759×10^{-2}	225	26
	(0.3857, 0.6143)	2.5465×10^{-2}		
	(0.6500, 0.3500)	0.0000		
	(0.9977, 2.2805×10^{-3})	-0.2278		
(0.75, 0.25)	(7.6194×10^{-2} , 0.9238)	-0.2366	251	29
	(0.3185, 0.6815)	-5.9660×10^{-2}		
	(0.7500, 0.2500)	0.0000		
	(0.9970, 2.9876×10^{-3})	-0.1575		

Table 10.

Stationary points for water(1)–benzonitrile(2) system of [Problem 10](#) at various feed points at 25 °C and 1 atm

Feed (z_1, z_2)	Stationary points (x_1, x_2)	D	Root inclusion tests	Solution time (s)
(0.05, 0.95)	(0.0500, 0.9500)	0.0000	266	30
	(0.7513, 0.2487)	0.1205		
	(0.9997, 3.2783×10^{-4})	-9.2920×10^{-2}		
(0.10, 0.90)	(0.1000, 0.9000)	0.0000	264	30
	(0.2383, 0.7617)	-3.5325×10^{-3}		
	(0.9997, 2.5340×10^{-4})	-0.3258		

Feed (z_1, z_2)	Stationary points (x_1, x_2)	D	Root inclusion tests	Solution time (s)
(0.20, 0.80)	(0.1156, 0.8844)	-9.0701×10^{-4}	266	30
	(0.2000, 0.8000)	0.0000		
	(0.9997, 2.4742×10^{-4})	-0.3475		
(0.25, 0.75)	(9.6405×10^{-2} , 0.9036)	-4.6336×10^{-3}	242	28
	(0.2500, 0.7500)	0.0000		
	(0.9997, 2.5537×10^{-4})	-0.3236		
(0.30, 0.70)	(8.5011×10^{-2} , 0.9150)	-1.0332×10^{-2}	225	26
	(0.3000, 0.7000)	0.0000		
	(0.9997, 2.6362×10^{-4})	-0.3012		
(0.35, 0.65)	(7.7947×10^{-2} , 0.9221)	-1.6598×10^{-2}	212	26
	(0.3500, 0.6500)	0.0000		
	(0.9997, 2.7071×10^{-4})	-0.2837		
(0.40, 0.60)	(7.3539×10^{-2} , 0.9265)	-2.2359×10^{-2}	231	27
	(0.4000, 0.6000)	0.0000		
	(0.9997, 2.7614×10^{-4})	-0.2716		
(0.45, 0.55)	(7.0822×10^{-2} , 0.9292)	-2.7046×10^{-2}	221	26
	(0.4500, 0.5500)	0.0000		
	(0.9997, 2.7996×10^{-4})	-0.2639		
(0.50, 0.50)	(6.9111×10^{-2} , 0.9309)	-3.0688×10^{-2}	217	26

Feed (z_1, z_2)	Stationary points (x_1, x_2)	D	Root inclusion tests	Solution time (s)
	(0.5000, 0.5000)	0.0000		
	(0.9997, 2.8257×10^{-4})	-0.2592		
(0.60, 0.40)	(6.6090×10^{-2} , 0.9339)	-3.9061×10^{-2}	217	26
	(0.6000, 0.4000)	0.0000		
	(0.9997, 2.8762×10^{-4})	-0.2515		
(0.65, 0.35)	(6.3181×10^{-2} , 0.9368)	-4.9320×10^{-2}	220	26
	(0.6500, 0.3500)	0.0000		
	(0.9997, 2.9306×10^{-4})	-0.2447		
(0.75, 0.25)	(5.0253×10^{-2} , 0.9497)	-0.1187	253	29
	(0.7500, 0.2500)	0.0000		
	(0.9997, 3.2696×10^{-4})	-0.2141		

Table 11.

Stationary points for acetonitrile(1)–benzene(2)–*n*-heptane(3) system of [Problem 11](#) at various feed points at 45 °C and 1 atm

Feed (z_1, z_2, z_3)	Stationary points (x_1, x_2, x_3)	D
(0.40, 0.05, 0.55)	(0.2215, 4.8013×10^{-2} , 0.7304)	-5.1389×10^{-3}
	(0.4000, 0.0500, 0.5500)	0.0000
(0.45, 0.05, 0.50)	(0.1919, 4.7275×10^{-2} , 0.7608)	-1.5255×10^{-2}
	(0.4500, 0.0500, 0.5000)	0.0000

Feed (z_1, z_2, z_3)	Stationary points (x_1, x_2, x_3)	D
(0.60, 0.05, 0.35)	(0.1320, 4.6723×10^{-2} , 0.8213)	-8.0816×10^{-2}
	(0.6000, 0.0500, 0.3500)	0.0000
(0.70, 0.05, 0.25)	(0.1118, 4.9260×10^{-2} , 0.8389)	-0.1334
	(0.7000, 0.0500, 0.2500)	0.0000
(0.50, 0.10, 0.40)	(0.1720, 9.5257×10^{-2} , 0.7327)	-2.8631×10^{-2}
	(0.5000, 0.1000, 0.4000)	0.0000
(0.55, 0.10, 0.35)	(0.1536, 9.5606×10^{-2} , 0.7508)	-4.7718×10^{-2}
	(0.5500, 0.1000, 0.3500)	0.0000
(0.65, 0.10, 0.25)	(0.1309, 0.1006, 0.7685)	-8.6916×10^{-2}
	(0.6500, 0.1000, 0.2500)	0.0000
(0.45, 0.15, 0.40)	(0.2023, 0.1460, 0.6518)	-1.1811×10^{-2}
	(0.4500, 0.1500, 0.4000)	0.0000
(0.50, 0.15, 0.35)	(0.1805, 0.1465, 0.6731)	-2.3794×10^{-2}
	(0.5000, 0.1500, 0.3500)	0.0000
(0.60, 0.15, 0.25)	(0.1548, 0.1538, 0.6914)	-5.0477×10^{-2}
	(0.6000, 0.1500, 0.2500)	0.0000
(0.45, 0.20, 0.35)	(0.2154, 0.1989, 0.5857)	-8.6429×10^{-3}
	(0.4500, 0.2000, 0.3500)	0.0000

Feed (z_1, z_2, z_3)	Stationary points (x_1, x_2, x_3)	D
(0.55, 0.20, 0.25)	(0.1863, 0.2085, 0.6051)	-2.4032×10^{-2}
	(0.5500, 0.2000, 0.2500)	0.0000

Table 12.

Stationary points for water(1)–citric acid(2)–2-butanol(3) system of [Problem 12](#) at various feed points at 25 °C and 1 atm

Feed (z_1, z_2, z_3)	Stationary points (x_1, x_2, x_3)	D
(0.10, 0.05, 0.85)	(6.9754×10^{-2} , 1.0768×10^{-2} , 0.9195)	-4.4700×10^{-3}
	(0.1000, 0.0500, 0.8500)	0.0000
(0.20, 0.05, 0.75)	(0.1752, 2.7558×10^{-2} , 0.7972)	-2.3699×10^{-4}
	(0.2000, 0.0500, 0.7500)	0.0000
	(0.2530, 0.1748, 0.5722)	-5.9447×10^{-3}
(0.25, 0.05, 0.70)	(0.2500, 0.0500, 0.7000)	0.0000
	(0.2657, 5.5224×10^{-2} , 0.6790)	-3.1317×10^{-4}
(0.30, 0.05, 0.65)	(0.3000, 0.0500, 0.6500)	0.0000
	(0.3048, 2.4285×10^{-2} , 0.6709)	-4.9534×10^{-3}
(0.05, 0.10, 0.85)	(2.1910×10^{-2} , 3.3676×10^{-3} , 0.9747)	-4.7349×10^{-2}
	(0.0500, 0.1000, 0.8500)	0.0000
(0.15, 0.10, 0.75)	(8.3523×10^{-2} , 7.0909×10^{-3} , 0.9094)	-2.1739×10^{-2}

Feed (z_1, z_2, z_3)	Stationary points (x_1, x_2, x_3)	D
	(0.1500, 0.1000, 0.7500)	0.0000
	(0.1544, 0.2908, 0.5548)	-1.7323×10^{-2}
(0.20, 0.10, 0.70)	(0.1257, 1.1877×10^{-2} , 0.8624)	-1.0321×10^{-2}
	(0.2000, 0.1000, 0.7000)	0.0000
	(0.2177, 0.1838, 0.5985)	-1.8376×10^{-3}
(0.05, 0.15, 0.80)	(1.8972×10^{-2} , 2.1628×10^{-3} , 0.9789)	-9.0052×10^{-2}
	(3.6280×10^{-2} , 0.3989, 0.5649)	-3.6271×10^{-2}
	(0.0500, 0.1500, 0.8000)	0.0000
(0.10, 0.15, 0.75)	(4.3189×10^{-2} , 3.2072×10^{-3} , 0.9536)	-6.5345×10^{-2}
	(8.9494×10^{-2} , 0.3230, 0.5875)	-1.1929×10^{-2}
	(0.1000, 0.1500, 0.7500)	0.0000
(0.15, 0.15, 0.70)	(7.3257×10^{-2} , 5.1272×10^{-3} , 0.9216)	-4.1534×10^{-2}
	(0.1500, 0.1500, 0.7000)	0.0000
	(0.1506, 0.2353, 0.6141)	-1.5662×10^{-3}
(0.05, 0.20, 0.75)	(1.8393×10^{-2} , 1.7716×10^{-3} , 0.9798)	-0.1192
	(4.2907×10^{-2} , 0.3218, 0.6353)	-4.3405×10^{-3}
	(0.0500, 0.2000, 0.7500)	0.0000
(0.10, 0.20, 0.70)	(4.1828×10^{-2} , 2.8004×10^{-3} , 0.9554)	-8.2501×10^{-2}
	(9.6537×10^{-2} , 0.2567, 0.6467)	-4.2773×10^{-4}
	(0.1000, 0.2000, 0.7000)	0.0000

Feed (z_1, z_2, z_3)	Stationary points (x_1, x_2, x_3)	D
(0.20, 0.20, 0.60)	(0.1095, 9.3104×10^{-3} , 0.8812)	-1.4729×10^{-2}
	(0.1878, 0.1126, 0.6996)	1.9549×10^{-3}
	(0.2000, 0.2000, 0.6000)	0.0000
(0.10, 0.30, 0.60)	(4.7113×10^{-2} , 3.3396×10^{-3} , 0.9495)	-5.7713×10^{-2}
	(0.1080, 0.1568, 0.7352)	6.8336×10^{-3}
(0.10, 0.50, 0.40)	(0.1000, 0.3000, 0.6000)	0.0000
	(0.1000, 0.5000, 0.4000)	0.0000
	(0.1006, 1.6405×10^{-2} , 0.8830)	0.2374
(0.10, 0.55, 0.35)	(0.1000, 0.5500, 0.3500)	0.0000
	(0.1085, 3.7935×10^{-3} , 0.8877)	0.3639
(0.05, 0.60, 0.35)	(0.0500, 0.6000, 0.3500)	0.0000
	(7.0593×10^{-2} , 1.6098×10^{-2} , 0.9133)	0.3981

4. Conclusions

Using the interval Newton method with the excess Gibbs energy model of NRTL, the phase stability has been predicted for the 10 binary and 2 ternary systems. Although some of the systems are highly non ideal and the local composition model of NRTL may have multiple roots, the interval method has predicted successfully all the stationary points in the tangent plane distance function.

List of symbols

D

tangent plane distance

$f(x)$

real nonlinear equation

g^E

reduced molar excess Gibbs energy

g_m

molar Gibbs energy of mixing surface

G^E

molar excess Gibbs energy

G_{ik}

NRTL binary interaction parameters

ΔG_{mix}

Gibbs energy of mixing

n

number of components

R

gas constant

T

absolute temperature

x

mole fraction

z

feed mole fraction

Greek letters

α_{ik}, α_{ki}

NRTL binary interaction parameters

τ_{ik}

NRTL binary interaction parameters

Subscripts

i

number of components

k

number of components

m

mixture

Superscripts

E

excess

k

iteration number

References

- [1] Y. Demirel and C. McDermott, *Chim. Acta Turc.* **12** (1984), pp. 173–188.
- [2] Y. Demirel and C. McDermott, *Chim. Acta Turc.* **15** (1987), pp. 329–350.
- [3] C. McDermott and Y. Demirel, *Rev. Inorg. Chem.* **10** (1989), pp. 226–233.
- [4] M.L. Michelsen, *Fluid Phase Equilibr.* **9** (1982), pp. 1–19. [Abstract](#)
- [5] M.L. Michelsen, *Fluid Phase Equilibr.* **9** (1982), pp. 21–40. [Abstract](#)
- [6] C.M. McDonald and C.A. Floudas, *AIChE J.* **41** (1995), pp. 1798–1817.
- [7] A.C. Sun and W.D. Seider, *Fluid Phase Equilibr.* **103** (1995), pp. 213–249. [Abstract](#)
- [8] G.A. Iglesias-Silva, A. Bonilla-Petriciolet, P.T. Eubank, J.C. Holste and K.R. Hall, *Fluid Phase Equilibr.* **210** (2003), pp. 229–245. [Abstract](#)
- [9] J.Z. Hua, J.F. Brennecke and M.A. Stadtherr, *Fluid Phase Equilibr.* **116** (1996), pp. 52–59. [Abstract](#)
- [10] J.Z. Hua, J.F. Brennecke and M.A. Stadtherr, *Ind. Eng. Chem. Res.* **37** (1998), pp. 1519–1527. [Abstract-Compendex](#) | [Full Text via CrossRef](#)
- [11] J.Z. Hua, R.W. Maier, S.R. Tessier, J.F. Brennecke and M.A. Stadtherr, *Fluid Phase Equilibr.* **158–160** (1999), pp. 607–615. [Abstract](#)
- [12] C.A. Schnepper and M.A. Stadtherr, *Comput. Chem. Eng.* **20** (1996), pp. 187–199. [Abstract](#)
- [13] S.R. Tessier, J.F. Brennecke and M.A. Stadtherr, *Chem. Eng. Sci.* **55** (2000), pp. 1785–1796. [Abstract](#)
- [14] C.Y. Gau and M.A. Stadtherr, *Comput. Chem. Eng.* **26** (2002), pp. 827–840. [Abstract](#)
- [15] M.A. Stadtherr and C.A. Schnepper, *AIChE Symp. Ser.* **91** (1995), pp. 356–359.
- [16] Y. Demirel and H. Gecegörmez, *Fluid Phase Equilibr.* **65** (1991), pp. 111–133. [Abstract](#)
- [17] L. Lintomen, R.T.P. Pinto, E. Batista, A.J.A. Meirelles and M.R.V. Maciel, *J. Chem. Eng. Data* **46** (2001), pp. 546–550. [Abstract-Compendex](#) | [Full Text](#)