

University of Nebraska - Lincoln

## DigitalCommons@University of Nebraska - Lincoln

---

Gautam Sarath Publications

Biochemistry, Department of

---

1-1-2004

### Hospital Universitario, "La Paz", Madrid, SPAIN

Michael G. Zeece

*University of Nebraska - Lincoln*, mzeece1@unl.edu

Gautam Sarath

*University of Nebraska - Lincoln*, Gautam.sarath@ars.usda.gov

John P. Markwell

*University of Nebraska - Lincoln*, jmarkwell2@unl.edu

N. Palmer

*University of Nebraska-Lincoln*, nathan.palmer@ars.usda.gov

J. Steckelberg

*University of Nebraska - Lincoln*

*See next page for additional authors*

Follow this and additional works at: <https://digitalcommons.unl.edu/biochemistrysarath>



Part of the [Biochemistry, Biophysics, and Structural Biology Commons](#)

---

Zeece, Michael G.; Sarath, Gautam; Markwell, John P.; Palmer, N.; Steckelberg, J.; and Schultz, J., "Hospital Universitario, "La Paz", Madrid, SPAIN" (2004). *Gautam Sarath Publications*. 3.

<https://digitalcommons.unl.edu/biochemistrysarath/3>

This Article is brought to you for free and open access by the Biochemistry, Department of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Gautam Sarath Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

---

**Authors**

Michael G. Zeece, Gautam Sarath, John P. Markwell, N. Palmer, J. Steckelberg, and J. Schultz

## 521 Microarray Assessment of IgE-Binding Proteins in Lentils (*Lens culinaris L.*)

M. G. Zeece<sup>1</sup>, G. Sarath<sup>2</sup>, J. Markwell<sup>3</sup>, N. Palmer<sup>1</sup>, J. Steckelberg<sup>1</sup>, J. Schultz<sup>4</sup>, C. Pascual<sup>5</sup>; <sup>1</sup>Food Science, University of Nebraska, Lincoln, NE, <sup>2</sup>USDA-ARS, Lincoln, NE, <sup>3</sup>Biochemistry, University of Nebraska, Lincoln, NE, <sup>4</sup>University of Nebraska, Lincoln, NE, <sup>5</sup>Hospital Universitario, "La Paz", Madrid, SPAIN.

**RATIONALE:** Proteomic technology has been shown to have distinct advantages for allergen characterization and identification. Therefore, we have used 2-D electrophoresis and protein microarrays to characterize the profile of allergens and assess their level in lentils.

**METHODS:** Lentil varieties were extracted and centrifuged to obtain 2S, 7S and 11S fractions. The profile of IgE-binding proteins in these fractions was assessed using lentil-allergic sera, by immunoblotting of 2-D electrophoretic separations. The level of IgE-binding was also assessed using protein microarrays. Arrays of covalently attached proteins were first incubated with lentil-allergic sera, followed by incubation with fluorescently-labeled anti-IgE detection antibody. Laser-induced fluorescence was used to detect IgE-binding.

**RESULTS:** Immunoblotting of electrophoretic separations showed several regions of IgE-binding that were consistently observed for all varieties. Major IgE-binding was observed for isoforms of 31 and 48 kDa spots. These proteins are of particular interest because of their low abundance, but high IgE-binding. Comparatively less IgE-binding was found for 12 and 14 kDa spots. Microarray analysis showed that IgE-binding of the 11S fraction was significantly greater than all other fractions. Microarrays enabled quantitative assessment of IgE-binding.

**CONCLUSIONS:** The application of proteomic technologies resulted in characterization of IgE-binding protein profile and determination of their level in lentils. Microarrays represent a promising tool for allergen investigation, detection and diagnosis.

*Funding: University of Nebraska*

SUNDAY