

8-1-2018

# Safe use of Hydrogen Peroxide in the Organic Lab

Patrick Dussault

*University of Nebraska-Lincoln*, [pdussault1@unl.edu](mailto:pdussault1@unl.edu)

Follow this and additional works at: <http://digitalcommons.unl.edu/chemistryperoxides>

 Part of the [Higher Education Commons](#), [Organic Chemistry Commons](#), and the [Science and Mathematics Education Commons](#)

---

Dussault, Patrick, "Safe use of Hydrogen Peroxide in the Organic Lab" (2018). *Organic Peroxides: Safety Issues*. 3.  
<http://digitalcommons.unl.edu/chemistryperoxides/3>

This Article is brought to you for free and open access by the Chemistry, Department of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Organic Peroxides: Safety Issues by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

## Safe use of Hydrogen Peroxide in the Organic Lab

Pat Dussault

Department of Chemistry, University of Nebraska-Lincoln

e-mail: pdussault1@unl.edu phone: 402-472-6951

(August 2018)

### Outline

---

Leading references

Toxicity

Physical properties

Chemical properties

Stability and incompatibilities

Confinement/pressure

References to accidents involving H<sub>2</sub>O<sub>2</sub>

### Leading references

*Applications of Hydrogen Peroxide and Derivatives* offers a very useful introduction, including discussions of: methods for preparation; properties; use as an oxidant in the presence of catalysts; synthetic application; and, environmental applications.<sup>1</sup> The web sites of several suppliers and a paper industry trade group offer information on: physical properties, safety, guidelines for handling/storage/usage, and regulatory requirements.<sup>2</sup> Information on aq. H<sub>2</sub>O<sub>2</sub> solutions is available in the American Chemical Society *Reagent Chemicals* series.<sup>3</sup> Safety data sheets for commercially available solutions are available from all major vendors;<sup>2</sup> a link to a safety data sheet portal is provided.<sup>4</sup> A review describing industrial synthesis of H<sub>2</sub>O<sub>2</sub> is available.<sup>5</sup>

### Toxicity

Exposure to odorless (no odor threshold is known) hydrogen peroxide vapors has resulted in injuries to several dozen employees.<sup>6</sup> H<sub>2</sub>O<sub>2</sub> (vapor, mist, or aerosol) is considered to hold immediate danger to life and health (IDLH) at 75 ppm;<sup>7</sup> the threshold limit value (TLV) for time-weighted exposure is 1.0 ppm.<sup>8</sup> Dermal exposure to 30% H<sub>2</sub>O<sub>2</sub> can produce skin damage in a few minutes and serious eye damage in only a few seconds.<sup>9,10</sup> Although relatively modest acute toxicities (LD<sub>50</sub>) are reported for oral and dermal exposure in rodent models (2-4 g/kg and

> 10 g/kg, respectively),<sup>1</sup> accidental human ingestion of one pint of 30% aq. H<sub>2</sub>O<sub>2</sub> resulted in a near fatality.<sup>11</sup>

### Physical properties

Hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>, [7722-84-1]), is an odorless and colorless molecule with a boiling point of approximately 150 °C.<sup>1,2</sup> Gaseous mixtures rich in H<sub>2</sub>O<sub>2</sub> can explode violently and without warning; under no conditions should ambient pressure distillation be attempted. The melting point of highly concentrated H<sub>2</sub>O<sub>2</sub> is just below 0 °C but the freezing point of aqueous solutions is typically much lower.<sup>1</sup> Cooling of dilute aqueous solutions typically results in separation of water ice (and some concentration of the hydrogen peroxide) while concentrated solutions can undergo supercooling.<sup>12</sup> A useful summary of physical properties of H<sub>2</sub>O<sub>2</sub> solutions is available.<sup>2a</sup>

### Chemical properties

Acidity: The pK<sub>a</sub> of H<sub>2</sub>O<sub>2</sub> is ~ 12 and concentrated aqueous solutions can have pH values as low as 4.3.<sup>1</sup> As a result, self-catalyzed reactions of 30% H<sub>2</sub>O<sub>2</sub> with carbonyls to form 1,1-dihydroperoxides can occur.<sup>13</sup>

As oxidant: H<sub>2</sub>O<sub>2</sub> is a powerful oxidant, with an E<sub>0</sub> (1.8 V) falling between KMnO<sub>4</sub> and O<sub>3</sub>.<sup>1,2</sup> Although catalysts are often required to achieve high rates of oxidation, exposure of concentrated H<sub>2</sub>O<sub>2</sub> solutions to flammable or combustible materials (including wood pallets, paper) can result in fires and must be avoided.<sup>2,11</sup>

### Stability and incompatibilities

Safe Handling and Storage: Many of the hazards, warnings, and mitigation principles described in the following section are nicely laid out in technical bulletins, many of which are maintained on vendor web sites.<sup>2a-d, 14</sup> The short version: handling, storage, and/or transport of hydrogen peroxide must take into account decomposition (and attendant generation of oxygen) but also the high reactivity towards many organic materials, (including simple combustibles such as paper and wood), many inorganics, (particularly reducing agents), many metals (why container choice can be important), and many redox-active metal ions.

Incompatibilities: Concentrated solutions of H<sub>2</sub>O<sub>2</sub> can be highly reactive towards organics and combustible materials and can decompose with considerable force,<sup>1,2, 15</sup> mixtures of hydrogen

peroxide and organics may explode upon heating even in the absence of metals.<sup>16</sup> Hydrogen peroxide should never be used in ketone solvents or stored in the presence of aliphatic ketones or almost any aldehydes. In the presence of even moderate amounts of strong acids, such mixtures can be converted to highly explosive 1,2,4,5-tetraoxanes (ketone peroxide dimers), or 1,2,4,5,7,8-hexaoxonanes (trimers), or polymeric peroxides,<sup>17</sup> and formation of these and other ketone/ H<sub>2</sub>O<sub>2</sub> adducts has been observed even in the absence of acid.<sup>13,18</sup> The curious stabilization of H<sub>2</sub>O<sub>2</sub> solutions by small amounts of aryl alkyl ketones is discussed below.

Reactions of hydrogen peroxide, including disproportionation, are often catalyzed by redox-active transition metals (Fe<sup>+2/+3</sup> and Cu<sup>+/+2</sup> are two of the best known systems) and exposure of concentrated solutions to metal ions or precursors of metal ions should be avoided at all costs. Thus, transfer of aq. H<sub>2</sub>O<sub>2</sub> should employ Teflon needles; if it is absolutely necessary to use a metal needle for extremely small volumes, then approximate amount of H<sub>2</sub>O<sub>2</sub> should be transferred first to a small vial and any leftover reagent discarded into running water.

Cleaning solutions based upon mixtures of H<sub>2</sub>O<sub>2</sub> and strong acids. for example, sulfuric ("Piranha" solution) or nitric) are remarkably corrosive, and call for great care in terms of procedure, personal protective equipment, and choice of vessel/container (the solutions are very destructive of many plastics).<sup>19</sup> Under strongly acidic conditions, H<sub>2</sub>O<sub>2</sub> can undergo relatively rapid decomposition with liberation of oxygen;<sup>20</sup> bottles must be vented to avoid over pressurization. Acetone and any other low-molecular weight carbonyl compound must not be introduced into acidic solutions of peroxides: see: <http://cenblog.org/the-safety-zone/2015/01/piranha-solution-explosions/>

#### Stabilization of aqueous solutions:

Highly pure aqueous solutions may lose only a few percent of activity each year.<sup>2</sup> However, in the words of Shanley and Greenspan "Hydrogen peroxide is remarkable for the number and variety of decomposition catalysts and for the minute quantities required to give large effect."<sup>14</sup> These and other authors note a remarkable acceleration in the rate of decomposition under basic conditions. Commercially marketed aqueous solutions of hydrogen peroxide typically contain one or more inorganic or organic stabilizers. A recent report describing aryl alkyl ketone stabilizers provides an entry to this discussion.<sup>21</sup>

Solutions of H<sub>2</sub>O<sub>2</sub> in organic solvents: Hydrogen peroxide has good solubility in many organic solvents and experimenters unable to access the highly concentrated (e.g., 90%, > 30 M)<sup>2</sup>

solutions described in many older synthetic procedures<sup>14</sup> have sometimes turned to organic solutions prepared via extraction of 30 or 50% aqueous solutions with ether or dichloromethane.<sup>22,23,24</sup> These "anhydrous" solutions, which are typically 1-2 M in H<sub>2</sub>O<sub>2</sub>, should be considered inherently unstable;<sup>2d</sup> our group, for example, limits preparation to a maximum of 10 mL. A recent article describes safety issues related to generation and ignition of O<sub>2</sub> head gas within organic solutions of H<sub>2</sub>O<sub>2</sub>.<sup>25</sup> Attempts to further concentrate ethereal solutions of H<sub>2</sub>O<sub>2</sub> through evaporation or distillation are ineffective and represent needless hazard.<sup>26</sup> Similarly, ethereal solutions of H<sub>2</sub>O<sub>2</sub> should never be stored, but immediately consumed or quenched. The urea•H<sub>2</sub>O<sub>2</sub> complex, as well as salts of perborates and percarbonates, are sometimes applied as substitutes for H<sub>2</sub>O<sub>2</sub>,<sup>27,28</sup> however, these have limited solubility in aprotic solvents.<sup>29</sup> A recently described alternative approach for generation of dry H<sub>2</sub>O<sub>2</sub> involves dissolution of amino acid perhydrates into organic solvents.<sup>30</sup> Note that several recent publications describe the incompatibility of even dilute solutions of H<sub>2</sub>O<sub>2</sub> with acetone and other ketones or aldehydes.<sup>14</sup> A statement from Solvay's guide on handling and storage is useful to remember: "Investigation of new applications involving hydrogen peroxide and other chemicals should be carried out on a small scale, with adequate precautions taken for dealing with uncontrolled and potentially explosive reactions."<sup>2d</sup>

#### **Confinement/pressure/temperature.**

H<sub>2</sub>O<sub>2</sub> readily decomposes to liberate O<sub>2</sub> and the resulting enrichment of oxygen in the headspace of containers and reactors can dramatically increase fire hazards.<sup>1</sup> Over the range of 20-100 °C, the rate of decomposition increases more than two-fold for each rise of 10 °C and confined reactions should be avoided unless adequate precautions have been taken to avoid unintentional over pressurization.<sup>2</sup> Solutions of hydrogen peroxide are best stored with venting caps. Guidelines for safe storage and handling are available.<sup>2a-d.</sup>

**Literature on accidents.** The listed references discuss accidents related to hydrogen peroxide.<sup>6,9,11,15</sup>

Acknowledgments: We are grateful to NSF for support of our research on ozonolysis and peroxide chemistry (CHE- 0749916, 1057982, 1469414). We thank Prof. Paul Royster, UNL Libraries, for assistance in posting this material onto the Digital Commons and Alissa Horn and Moriah Locklear for useful suggestions.

- 
- <sup>1</sup> Jones, C. W. *Applications of Hydrogen Peroxide and Derivatives*, RSC Clean Technology Monographs, Royal Society of Chemistry, Cambridge, 1999.
- <sup>2</sup> a. <http://www.h2o2.com/technical-library/default.aspx?pid=9&name=Technical-Library> (Accessed August 2018); b. <http://h2o2.evonik.com/product/h2o2/en/about-hydrogen-peroxide/basic-information/safety-and-handling/pages/default.aspx> (Accessed August 2018); c. <http://www.peroxychem.com/chemistries/hydrogen-peroxide> (accessed August 2018); d. [https://www.solvay.co.th/en/binaries/H2O2\\_Handling%20and%20Storage-191789.pdf](https://www.solvay.co.th/en/binaries/H2O2_Handling%20and%20Storage-191789.pdf); <http://www.solvay.us/en/markets-and-products/Literature/Peroxides-TDS.html> (both accessed August 2018).
- <sup>3</sup> "Hydrogen Peroxide Solutions" *ACS Reagent Chemicals* <https://pubs.acs.org/doi/abs/10.1021/acsreagents.4167> (accessed August 2018)
- <sup>4</sup> <http://icca.cefic.org/en/Home/Global-Product-Strategy/global-product-strategy/chemical-information-search/> (accessed August 2018)
- <sup>5</sup> "Hydrogen peroxide synthesis: an outlook beyond the anthraquinone process" Campos-Martin, J. M.; Blanco-Brieva, G.; Fiero, J. L. G. *Angew. Chem. Int. Ed.* **2006**, *45*, 6962-84.
- <sup>6</sup> <https://www.osha.gov/pls/imis/accidentsearch.html> (August 2018; for example, can search for "hydrogen peroxide")
- <sup>7</sup> a. "Medical Management Guidelines for Hydrogen Peroxide" ATSDR Toxic Substances Portal. <https://www.atsdr.cdc.gov/mmg/mmg.asp?id=304&tid=55>; b. <http://www.cdc.gov/niosh/idlh/intridl4.html> (both accessed August 2018)
- <sup>8</sup> <http://www.atsdr.cdc.gov/MMG/MMG.asp?id=304&tid=55> (accessed August 2018)
- <sup>9</sup> "Management of 35% hydrogen peroxide exposure to naked eyes: a case report and review" Sreeram, S. R.; Singh, P.; Thomas, P.; Unni, K. N.; Tharani, S. *Int. J. Pharma and Bio Sciences*, **2015**, *6*, 417-420.
- <sup>10</sup> [https://www.gov.uk/government/uploads/system/uploads/attachment\\_data/file/337708/Hydrogen\\_Peroxide\\_Toxicological\\_Overview\\_phe\\_v1.pdf](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/337708/Hydrogen_Peroxide_Toxicological_Overview_phe_v1.pdf) (accessed August 2018)
- <sup>11</sup> "Hydrogen Peroxide Accidents And Incidents: What We Can Learn From History" Greene, B.; Baker, D. L.; Frazier, W. White Sands Test Facility internal document (2004), <https://ntrs.nasa.gov/archive/nasa/casi.ntrs.nasa.gov/20050217417.pdf> (accessed August 2018)
- <sup>12</sup> "The supercooling of aqueous hydrogen peroxide" Cooper, K. A.; Watkinson, J. G. *Trans. Faraday Soc.*, **1957**, *53*, 635-641.
- <sup>13</sup> "A facile catalyst-free synthesis of gem-dihydroperoxides with aqueous hydrogen peroxide" Tada, N.; Cui, L.; Okubo, H.; Miura, T.; Itoh, A. *Chem. Commun.* **2010**, *46*, 1772-4; see also Espinosa-Fuentes, E.A., Pacheco-Londoño, L.C., Barreto-Cabán, M.A. and Hernández-Rivera, S.P., 2012. Novel uncatalyzed synthesis and characterization of diacetone diperoxide. *Propellants, Explosives, Pyrotechnics*, **2012**, *37*, 413-421.
- <sup>14</sup> "Hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) safe storage and handling" TIP 0606-24; TAPPI; [https://www.fpl.fs.fed.us/documnts/pdf2010/fpl\\_2010\\_anderson001.pdf](https://www.fpl.fs.fed.us/documnts/pdf2010/fpl_2010_anderson001.pdf) (accessed August 2018)

---

<sup>15</sup> "Highly Concentrated Hydrogen Peroxide: Physical and Chemical Properties" Shanley, E. S.; Greenspan, F. D. *Ind. Eng. Chem.* **1947**, *39*, 1536.

<sup>16</sup> <http://cenblog.org/the-safety-zone/2017/02/from-the-archives-chemists-move-wall-with-hydrogen-peroxide-explosion/> (accessed August 2018)

<sup>17</sup> P. R. Story, B. Lee, D. D. Denson, P. Busch, *J. Org. Chem.* **1970**, *35*, 3059, and references within.

<sup>18</sup> Oxley, J.C., Brady, J., Wilson, S.A. and Smith, J.L., 2012. The risk of mixing dilute hydrogen peroxide and acetone solutions. *J. Chem. Health Safety* **2012**, *19*, 27-33;

<sup>19</sup> For a useful safety summary, see: <https://www.drillinois.edu/SafetyLibrary/PiranhaSolutions> (accessed August 2018)

<sup>20</sup> In a solution of 4M nitric acid at elevated temperatures, the half-life for hydrogen peroxide can be less than two hours: "Rate of Hydrogen Peroxide Decomposition in Nitric Acid Solutions" Miner, F. J.; Hagan, R. G. *Ind. Eng. Chem. Proc. Des. Dev.* **1972** *11*, 547-549; DOI: 10.1021/i260044a017

<sup>21</sup> Terent'ev, A.O.; Pastukhova, Z.Y.; Yaremenko, I.A.; Bruk, L.G.; Nikishin, G.I., Promising hydrogen peroxide stabilizers for large-scale application: unprecedented effect of aryl alkyl ketones. *Mendeleev Communications*, **2016**, *26*, 329.

<sup>22</sup> Extraction into ether has been long known. See, for example: *J. Chem. Soc. Industry*, **1900**, *19*, 777.

<sup>23</sup> In our experience, extraction of 1-2 mL of 30% aq. H<sub>2</sub>O<sub>2</sub> with 10 mL of diethyl ether furnishes a 1-2 M ethereal solution.

<sup>24</sup> "Selective Oxidation by Peroxide-based Reagents" Nagata, R.; Saito, I. *Synlett* **1990**, 291-300.

<sup>25</sup> "Safe Scale-Up of Oxidation by Hydrogen Peroxide in Flammable Solvents" Astbury, G. R. *Organic Process Research & Development* **2002** *6*, 893-895; DOI: 10.1021/op025574y

<sup>26</sup> "Concerning the preparation of anhydrous hydrogen peroxide" Dove J. E.; Riccick, J. *Can. J. Chem.* **1968**, *46*, 330-332, doi 10.1139/v68-050

<sup>27</sup> "Oxidation Reactions Using Urea-Hydrogen Peroxide; A Safe Alternative to Anhydrous Hydrogen Peroxide" Cooper, M. S.; Heaney, H.; Newbold, A. J.; Sanderson, W. R. *Synlett*, **1990**, 533-5, DOI: 10.1055/s-1990-21156.

<sup>28</sup> "The 1,4-diaza[2.2.2] Bicyclooctane—hydrogen peroxide complex as a source of anhydrous hydrogen peroxide: the preparation of bis(trialkylsilyl) peroxides" Cookson, P. G.; Davies, A. G.; Fazal, N. *J. Organomet. Chem.*, **1975**, *99*, C31.

<sup>29</sup> "Sodium perborate and sodium percarbonate: Cheap, safe and versatile oxidising agents for organic synthesis" McKillop, A.; Sanderson, W. R. *Tetrahedron*, **1995**, *51*, 6145–6166.

<sup>30</sup> "Preparation of pure hydrogen peroxide and anhydrous peroxide solutions from crystalline serine perhydrate" Wolanov, Y., et al, *Tetrahedron*, **2010**, *66*, 5130- 5133, <http://www.sciencedirect.com/science/article/pii/S0040402010006915>