

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

Nutrition & Health Sciences Dissertations & Theses

Nutrition and Health Sciences, Department of

---

Spring 5-2010

## UNSATURATED FATTY ACIDS REPRESS THE EXPRESSION OF ADIPOSE FATTY ACID-BINDING PROTEIN, aP2, IN RAW 264.7 MACROPHAGE

Sara L. Coleman

University of Nebraska at Lincoln, sara34@gmail.com

Follow this and additional works at: <https://digitalcommons.unl.edu/nutritiondiss>



Part of the [Dietetics and Clinical Nutrition Commons](#)

---

Coleman, Sara L., "UNSATURATED FATTY ACIDS REPRESS THE EXPRESSION OF ADIPOSE FATTY ACID-BINDING PROTEIN, aP2, IN RAW 264.7 MACROPHAGE" (2010). *Nutrition & Health Sciences Dissertations & Theses*. 5.

<https://digitalcommons.unl.edu/nutritiondiss/5>

This Article is brought to you for free and open access by the Nutrition and Health Sciences, Department of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Nutrition & Health Sciences Dissertations & Theses by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

UNSATURATED FATTY ACIDS REPRESS THE  
EXPRESSION OF ADIPOSE FATTY ACID-BINDING  
PROTEIN, aP2, IN RAW 264.7 MACROPHAGE

by

Sara L. Coleman

A THESIS

Presented to the Faculty of

The Graduate College at the University of Nebraska

In Partial Fulfillment of Requirements

For the Degree of Master of Science

Major: Nutrition & Health Sciences

Under the Supervision of Professor Ji-Young Lee

Lincoln, Nebraska

May 2010

# UNSATURATED FATTY ACIDS REPRESS THE EXPRESSION OF ADIPOSE FATTY ACID-BINDING PROTEIN, aP2, IN RAW 264.7 MACROPHAGE

Sara L. Coleman, M.S.

University of Nebraska, 2010

Adviser: Ji-Young Lee

Adipocyte fatty acid binding protein, aP2, present in macrophages has been implicated in the integration of lipid metabolism and inflammatory response, contributing to development of insulin resistance and atherosclerosis. We investigated the modulation of aP2 expression by inflammatory insults and fatty acids in RAW 264.7 macrophages. When the cells were incubated with lipopolysaccharides (LPS; 100 ng/ml) or 10 ng/ml of tumor necrosis factor  $\alpha$  for 18 h, aP2 mRNA and protein levels were drastically increased. Unsaturated fatty acids (100  $\mu$ M of 18:1, 18:2, 18:3, 20:5 in complex with BSA), but not saturated fatty acids (16:0), significantly repressed the basal aP2 expression and abolished induction of aP2 expression by LPS. Trichostatin A (TSA), a histone deacetylase (HDAC) inhibitor, increased aP2 mRNA levels but abolished the repressive effect of 18:2 on aP2 expression in unstimulated and LPS-stimulated macrophages. Cells transfected with siRNA targeting aP2 showed a relationship with cyclooxygenase-2 (COX-2), but not other pro-inflammatory mediators. In summary, our data suggest that unsaturated fatty acids may inhibit the basal as well as LPS-induced aP2 expression by mechanisms involving HDAC in RAW 264.7 macrophages and that there may be a link between COX-2 and aP2-repression by unsaturated fatty acids.

## Table of Contents

	Page
List of Tables	i
List of Figures	ii
Introduction	1
Literature Review	4
Obesity and Inflammation	4
Inflammatory Signaling Pathways	5
PPAR $\gamma$	6
NF- $\kappa$ B Pathway	7
MAPK and AP-1 Pathway	8
Inflammatory Cytokines	9
COX-2 and Anti-Inflammatory Drugs	10
Fatty Acids	11
Fatty Acid Metabolism	11
Fatty Acid Structure	12
Adipose Fatty Acid-Binding Protein, aP2	13
Histone Modification	15
Histone Deacetylase	16
Materials and Methods	18
Results	23
Discussion	32
References	37

## List of Tables

Table	Page
1. Primers for quantitative real-time PCR	22

## List of Figures

Figure	Page
1. Elevated aP2 levels by LPS	24
2. a) Elevated aP2 mRNA levels abrogated by unsaturated fatty acids	25
b) Elevated aP2 protein levels abrogated by unsaturated fatty acids	25
3. Abrogation of 18:2-mediated aP2 repression by TSA	27
4. a) Western-Blot Analysis of Cytoplasmic Fraction for aP2 knock-down	28
b) HDAC Activity	28
5. siRNA knock-down of aP2 mRNA expression	30
6. Effect of aP2 and FA on pro-inflammatory gene expression	31

## Introduction

The link between lipid metabolism and inflammatory pathways has been suggested as a significant component in the development of insulin resistance, atherosclerosis, and type 2 diabetes (1-3). Dietary fat is an important macronutrient for the growth and development of all organisms (4). Excessive levels of dietary fat or an imbalance of saturated versus unsaturated fat have been implicated in the onset and progression of several chronic diseases, including coronary artery diseases, atherosclerosis, diabetes, obesity, and cancer (4). Specifically, unsaturated fatty acids are known to have positive health benefits such as reductions in atherosclerosis, inflammation, and plasma free fatty acid levels (5). However, the mechanisms for the anti-inflammatory properties of unsaturated fatty acids are poorly understood.

Fatty acid binding proteins (FABP) are implicated as players in the inflammatory response as well as lipid signaling cascades (2, 3). FABP are intracellular lipid chaperones that have a role in import, storage, and export of fatty acids because they reversibly bind hydrophobic ligands, such as saturated/ unsaturated long-chain fatty acids with high affinity (2).

The most characterized FABP is adipocyte fatty acid binding protein (aP2/FABP4), which is found in adipocytes at high levels and in macrophage cells to a lesser extent (2). Studies have shown that aP2 deficiency gives partial protection against atherosclerotic lesion formation in apolipoprotein E knockout mice, as well as, systemic insulin resistance, dyslipidemia, and lipotoxicity in *aP2<sup>-/-</sup> ob/ob* mice (2, 6).

The chronic low-grade inflammation associated with obesity is perpetuated by macrophage infiltration into lipid-engorged adipose tissue; this results in a release of various inflammatory cytokines (7, 8). Accompanying an obesity-induced inflammatory state are elevated circulating levels of aP2 (9).

The majority of research on the regulation of aP2 expression has been in adipocytes. Various cis-acting elements of the aP2 gene have been suggested, including an activator protein -1 (AP-1) sequence, and CAAT/enhancer-bind protein (C/EBP) binding element (10, 11). Recently it was demonstrated that macrophage aP2 can be regulated by phorbol 12-myristate 13- acetate, lipopolysaccharide, oxidized low-density lipoprotein and PPAR $\gamma$  ligands (12).

One way to affect gene expression is through the involvement of coactivator/ corepressor complexes (13). Histone deacetylases (HDAC) are essential components of co-repressor complexes as they provide the enzymatic activity for active repression(14). HDAC are a class of enzymes that remove acetyl groups from an  $\epsilon$ -N-acetyl lysine amino acid on a histone. Its action is opposite to that of histone acetyltransferase (HAT) (15). Removal of acetyl groups gives chromatin a tightly-bound structure that prohibits transcription. Inclusion of acetyl groups decreases the affinity of histones for DNA and allows transcription to occur (15). There are eighteen known HDAC, which are divided into four classes. Class 1 HDAC are generally detected in the nucleus and ubiquitously expressed. Specifically, HDAC3 is unique among class 1 HDAC in that it can shuttle between the nucleus and cytoplasm and form complexes(15). Transcription factors that have been shown to interact with the corepressor complex for repressive effects on gene transcription include: AP-1 (16-18), nuclear factor  $\kappa$ B (NF- $\kappa$ B) (16, 17, 19), liver X



receptor  $\alpha$  (20), and thyroid hormone receptor (13, 21, 22). Upon activation, the co-repressor complex is exchanged with co-activators that lead to transcription (14).

Here we demonstrate histone modification involving HDAC as a mechanism for how fatty acids can affect aP2 expression in murine RAW 264.7 macrophage. Though therapeutic drugs are currently being developed to exploit aP2 for the treatment of diabetes and atherosclerosis (3, 12), the mechanisms for how aP2 expression is controlled under basal and activated conditions in macrophage is still unknown. Elucidation of aP2 regulatory mechanisms would allow for more accurate manipulation of pathways by dietary or pharmaceutical agents in order to positively influence health status.

## Literature Review

### Obesity and Inflammation

Obesity, which is defined as a body mass index (BMI) greater than or equal to 30 kg per m<sup>2</sup>, predisposes an individual to develop insulin resistance, cardiovascular disease, diabetes, hypertension, high cholesterol, stroke, and certain cancers (23-25). The obesity epidemic reached a peak in 2003, and currently more than one-third of adults over age 20 in the United States are considered obese (24). Much of the complications associated with obesity are thought to be a product of the accompanying chronic inflammatory state in the adipose tissue (1, 7, 23, 25).

Adipose tissue is now being recognized as an endocrine organ, due to its ability to release pro- and anti-inflammatory protein factors, known as adipokines (26, 27). A chronic inflammatory response, characterized by elevated adipokine production and the activation of some pro-inflammatory signaling pathways, causes an induction of several biological markers of inflammation (28). These variations in gene expression of white adipose tissue are attributed to the infiltration of monocytes into lipid-engorged adipose tissue, which has been seen in both mouse models of obesity and obese humans (8, 28). It has been proposed that monocyte chemoattractant protein- 1 (MCP-1) (8), leptin (29), or activated endothelial cells (29, 30) could contribute to the macrophage infiltration into adipose tissue. Macrophages in the adipose tissue are responsible for most of the locally-produced cytokines such as tumor necrosis factor  $\alpha$  (TNF $\alpha$ ) and interleukin-6 (IL-6). TNF $\alpha$  is known to interfere with energy metabolism, especially lipid homeostasis by

increasing free fatty acid production (31). *In vitro* studies using adipocyte cell lines support the role of IL-6 as a factor to cause insulin resistance in adipocytes (32).

It is believed that the increases in local cytokine production from macrophage in lipid-engorged macrophages is the cause or possibly the consequence of the low-grade inflammation associated with obesity (28, 33, 34). The connection between adipose tissue and the immune system is clear, but why such a link exists is up for speculation.

Coppack postulates that with the immune system using about 15% of resting metabolic rates, it would need to have a link to the availability of energy, especially the size of the adipose energy stores (35). This idea gives evolutionary purpose to the cross-talk between the immune and metabolic systems.

## **Inflammatory Signaling Pathways**

Toll-like receptors (TLRs) play a critical role in the activation of the innate immune response in mammals by recognizing conserved pathogen-associated molecular patterns on invading bacteria (36). There are currently 13 known members of the TLR family in mammals, with TLR4 being a subclass (36). TLR4 can be activated by lipopolysaccharide (LPS), originating from the outer-membrane of gram negative bacteria, and non-bacterial agents such as saturated fatty acids (36, 37). The activation of TLR4 induces upregulation of intracellular inflammatory pathways such as c-Jun NH<sub>2</sub>-terminal kinase (JNK), nuclear factor- $\kappa$ B (NF- $\kappa$ B), and mitogen activated protein kinase (MAPK) pathways. The MAPK pathway can then lead to activation of the transcription factor activating protein 1 (AP-1) (36, 38).

TLR signaling in response to pathogen contact can contribute to the development of atherosclerosis (38). Loss of TLR4 leads to a protection against diet-induced obesity, I Kappa B Kinase  $\beta$  (IKK $\beta$ ), and JNK expression, and insulin resistance (36). Activation of TLR4 by saturated fatty acids, not unsaturated fatty acids, can induce the expression of cyclooxygenase-2 (COX-2), a marker for inflammation (39). Tsukuma *et al.* have suggested TLR4 may be a candidate for participation in insulin resistance induced by a saturated fatty acid-rich diet (36). It has been observed that saturated fatty acids can directly interact with the immune modulation and inflammation response through the activation of TLRs in macrophage (39).

Activation of the innate immune system can also be a result of TNF $\alpha$ . Binding of TNF $\alpha$  to its cell surface receptor induces the inflammatory response. TNF $\alpha$  is a major adipokine and its expression is highly induced in obesity (32, 35). It is known to interfere with insulin signaling after receptor binding on muscle cells or hepatocytes (32). TNF $\alpha$  causes an increase in free fatty acids, which encourage the development of insulin resistance in persons with metabolic disorders (31). Activation of a macrophage by either LPS or TNF $\alpha$  leads to downstream convergence onto similar inflammatory pathways. One route is through the IKK which signals the NF-KB pathway(40). Secondly, is through JNK and the AP-1 pathway (40).

### ***Peroxisome Proliferator Activator Receptor $\gamma$***

PPAR $\gamma$  is a ligand-activated member of the nuclear receptor super family known to be involved in gene regulation of inflammatory and metabolic pathways (41, 42).

PPAR $\gamma$  was initially discovered as the master regulator in adipose cells (43). It is essential for the regulation of adipogenesis and is required for maintenance of mature adipocyte function, including triglyceride synthesis and storage (43-45). PPAR $\gamma$  is also expressed in activated monocytes/ macrophage and its agonist can inhibit the production of inflammatory cytokines by antagonizing the activities of transcription factors such as AP-1 and NF- $\kappa$ B (46, 47). One such antagonizing mechanism is through the ligand-dependant sumoylation of the PPAR $\gamma$  ligand-binding domain which inhibits the removal of a repressor complex from inflammatory gene promoters (14, 48). Alternatively, anti-inflammatory PPAR $\gamma$  activity can be inhibited by TNF $\alpha$  and this inhibition is thought to be involved in the pathogenesis of insulin resistance, atherosclerosis, and inflammation (6, 49). Naturally occurring ligands for PPAR $\gamma$  include long-chain fatty acids; other known synthetic PPAR $\gamma$  ligands include the thiazolidinedione class of anti-diabetic drugs (43). It is unclear as to whether aP2 enhances or inhibits the transcriptional activity of PPAR $\gamma$ , as experiments showing both have been reported (50, 51). A culmination of recent data suggests PPAR $\gamma$  agonists may regulate inflammation by the activation of PPAR $\gamma$  or alternatively through a PPAR $\gamma$ -independent mechanism (42, 44).

### ***NF- $\kappa$ B Pathway***

Production of pro-inflammatory cytokines is regulated largely by NF- $\kappa$ B (40, 42, 52, 53). It plays a pivotal role in the regulation over 100 genes, including TNF $\alpha$ , MCP-1, and cyclooxygenase 2 (COX-2) (54). In the canonical NF- $\kappa$ B pathway, p65 and p50 form a dimer and its activation is regulated by kappa B inhibitor protein (IkB) (14).

Following stimulation by inflammatory stimuli such as LPS, rapid proteasomeal degradation of I $\kappa$ B occurs after being phosphorylated by I $\kappa$ B kinase (IKK) (40). This allows NF- $\kappa$ B to shuttle from the cytoplasm into the nucleus, where it then promotes the transcription of target inflammatory genes (40).

Dietary agents have been shown to affect the NF- $\kappa$ B pathway. Fraxinellone, derived from root bark of *Dictamnus dasycarpus*, inhibits LPS-induced inflammatory response by negatively regulating NF- $\kappa$ B in RAW 264.7 macrophages (55). In contrast, *in vitro* culture of monocytes under high glucose condition compared to normal glucose led to the activation of NF- $\kappa$ B and significantly increased the expression of inflammatory cytokines including TNF $\alpha$  and MCP-1 (54).

### ***MAPK and AP-1 Pathway***

Upon binding of a peptide:MHC complex to a T-cell receptor, a cascade of proteins initiate the activation of the MAPK pathway (56). In the MAPK pathway, a three-kinase relay system, or MAPK cascade, involves three phosphorylation events that can lead to new gene transcription (56). The three most characterized groups of MAPK are extracellular signal-regulated kinases (ERK), p38 MAPK, and JNK (57). ERK are robustly activated by growth factors and phorbol esters but only weakly activated by cytokines and environmental stress (57). In contrast, p38 MAPK and JNK are strongly activated by cytokines and environmental stress but are poorly activated by growth factors and phorbol esters (57, 58). ERK works with the transcription factor Elk-1 to initiate transcription of Fos, a dimer component of AP-1 (56). JNK controls transcriptional activity of AP-1 via phosphorylation of c-Jun, allowing translocation of c-

Jun to the nucleus (56, 58). In the nucleus, c-Jun can dimerize with Fos to form AP-1 (56). In LPS-induced macrophages, phosphorylation of JNK was shown to initiate an inflammatory responses (59). Furthermore, JNK activity is significantly elevated in tissues of type 2 diabetic patients and in animals of obesity and diabetes (58).

AP-1 is a transcription factor that is a heterodimer of c-Jun, c-Fos, or activating transcription factor ATF proteins (60). It is activated by growth factors, neurotransmitters, cytokines, and bacterial or virus infections and is known to be involved in the transcriptional regulation of the inflammatory response (59).

## **Inflammatory Cytokines**

Cytokines are a group of small, diverse proteins that are released by cells in the body in response to activating stimulus, such as pathogen invasion. They induce responses by binding to specific receptors and exert their effects in an autocrine, paracrine, or endocrine manner (35). The cytokines secreted by macrophage include: IL-1 $\beta$ , IL-6, IL-12, and TNF- $\alpha$ . IL-1 $\beta$  is responsible for activating vascular endothelium and lymphocytes (56). It is the cytokine responsible for the induction of fever (56). IL-6 is also responsible for lymphocyte activation and increases antibody production for the adaptive immune response (56). TNF $\alpha$  activates vascular endothelium and increases vascular permeability, which leads to increased entry of IgG, complement, and cells to tissues, and increases fluid drainage to lymph nodes (56). These cytokines are commonly used as experimental markers of an inflammatory response.

### ***COX- 2 and Anti-Inflammatory Drugs***

Cyclooxygenase (COX) catalyses the committed conversion of arachidonic acid to prostaglandin endoperoxide  $H_2$  ( $PGH_2$ ) and exists in two isoforms (61-66). These two isoforms of COX, i.e., COX-1 and COX-2, have different amino acid sequences but catalyze the same reaction. Both forms are membrane-bound, endoplasmic reticulum (ER)- resident, heme-containing glycoproteins that function as homodimers (63). COX-1 is constitutively expressed in most cell types (65, 66). In contrast, COX-2 expression is typically transient and is associated with inflammation (39, 64). The expression of COX-2 is a key element in various pathological conditions including arthritis, cardiovascular disease, and cancer (64).

Nonsteroidal anti-inflammatory drugs (NSAIDs) are a class of drugs that non-specifically target COX (61). They are prescribed for the treatment of inflammatory diseases such as arthritis and osteoarthritis. NSAIDs had initial success; however, long term therapy showed gastrointestinal (GI) toxicities due to the general inhibition of both COX-1 and COX-2. COX-1 expressed throughout the GI tract has protective effects on the GI mucosa by producing prostaglandin  $E_2$  ( $PGE_2$ ) (63). To avoid this side effect, research shifted toward selective COX-2 inhibitors. Selective inhibitors for COX-2, a class of drugs known as coxibs, show a decrease in inflammation. While, COX-2 knock-out mice have a marked protection from intestinal and skin tumors (66). Conversely, over-expression of COX-2 is observed in various malignant tissues (62). Depending on the cell type, COX-2 expression can be rapidly induced by the bacterial endotoxin LPS, cytokines such as IL-1, IL-2, and  $TNF\alpha$ , and the tumor promoter, phorbol 12-myristate 13- acetate (PMA) (61). Though, even after stimulation, protein levels of COX-2 only



reach about 50% of COX-1 (61). After initial studies on COX-2 inhibitors, concerns arose regarding adverse cardiovascular events. Studies demonstrated that selective COX-2 inhibitors tip the natural balance between prothrombotic thromboxane A<sub>2</sub> and antithrombotic prostacyclin, which potentially increases the possibility of thrombosis (63). Research is on-going to find a selective COX-2 inhibitor that is not accompanied by severe side effects.

## **Fatty Acids**

In the western diet, lipids represent more than 40% of the daily caloric intake, while nutritional advice recommends to limit lipid intake to 30% (26, 67). Fatty acids in the body can be exogenously brought in from the diet or can be generated from acetyl-CoA in a process called de novo lipogenesis (4, 68). A high fat diet is understood to greatly contribute to the increase of obesity prevalence as fatty acids are stored in fat cells composing adipose tissue (69). Carbohydrate (i.e. glucose) can be converted through de novo synthesis to fatty acids, which can also be stored in the form of triacylglycerol in adipose tissue (68).

### ***Fatty Acids Metabolism***

Mechanisms for the absorption of fatty acids differ according to the size and characteristic of the fatty acid. Long-chain fatty acids mix with bile acids in the intestinal lumen to form micelle particles (70). Micelles interact with the brush border of the intestine to allow fatty acid to diffuse from the micelle into the enterocyte, moving down

a concentration gradient (70). Fatty acids can also enter the cell through protein transporters, such as fatty acid transport protein (FATP) (71). From the enterocyte, fatty acids are packaged into chylomicrons that are then excreted into the lymph system (4, 69). Conversely, small-chain fatty acids have the ability to move directly from the intestinal lumen into portal circulation where they bind and are transported by albumin to the liver (70).

The liver and adipose are the two most important tissues in fatty acid metabolism. The liver is responsible for the uptake and catabolism of fatty acids as well as their repackaging into very low density lipoprotein (VLDL) in the form of triacylglycerol (70). The liver can also generate fatty acid from non-lipid precursors such as glucose (68). Fatty acids are delivered to adipose cells for storage. They can be delivered by VLDL, which would carry fatty acids originating from the liver, whereas dietary fatty acids are delivered to adipose cells by chylomicrons (70).

### ***Fatty Acid Structure***

The basic structure of fatty acids is a hydrophobic polycarbon chain which can vary in chain length with different degrees of saturation. Short- and medium-chain fatty acids, those with less than 6 or 14 carbons, respectively, are mainly saturated and used predominantly for energy supply. Long-chain fatty acids can be saturated as well as mono- or polyunsaturated, depending on the number of double bonds. They can serve functions apart from energy supply, including structural components of cells, gene regulators, involvement in cell growth and differentiation, or modulation of lipid, carbohydrate, and protein metabolism (4). The two major classes of long-chain

polyunsaturated fatty acids (PUFA) are  $\omega$ -3 and  $\omega$ -6 fatty acids, where the  $\omega$ -number indicates the position of the first double bond counted from the methyl group of the hydrocarbon chain. Current recommendations are to consume a higher percentage of  $\omega$ -3 than  $\omega$ -6 PUFA (72). Linoleic acid (18:2,  $\omega$ -6) is an essential PUFA and the precursor for arachidonic acid (20:4,  $\omega$ -6), while  $\alpha$ -linolenic acid (18:3,  $\omega$ -3) is the precursor for eicosapentaenoic acid (20:5,  $\omega$ -3) and docosahexaenoic acid (22:6,  $\omega$ -3). PUFA contribute roughly 7% of total energy intake in the diets of adults with linoleic acid (18:2,  $\omega$ -6) being the most abundant in the diet (72). It has been established that fatty acids, specifically PUFA, can influence gene and protein expression (4, 26, 59, 73). Specifically, gamma- linoleic acid, which is converted in the body from linoleic acid by delta-6-desaturase, has the ability to inhibit the response generated by an inflammatory insult through inactivation of inflammatory pathways in RAW 264.7 macrophage (59). Unsaturated fatty acids, in general, are thought to have anti-inflammatory properties and many studies have given support to their inclusion in the diet (72, 74, 75).

### **Adipose Fatty Acid-Binding Protein, aP2**

Fatty-acid binding proteins (FABP) are cytoplasmic proteins with low molecular weight (~15 KDa) that bind fatty acids with high affinity in a 1:1 complex with an interior ligand binding domain (76). They are implicated as players in the inflammatory response as well as lipid signaling cascades (2, 3). FABP function as intracellular lipid chaperones with a role in import, storage, and export of fatty acids because they reversibly bind hydrophobic ligands, such as saturated/unsaturated long-chain fatty acids

with high affinity (2, 77). There are nine known FABP identified in different cell types (78). The most characterized FABP is adipocyte fatty acid-binding protein (aP2/FABP4) which is found in adipocytes at high levels and in macrophages to a lesser extent (2). Specific functions of aP2 other than a lipid chaperon are unclear; however, Thompson *et al.* (76) suggested that aP2 may act as a fatty acid sensor affecting cellular metabolism via protein-protein interactions as demonstrated by experiments involving Janus kinase 2 (JAK2) (76).

Mouse studies have shown that aP2 deficiency gives partial protection against atherosclerotic lesion formation in apolipoprotein E knockout mice, as well as systemic insulin resistance, dyslipidemia, and lipotoxicity in *aP2<sup>-/-</sup> ob/ob* mice (2, 6). Molecular mechanisms underlying the effect of aP2 on the pathogenesis of the diseases are still under investigation. aP2's involvement in these diseases may be, in part, from the ability of an inflammatory insult such as LPS to dramatically increase aP2 expression in RAW 264.7 macrophage; triglycerides in macrophage are also increased when treated with LPS (79). This paralleled increase suggests aP2 as an intermediary in the link between the immune and metabolism systems. Pharmacological agents specifically targeting aP2 are currently being investigated as a treatment for diabetes and atherosclerosis (3, 12).

An AP-1 sequence (80, 81), and C/EBP binding element (10, 11) have both been suggested as cis-acting elements of the aP2 gene. It was demonstrated that macrophage aP2 can be regulated by phorbol 12-myristate 13-acetate (PMA), LPS, oxidized low-density lipoprotein and PPAR $\gamma$  ligands (12). The mechanisms for how aP2 expression is controlled under basal and activated conditions in macrophage is still unknown.

## **Histone Modifications**

Eukaryotic cell DNA is packaged into structures called nucleosomes (82). Each nucleosome is comprised of eight core histone proteins and the associated wrapped DNA (82). Histones are small, positively charged proteins and have close association with DNA. Eukaryotic cells commonly contain five abundant histone proteins: H1, H2A, H2B, H3, and H4 (82). H2A, H2B, H3, and H4 are considered the core histones because two copies of each form the protein core around which the nucleosome DNA is wrapped (82). H1 binds to the linker DNA and is referred to as a linker histone. Histone proteins tend to have a high concentration of positively charged amino acids, including lysine and arginine (82). By assembling into nucleosomes, DNA is approximately compacted 6-fold (82). The core DNA is a 147 base pair section that is wrapped around the histone octamer and then linked to the next histone octamer by a variable 20 – 60 base pair region of DNA (82).

Chromatin greatly impedes transcription. Epigenetic modifications of chromatin and DNA have been recognized as important permissive and suppressive factors in controlling gene transcription (83). Epigenetic mechanisms result in heritable modification in the expression of genes that are independent from DNA coding variability (83). Two major epigenetic mechanisms are the post-translational modification of histone proteins in chromatin and the methylation of DNA (83). Post-translational modifications of histones include, but are not limited to, lysine acetylation, lysine and arginine methylation, serine and threonine phosphorylation, and lysine ubiquitination and sumoylation (84). Emerging evidence suggest a role of epigenetics in human pathologies, including inflammation (83). Additionally, it is known that nutrition and environment have the ability to influence the epigenetic profile.

### *Histone Deacetylases*

One way to affect gene expression is through the involvement of coactivator/corepressor complexes (13). Histone deacetylases (HDAC) are essential components of co-repressor complexes as they provide the enzymatic activity for active repression (14). HDAC are a class of enzymes that remove acetyl groups from an  $\epsilon$ -N-acetyl lysine amino acid on a histone (15). Its action is opposite to that of histone acetyltransferase (HAT) which act as coactivators (15). Removal of acetyl groups gives chromatin a tightly-bound structure that prohibits transcription. Inclusion of acetyl groups decreases the affinity of histones for DNA and allows transcription to occur (15). There are eighteen known HDAC, which are divided into four classes (15). Class 1 HDAC are generally detected in the nucleus and ubiquitously expressed (15). Specifically, HDAC3 is unique among class 1 HDAC in that it can shuttle between the nucleus and cytoplasm and form complexes (15).

Transcription factors that have been shown to interact with the corepressor complex for repressive effects on gene transcription include AP-1 (16-18), NF- $\kappa$ B (16, 17, 19), liver X receptor  $\alpha$  (20), and thyroid hormone receptor (13, 21, 22). Upon activation, the co-repressor complex is exchanged with co-activators that lead to transcription (14). Using chromatin immunoprecipitation (ChIP), it was shown that high glucose conditions, similar to a diabetic state, could increase the recruitment of HAT to the promoter regions of inflammatory genes such as TNF $\alpha$  and COX-2 and lead to acetylation of nucleosomal factors histone H3 and H4 (54).

HDAC inhibitors are currently under investigation as anti-cancer agents because overexpression of HDACs has been observed in various cancers (15). Trichostatin A (TSA) is a potent and nonselective inhibitor of HDAC (15, 85). TSA treatment has been seen to change the expression of only about 2% of genes (85). TSA is widely used to study the role of histone acetylation on gene expression.

## Materials and Methods

### *Cell Culture and Fatty Acid Preparation*

Murine RAW 264.7 macrophages (ATCC) were cultured in RPMI 1640 supplemented with 10% fetal bovine serum, 100 U/ml penicillin/ 100 ng/ml streptomycin, 1x vitamins, and 2 mM L-glutamine in a humidified chamber at 37°C with 5% CO<sub>2</sub>. Cells were plated at  $5 \times 10^5$  per well in a 12-well plate to extract RNA and  $1.5 \times 10^6$  per well in a 6-well plate for protein extraction. All cell culture supplies were purchased from MediaTech.

Two mmol/L of fatty acid-poor and endotoxin-free bovine serum albumin (BSA; Calbiochem) was prepared in phosphate buffered saline (PBS). Sodium salts of fatty acids (Nu-Chek) were dissolved in the 2 mmol/L BSA solution to a final concentration of 5 mmol/L. Fatty acids prepared included: myristic acid (14:0), palmitic acid (16:0), palmitoleic acid (16:1), oleic acid (18:1), linoleic acid (18:2), linolenic acid (18:3), arachidonic acid (20:4), eicosapentaenoic acid (20:5) and docosahexaenoic acid (22:6). The fatty acid and BSA mixture was purged with N<sub>2</sub>, sonicated in a cool water bath until the solution became clear to form BSA/fatty acid complex (approximate molar ratio = 1:2.5), filter sterilized through a Millex®-GV 0.22 µm filter unit (Millipore), and diluted with cell medium to reach a final concentration of 100 µmol/L.

Raw 264.7 macrophages were incubated with BSA only (control) or 100 µM of a fatty acid for 6 h followed by BSA or fatty acid with 100 ng/mL LPS. For the inhibition of HDAC, RAW 264.7 macrophages were incubated in RPMI 1640 containing 500 nM TSA (Sigma-Aldrich) dissolved in dimethyl sulfoxide (DMSO; Sigma-Aldrich) and 100



$\mu\text{M}$  fatty acids for 6 hours followed by 14 hours in RPMI 1640 containing 500 nM TSA, 100  $\mu\text{M}$  fatty acid, and 100 ng/ml LPS. For the experiments using the inhibitor, control cells were incubated with the same amount of DMSO vehicle. Macrophage activation was by LPS and  $\text{TNF}\alpha$ . Cells were incubated with 100 ng/mL LPS (Sigma-Aldrich) or 2.5ng/ml, 5.0ng/ml, and 10ng/ml  $\text{TNF}\alpha$  (eBioscience) for 18 hours and RNA was collected and analyzed by real-time PCR.

### ***Small Interfering RNA (siRNA) Transfection***

RAW 264.7 macrophages were transfected with Silencer<sup>®</sup> Negative Control siRNA (Ambion) (control) or Dharmacon siGenome SMART pool siRNA targeting aP2 (Thermo Scientific) for gene knockdown using DharmaFECT<sup>®</sup> 1 transfection reagent (Thermo Scientific) according to the manufacture's protocol. . In brief, 2.5  $\mu\text{l}$  of transfection reagent DharmaFECT 1 was diluted into 97.5  $\mu\text{l}$  of cell medium void of antibiotics and fetal bovine serum. Control and siRNA (2 or 3  $\mu\text{mol/L}$ ) were prepared in RNase-free sterile water and 50  $\mu\text{l}$  of the diluted siRNA solution was mixed with 50 $\mu\text{l}$  of cell medium void of antibiotics and fetal bovine serum. Subsequently, the media containing transfection agent and siRNA were combined and incubated for 20 min at room temperature. The siRNA and transfection agent complex was then diluted to desired volume (1 ml). When recovering protein, reaction volumes were doubled and diluted siRNA concentration was 2.5 $\mu\text{mol/L}$ . Transfection occurred for 24 hours, after which cells were incubated with 100  $\mu\text{M}$  fatty acids for 6 hours and then 100  $\mu\text{M}$  fatty acid and 100 ng/ml LPS for 18 hours.

### ***Total RNA Isolation and Quantitative Real-time PCR***

Total RNA was extracted from cells using TRIzol reagent (Invitrogen) following manufacturer's protocol or using the RNeasy Kit (Qiagen). Reverse transcription for cDNA synthesis and quantitative real-time PCR analysis were performed as previously described (86, 87). Primers were designed according to GenBank database using the Primer Express software (Applied Biosystems). The following primers were used for real-time PCR analysis: GAPDH, aP2, COX2, IL-1 $\beta$ , IL-6, and TNF $\alpha$  (Table 1).

### ***HDAC Activity***

Raw 264.7 macrophages were transfected with scrambled control or aP2 siRNA and incubated with 100  $\mu$ M fatty acid for 6 h then 100 ng/ml LPS was added to fatty acids for an additional 18 h. Nuclear extracts were collected using Nuclear Extract Kit (Active Motif) according to manufacturer's protocol. Protein concentration was measured using BCA Protein Assay Kit (Thermo Scientific) and 10  $\mu$ g of nuclear extract was analyzed for HDAC activity using a HDCA Activity Assay Kit (Cayman Chemical) according to the manufacturer's protocol.

### ***Western Blot Analysis***

Cell lysate was prepared and Western blot analysis was performed as previously described (88). Antibodies used for analysis were as follow: aP2/FABP4 (Santa-Cruz), and  $\beta$ -actin (Sigma-Aldrich).  $\beta$ -Actin was used as a loading control to normalize the data.

### ***Statistical Analysis***

ANOVA and Newman-Keuls Multiple comparison with Welch's correction for unequal variance when appropriate were used to identify statistically significant differences of treatments with  $P < 0.05$  considered significant by GraphPad InStat 5 (GraphPad Software, Inc.). Data are expressed as mean  $\pm$  SEM.

**Table 1.** Real-time PCR primers for mRNA expression

	Forward Primer 5' → 3'	Reverse Primer 5' → 3'
GAPDH	TGTGTCCGTCGTGGATCTGA	CCTGCTTCACCACCTTCTTGAT
aP2	CACCATCCGGTCAGAGAGTACTT	CGGTGATTTTCATCGAATTCCA
COX2	AAAGGTTCTTCTACGGAGAGAGTTCA	TGGGCAAAGAATGCAAACATC
IL-1 $\beta$	GTCACAAGAAACCATGGCACAT	GCCCATCAGAGGCAAGGA
IL-6	CTGCAAGAGACTTCCATCCAGTT	AGGGAAGGCCGTGGTTGT
TNF $\alpha$	GGCTGCCCCGACTACGT	ACTTTCTCCTGGTATGAGATAGCAAAT


## Results

### ***Expression of aP2 was induced by LPS stimulation in RAW 264.7 macrophages***

Raw 264.7 macrophages were treated with 100 ng/ml LPS or 2.5 ng/ml, 5.0 ng/ml, and 10 ng/ml TNF $\alpha$  for 18 hours to evaluate the effect of an inflammatory insult on aP2 expression and quantitative real-time PCR analysis was conducted. LPS drastically increased aP2 mRNA abundance by ~8-fold compared with the control (Figure 1). Cells treated with TNF $\alpha$  also showed a ~2-fold increase in aP2 mRNA expression relative to the control (data not shown).


### ***Unsaturated fatty acids (UFA) inhibited LPS-induced aP2 expression in RAW 264.7 macrophages***

To investigate whether fatty acids could alter aP2 expression in response to an inflammatory insult, RAW 264.7 macrophage were incubated with fatty acids including myristic acid (14:0), palmitic acid (16:0), palmitoleic acid (16:1), oleic acid (18:1), linoleic acid (18:2), linolenic acid (18:3), arachidonic acid (20:4), eicosapentaenoic acid (20:5) and docosahexaenoic acid (22:6) in the absence or presence of LPS. Consistent with previous data in Figure 1, LPS increased aP2 expression in the BSA control. Without LPS treatment, no significant difference in aP2 mRNA levels was seen in all the fatty acid treatment groups (Figure 2A). In contrast, when cells were activated by LPS, unsaturated fatty acids significantly decreased aP2 expression whereas saturated fatty acids did not alter the expression compared with the control. Western blot analysis confirmed the inhibition of LPS-induced aP2 protein levels by unsaturated fatty acids




**Figure 1. Elevated aP2 mRNA levels by LPS in RAW 264.7 macrophages.** Cells were treated with 100 ng/ml LPS for 18 hours, after which quantitative real-time PCR was conducted to measure aP2 mRNA abundance.  $n = 9$ , Mean  $\pm$  SEM. \*,  $P < 0.001$ .

A



B



**Figure 2. Elevated aP2 expression by LPS was abolished by unsaturated fatty acids in RAW 264.7 macrophages.** Cells were treated with 100  $\mu$ M fatty acid complexed with BSA for 12 hours and subsequently activated by 100 ng/ml LPS for 18 hours. A. aP2 mRNA expression was measured by real time PCR.  $n=3-4$ , Mean  $\pm$  SEM. Bars with different letters are significantly different compared with control within the absence or presence of LPS treatment ( $P < 0.05$ ). B. aP2 protein was measured by western blot analysis and  $\beta$ -actin was used as a loading control.

(Figure 2B). Palmitic acid markedly increased aP2 protein levels in the presence of LPS despite no significant change in mRNA levels.


### ***TSA abrogated UFA-mediated repression of basal and LPS-induced aP2 expression***

To evaluate whether alterations in histone acetylation state is involved in the inhibitory effect of UFA on aP2 expression, RAW 264.7 macrophages were treated with TSA, an HDAC inhibitor (15), together with fatty acids, after which aP2 mRNA levels were quantified by real-time PCR. In consistent with our previous observations, LPS increased aP2 mRNA levels and 18:2 significantly lowered basal as well as LPS-induced aP2 expression in the absence of TSA (Figure 3). In the presence of TSA, however, the repressive effect of 18:2 on basal and LPS-induced aP2 expression was abolished. The data indicate that 18:2 could inhibit basal and LPS-induced aP2 expression by modulating HDAC.


### ***Palmitic acid decreased HDAC activity in RAW 264.7 macrophages***

To evaluate the effect of fatty acids and aP2 on HDAC activity, Raw 264.7 macrophages were transfected with scrambled control or siRNA targeting aP2 for 24 h and then treated with 100  $\mu$ M fatty acid for 24 h. aP2 protein levels were markedly diminished by siRNA in all the groups (Figure 4A). Compared to control, 16:0 significantly reduced HDAC activity regardless of aP2 deficiency (Figure 4B). Although knockdown of aP2 did not alter HDAC activity within each fatty acid treatment, HDAC activity was significantly elevated by 18:2 compared with control when aP2 was deficient.






**Figure 3. Abrogation of repression of basal and LPS-induced aP2 mRNA levels by TSA in RAW 264.7 macrophages.** Cells were treated with 100  $\mu$ M fatty acid and 500 nM TSA for 6 hours, after which 100 ng/ml of LPS was added for additional 18 hours. aP2 mRNA levels was measured by quantitative real-time PCR analysis. n=8-9, Mean  $\pm$  SEM. \*,  $P < 0.05$  compared with control in the same treatment. \$,  $P < 0.05$  compared with 18:2 in the same treatment.




**Figure 4. Inhibition of HDAC Activity by palmitic acid in RAW 264.7 macrophages.** Cells were transfected with 125 nM of scrambled control (sc) or siRNA targeting aP2 for 24 h. They were subsequently treated with 100  $\mu$ M fatty acid for 6 h after which 100 ng/ml LPS was added for additional 18 h. A. Nuclear extracts were collected to measure HDAC activity  $n = 4$ , Mean  $\pm$  SEM. B. Cytoplasmic fractions were subjected to Western blot analysis to assess aP2 knockdown.  $\beta$ -actin was used as a loading control.

***COX-2 was repressed by aP2 knockdown in LPS- stimulated RAW 2647 macrophages***

To investigate whether aP2 plays a role in the repressed pro-inflammatory gene expression by unsaturated fatty acids, RAW 264.7 macrophages were transfected with scrambled control or aP2 siRNA to knockdown aP2 expression. Cells were then treated with 100  $\mu$ M of 16:0 and 18:2 for 6 h, after which 100 ng/ml LPS was added to cell culture medium to activate macrophages for 18 h. Real-time PCR analysis was performed to measure mRNA levels of aP2 and pro-inflammatory markers including IL-1 $\beta$ , IL-6, TNF $\alpha$ , and COX-2. aP2 mRNA levels showed a reduction by 75-80% by siRNA transfection (Figure 5) . In the unstimulated macrophages, deficiency of aP2 by siRNA increased the pro-inflammatory mediator expression and the increase was drastic particularly in 16:0-treated cells (Figure 6). In contrast, when cells were stimulated by LPS, aP2 knockdown had a minimal effect on IL-1 $\beta$ , TNF $\alpha$  and IL-6 mRNA levels. Of interest was that COX-2 mRNA abundance were significantly reduced when aP2 was deficient.



**Figure 5. Knockdown of aP2 using siRNA in RAW 264.7 macrophages.** Cells were transfected with 100 nM of scrambled control or aP2 siRNA and then treated with 100  $\mu$ M fatty acid for 6 h. Subsequently, cells were activated by 100 ng/ml of LPS for 18 h. aP2 mRNA levels were measured by real-time PCR. n= 3, Mean  $\pm$  SEM



**Figure 6. Effect of aP2 and FA on pro-inflammatory gene expression in RAW 264.7 macrophages.** Cells were transfected with 100 nM of scrambled control or aP2 siRNA and then treated with 100  $\mu$ M fatty acid for 6 hours. Subsequently, cells were activated by 100 ng/ml of LPS for 18 hours. Expression of pro-inflammatory mediators including IL-6, IL-1 $\beta$ , TNF $\alpha$  and COX-2 were measured by real-time PCR. n = 3, Mean  $\pm$  SEM. \*,  $P < 0.05$  compared with scrambled control in the same fatty acid treatment. #,  $P < 0.05$  compared with BSA control.

## Discussion

aP2 deficiency in mice led the protection against the development of IR (89-91). Total and macrophage-specific aP2 knockout mice showed a marked reduction in atherosclerotic lesion formation in apolipoprotein E knockout (*apoE*<sup>-/-</sup>) mice (92-95). In humans, a functional genetic variant of aP2 gene, resulting in reduced aP2 expression, showed a significantly lowered risk for Type 2 diabetes and CVD (96). Additionally, administration of an aP2 inhibitor in ob/ob and *apoE*<sup>-/-</sup> mice reduced the development of type 2 diabetes and atherosclerosis, respectively (12). The health benefits from absence or reduction of aP2 are linked to inhibited inflammatory mediator production in macrophages. We found that aP2 expression in macrophages is repressed by unsaturated fatty acids and acetylation of histones is likely to play a role in this repression. In addition, aP2 may contribute to the induction of COX-2 expression by inflammatory insults in macrophages.

In murine macrophage cell line, absence of aP2 markedly diminished pro-inflammatory cytokines (92). When THP-1 macrophages were incubated with the aP2 inhibitor, the production of pro-inflammatory cytokines including monocyte chemoattractant protein-1 (MCP-1), IL-1 $\beta$  and IL-6 was significantly reduced (12). These studies suggest a significant role of aP2 in inflammatory signaling in macrophages. In the present study, both LPS and TNF $\alpha$  significantly increased aP2 expression in RAW 264.7 macrophages. Although cell surface receptors for LPS and TNF $\alpha$  are different, downstream mediators that can transduce their signals can converge onto similar pathways, including I Kappa B Kinases (IKK)/NF- $\kappa$ B and MAPK/AP-1 pathways.

Whether the induction of aP2 expression by LPS and TNF $\alpha$  is mediated through the pathways needs further investigation.

aP2 was initially thought to be controlled by PPAR $\gamma$  in adipocytes (97). It has been suggested by Genolet *et al.* (36) that PPAR $\gamma$  agonists, such as long-chain fatty acids, may regulate inflammatory pathways via both PPAR $\gamma$ -dependent and independent mechanisms. The observation that LPS increases aP2 in the present study is counterintuitive as LPS is known to inhibit PPAR $\gamma$  transcriptional activity (98). If PPAR $\gamma$  represents the major regulator for aP2 expression in macrophages, down-regulation of aP2 by LPS would be expected. Thus, another mechanism may exist by which LPS induces aP2 expression independent of PPAR $\gamma$ . AP-1 and C/EBP are shown to regulate aP2 expression in adipocytes and preadipocytes (80, 81, 99, 100). AP-1 is a more likely candidate for a transcriptional regulator of macrophage aP2 responding to inflammatory stimuli for the following reasons: first, AP-1 is one of the major transcriptional factors that are sensitive to inflammatory insults to produce pro-inflammatory mediators (101, 102); second, aP2 promoter contains AP-1 binding element; and third, LPS stimulation is shown to trigger signaling pathways leading to activation of MAPK (103-108) that can, in turn, activate AP-1. Therefore, MAPK/AP-1 pathway could be involved in the regulation of aP2 expression in macrophages.

Basal as well as LPS-induced aP2 expression in RAW 264.7 macrophage was repressed by unsaturated fatty acids but not by saturated fatty acids. Therefore, we sought to understand the mechanism for how unsaturated fatty acids can alter aP2 expression. Posttranslational modifications in histones such as acetylation, phosphorylation and ubiquitination are an important way to alter gene expression. Of the modifications,

histone acetylation is best understood. To gain an insight into a potential role of HDAC in the fatty acid regulation of aP2 expression, RAW 264.7 macrophages were treated with TSA. We observed that the repressive effects of the unsaturated fatty acids on the basal as well as LPS-induced aP2 expression were abrogated in the presence of TSA. The result is supportive to the idea that histone modifications, particularly acetylation, can be a mechanism by which fatty acids can alter aP2 expression in RAW 264.7 macrophages. Of interest is that basal aP2 expression was markedly increased by TSA regardless of the presence or absence of fatty acids suggesting that aP2 expression in RAW 264.7 macrophages is likely to be under active transrepression. Transcriptional factors namely AP-1, NF- $\kappa$ B and LXR $\alpha$  (17, 19, 109-111) have shown to exert active transrepression of gene expression through association with a co-repressor complex when they are not activated by their agonists. Nuclear receptor corepressors (NCoR) and silencing mediator for retinoid and thyroid receptor (SMRT) are essential components of a co-repressor complex and HDAC3 is a well-known HDAC isoform to interact with NCoR and SMRT (111-114). Therefore, we speculate that the co-repressor complex may be associated with an unknown transcription factor in the promoter of aP2 to repress basal gene expression in macrophages; and upon LPS stimulation, the co-repressor complex is dissociated to induce gene expression.

HDAC activity was measured to assess whether fatty acids alter the activity. Saturated fatty acids lowered HDAC activity. This observation is consistent with the observation that saturated fatty acids can increase transcription of inflammatory mediated such as COX-2 (39), in that a decrease in HDAC activity would coordinate with an increase in transcription. Furthermore, saturated fatty acids lowering HDAC activity is



also consistent with a trend seen in our data that saturated fatty acids can increase aP2 expression (data not shown). A similar parallel is seen with unsaturated fatty acids increasing HDAC activity. When macrophages were transfected with aP2 siRNA and treated with 18:2, there was a significant increase in HDAC activity. An increase in HDAC activity would coordinate with a repression in transcription. The trend toward significant for the negative control transfected cells treated with unsaturated fatty acid would suggest that unsaturated fatty acids are capable of affecting the HDAC activity. There was no significant differences in HDAC activity between the negative control and aP2 transected cells but this could be due to the fact the HDAC activity kit used measured universal HDAC activity and not a specific isoform. HDAC1 is the most abundance HDAC isoform in macrophage (data not shown), it is possible that fatty acids may not effect HDAC1 activity but could affect HDAC3 activity, which is the isoform believed to be part of the corepressor complexes that can affect transcription (115). An assay to measure universal HDAC activity would see HDAC1 overshadowing any effects that could be present in HDAC3. However, we speculate that HDAC3 as part of a co-repressor complex is influenced to continue or increase repression by unsaturated fatty acids but alternatively removed in the presence of saturated fatty acids. The absence of aP2 could further promote unsaturated fatty acids ability to influence the co-repressor complex.

In summary, we have shown a significant up-regulation of aP2 by an inflammatory insult, that unsaturated fatty acid repress both basal and induced aP2 expression, and that palmitic acid reduces HDAC activity in activated RAW 264.7 macrophage. This evidence suggests the involvement of HDAC and histone

deacetylation state in the mechanisms for how unsaturated fatty acids modulate a repression in LPS-activated macrophage. Lastly, a relationship between COX-2 and aP2 could be part of the mechanism for the anti-inflammatory effects of unsaturated fatty acids seen in macrophage. We have provided here new information on the mechanisms by which unsaturated fatty acids can regulate aP2 expression. Due to the high frequency of obesity in the United States, deciphering the mechanism for the anti-inflammatory properties of unsaturated fatty acid will allow for more appropriate and precise dietary interventions to prevent chronic metabolic and inflammatory diseases.

## Literature Cited

1. Hotamisligil GS. Inflammation and metabolic disorders. *Nature*. 2006 Dec 14;444:860-7.
2. Makowski L, Hotamisligil GS. Fatty acid binding proteins--the evolutionary crossroads of inflammatory and metabolic responses. *J Nutr*. 2004 Sep;134:2464S-8S.
3. Makowski L, Hotamisligil GS. The role of fatty acid binding proteins in metabolic syndrome and atherosclerosis. *Curr Opin Lipidol*. 2005 Oct;16:543-8.
4. Jump DB. Fatty acid regulation of gene transcription. *Crit Rev Clin Lab Sci*. 2004;41:41-78.
5. Lavie CJ, Milani RV, Mehra MR, Ventura HO. Omega-3 polyunsaturated fatty acids and cardiovascular diseases. *J Am Coll Cardiol*. 2009 Aug 11;54:585-94.
6. Makowski L, Boord JB, Maeda K, Babaev VR, Uysal KT, Morgan MA, Parker RA, Suttles J, Fazio S, et al. Lack of macrophage fatty-acid-binding protein aP2 protects mice deficient in apolipoprotein E against atherosclerosis. *Nat Med*. 2001 Jun;7:699-705.
7. Guilherme A, Virbasius JV, Puri V, Czech MP. Adipocyte dysfunctions linking obesity to insulin resistance and type 2 diabetes. *Nat Rev Mol Cell Biol*. 2008 May;9:367-77.
8. Kanda H, Tateya S, Tamori Y, Kotani K, Hiasa K, Kitazawa R, Kitazawa S, Miyachi H, Maeda S, et al. MCP-1 contributes to macrophage infiltration into adipose tissue, insulin resistance, and hepatic steatosis in obesity. *J Clin Invest*. 2006 Jun;116:1494-505.
9. Aeberli I, Beljean N, Lehmann R, l'Allemand D, Spinaz GA, Zimmermann MB. The increase of fatty acid-binding protein aP2 in overweight and obese children: interactions with dietary fat and impact on measures of subclinical inflammation. *Int J Obes (Lond)*. 2008 Oct;32:1513-20.
10. Christy RJ, Yang VW, Ntambi JM, Geiman DE, Landschulz WH, Friedman AD, Nakabeppu Y, Kelly TJ, Lane MD. Differentiation-induced gene expression in 3T3-L1 preadipocytes: CCAAT/enhancer binding protein interacts with and activates the promoters of two adipocyte-specific genes. *Genes Dev*. 1989 Sep;3:1323-35.
11. Herrera R, Ro HS, Robinson GS, Xanthopoulos KG, Spiegelman BM. A direct role for C/EBP and the AP-1-binding site in gene expression linked to adipocyte differentiation. *Mol Cell Biol*. 1989 Dec;9:5331-9.

12. Furuhashi M, Tuncman G, Gorgun CZ, Makowski L, Atsumi G, Vaillancourt E, Kono K, Babaev VR, Fazio S, et al. Treatment of diabetes and atherosclerosis by inhibiting fatty-acid-binding protein aP2. *Nature*. 2007 Jun 21;447:959-65.
13. Naar AM, Lemon BD, Tjian R. Transcriptional coactivator complexes. *Annu Rev Biochem*. 2001;70:475-501.
14. Bailey ST, Ghosh S. 'PPAR'ting ways with inflammation. *Nat Immunol*. 2005 Oct;6:966-7.
15. Bolden JE, Peart MJ, Johnstone RW. Anticancer activities of histone deacetylase inhibitors. *Nat Rev Drug Discov*. 2006 Sep;5:769-84.
16. Lee SK, Kim JH, Lee YC, Cheong J, Lee JW. Silencing mediator of retinoic acid and thyroid hormone receptors, as a novel transcriptional corepressor molecule of activating protein-1, nuclear factor-kappaB, and serum response factor. *J Biol Chem*. 2000 Apr 28;275:12470-4.
17. Perissi V, Aggarwal A, Glass CK, Rose DW, Rosenfeld MG. A corepressor/coactivator exchange complex required for transcriptional activation by nuclear receptors and other regulated transcription factors. *Cell*. 2004 Feb 20;116:511-26.
18. Zhang J, Kalkum M, Chait BT, Roeder RG. The N-CoR-HDAC3 nuclear receptor corepressor complex inhibits the JNK pathway through the integral subunit GPS2. *Mol Cell*. 2002 Mar;9:611-23.
19. Baek SH, Ohgi KA, Rose DW, Koo EH, Glass CK, Rosenfeld MG. Exchange of N-CoR corepressor and Tip60 coactivator complexes links gene expression by NF-kappaB and beta-amyloid precursor protein. *Cell*. 2002 Jul 12;110:55-67.
20. Wagner BL, Valledor AF, Shao G, Daige CL, Bischoff ED, Petrowski M, Jepsen K, Baek SH, Heyman RA, et al. Promoter-specific roles for liver X receptor/corepressor complexes in the regulation of ABCA1 and SREBP1 gene expression. *Mol Cell Biol*. 2003 Aug;23:5780-9.
21. McKenna NJ, O'Malley BW. Combinatorial control of gene expression by nuclear receptors and coregulators. *Cell*. 2002 Feb 22;108:465-74.
22. Rosenfeld MG, Glass CK. Coregulator codes of transcriptional regulation by nuclear receptors. *J Biol Chem*. 2001 Oct 5;276:36865-8.
23. Stienstra R, Duval C, Muller M, Kersten S. PPARs, Obesity, and Inflammation. *PPAR Res*. 2007;2007:95974.

24. Flegal KM, Carroll MD, Ogden CL, Curtin LR. Prevalence and trends in obesity among US adults, 1999-2008. *JAMA*. Jan 20;303:235-41.
25. Vachharajani V, Granger DN. Adipose tissue: a motor for the inflammation associated with obesity. *IUBMB Life*. 2009 Apr;61:424-30.
26. Oller do Nascimento CM, Ribeiro EB, Oyama LM. Metabolism and secretory function of white adipose tissue: effect of dietary fat. *An Acad Bras Cienc*. 2009 Sep;81:453-66.
27. Rajala MW, Scherer PE. Minireview: The adipocyte--at the crossroads of energy homeostasis, inflammation, and atherosclerosis. *Endocrinology*. 2003 Sep;144:3765-73.
28. Bastard JP, Maachi M, Lagathu C, Kim MJ, Caron M, Vidal H, Capeau J, Fève B. Recent advances in the relationship between obesity, inflammation, and insulin resistance. *Eur Cytokine Netw*. 2006 Mar;17:4-12.
29. Curat CA, Miranville A, Sengenès C, Diehl M, Tonus C, Busse R, Bouloumié A. From blood monocytes to adipose tissue-resident macrophages: induction of diapedesis by human mature adipocytes. *Diabetes*. 2004 May;53:1285-92.
30. Avogaro A, de Kreutzenberg SV. Mechanisms of endothelial dysfunction in obesity. *Clin Chim Acta*. 2005 Oct;360:9-26.
31. Chen X, Xun K, Chen L, Wang Y. TNF- $\alpha$ , a potent lipid metabolism regulator. *Cell Biochem Funct*. 2009 Oct;27:407-16.
32. Alexandraki K, Piperi C, Kalofoutis C, Singh J, Alaveras A, Kalofoutis A. Inflammatory process in type 2 diabetes: The role of cytokines. *Ann N Y Acad Sci*. 2006 Nov;1084:89-117.
33. Weisberg SP, McCann D, Desai M, Rosenbaum M, Leibel RL, Ferrante AW, Jr. Obesity is associated with macrophage accumulation in adipose tissue. *J Clin Invest*. 2003 Dec;112:1796-808.
34. Xu H, Barnes GT, Yang Q, Tan G, Yang D, Chou CJ, Sole J, Nichols A, Ross JS, et al. Chronic inflammation in fat plays a crucial role in the development of obesity-related insulin resistance. *J Clin Invest*. 2003 Dec;112:1821-30.
35. Coppack SW. Pro-inflammatory cytokines and adipose tissue. *Proc Nutr Soc*. 2001 Aug;60:349-56.
36. Tsukumo DM, Carvalho-Filho MA, Carvalheira JB, Prada PO, Hirabara SM, Schenka AA, Araujo EP, Vassallo J, Curi R, et al. Loss-of-function mutation in

- Toll-like receptor 4 prevents diet-induced obesity and insulin resistance. *Diabetes*. 2007 Aug;56:1986-98.
37. Chow JC, Young DW, Golenbock DT, Christ WJ, Gusovsky F. Toll-like receptor-4 mediates lipopolysaccharide-induced signal transduction. *J Biol Chem*. 1999 Apr 16;274:10689-92.
  38. Lapara NJ, 3rd, Kelly BL. Suppression of LPS-induced inflammatory responses in macrophages infected with *Leishmania*. *J Inflamm (Lond)*. 7:8.
  39. Lee JY, Sohn KH, Rhee SH, Hwang D. Saturated fatty acids, but not unsaturated fatty acids, induce the expression of cyclooxygenase-2 mediated through Toll-like receptor 4. *J Biol Chem*. 2001 May 18;276:16683-9.
  40. Skaug B, Jiang X, Chen ZJ. The role of ubiquitin in NF-kappaB regulatory pathways. *Annu Rev Biochem*. 2009;78:769-96.
  41. Szanto A, Nagy L. The many faces of PPARgamma: anti-inflammatory by any means? *Immunobiology*. 2008;213:789-803.
  42. Genolet R, Wahli W, Michalik L. PPARs as drug targets to modulate inflammatory responses? *Curr Drug Targets Inflamm Allergy*. 2004 Dec;3:361-75.
  43. Tontonoz P, Spiegelman BM. Fat and beyond: the diverse biology of PPARgamma. *Annu Rev Biochem*. 2008;77:289-312.
  44. Chawla A, Barak Y, Nagy L, Liao D, Tontonoz P, Evans RM. PPAR-gamma dependent and independent effects on macrophage-gene expression in lipid metabolism and inflammation. *Nat Med*. 2001 Jan;7:48-52.
  45. Stienstra R, Duval C, Keshtkar S, van der Laak J, Kersten S, Muller M. Peroxisome proliferator-activated receptor gamma activation promotes infiltration of alternatively activated macrophages into adipose tissue. *J Biol Chem*. 2008 Aug 15;283:22620-7.
  46. Ricote M, Li AC, Willson TM, Kelly CJ, Glass CK. The peroxisome proliferator-activated receptor-gamma is a negative regulator of macrophage activation. *Nature*. 1998 Jan 1;391:79-82.
  47. Jiang C, Ting AT, Seed B. PPAR-gamma agonists inhibit production of monocyte inflammatory cytokines. *Nature*. 1998 Jan 1;391:82-6.
  48. Pascual G, Fong AL, Ogawa S, Gamliel A, Li AC, Perissi V, Rose DW, Willson TM, Rosenfeld MG, Glass CK. A SUMOylation-dependent pathway mediates

- transrepression of inflammatory response genes by PPAR-gamma. *Nature*. 2005 Sep 29;437:759-63.
49. Ye J. Regulation of PPARgamma function by TNF-alpha. *Biochem Biophys Res Commun*. 2008 Sep 26;374:405-8.
  50. Tan NS, Shaw NS, Vinckenbosch N, Liu P, Yasmin R, Desvergne B, Wahli W, Noy N. Selective cooperation between fatty acid binding proteins and peroxisome proliferator-activated receptors in regulating transcription. *Mol Cell Biol*. 2002 Jul;22:5114-27.
  51. Helledie T, Antonius M, Sorensen RV, Hertzog AV, Bernlohr DA, Kolvaara S, Kristiansen K, Mandrup S. Lipid-binding proteins modulate ligand-dependent trans-activation by peroxisome proliferator-activated receptors and localize to the nucleus as well as the cytoplasm. *J Lipid Res*. 2000 Nov;41:1740-51.
  52. Marion-Letellier R, Dechelotte P, Iacucci M, Ghosh S. Dietary modulation of peroxisome proliferator-activated receptor gamma. *Gut*. 2009 Apr;58:586-93.
  53. Wahli W. A gut feeling of the PXR, PPAR and NF-kappaB connection. *J Intern Med*. 2008 Jun;263:613-9.
  54. Miao F, Gonzalo IG, Lanting L, Natarajan R. In vivo chromatin remodeling events leading to inflammatory gene transcription under diabetic conditions. *J Biol Chem*. 2004 Apr 23;279:18091-7.
  55. Kim JH, Park YM, Shin JS, Park SJ, Choi JH, Jung HJ, Park HJ, Lee KT. Fraxinellone inhibits lipopolysaccharide-induced inducible nitric oxide synthase and cyclooxygenase-2 expression by negatively regulating nuclear factor-kappa B in RAW 264.7 macrophages cells. *Biol Pharm Bull*. 2009 Jun;32:1062-8.
  56. Janeway C. *Janeway's Immunobiology*. 7th ed. New York: Taylor & Francis Group, LLC; 2008.
  57. Feng Y, Wen J, Chang CC. p38 Mitogen-activated protein kinase and hematologic malignancies. *Arch Pathol Lab Med*. 2009 Nov;133:1850-6.
  58. Yang R, Trevillyan JM. c-Jun N-terminal kinase pathways in diabetes. *Int J Biochem Cell Biol*. 2008;40:2702-6.
  59. Chang CS, Sun HL, Lii CK, Chen HW, Chen PY, Liu KL. Gamma-linolenic acid inhibits inflammatory responses by regulating NF-kappaB and AP-1 activation in lipopolysaccharide-induced RAW 264.7 macrophages. *Inflammation*. Feb;33:46-57.

60. Hess J, Angel P, Schorpp-Kistner M. AP-1 subunits: quarrel and harmony among siblings. *J Cell Sci.* 2004 Dec 1;117:5965-73.
61. Kang YJ, Mbonye UR, DeLong CJ, Wada M, Smith WL. Regulation of intracellular cyclooxygenase levels by gene transcription and protein degradation. *Prog Lipid Res.* 2007 Mar;46:108-25.
62. Kim SY, Jun TW, Lee YS, Na HK, Surh YJ, Song W. Effects of exercise on cyclooxygenase-2 expression and nuclear factor-kappaB DNA binding in human peripheral blood mononuclear cells. *Ann N Y Acad Sci.* 2009 Aug;1171:464-71.
63. Mbonye UR, Wada M, Rieke CJ, Tang HY, Dewitt DL, Smith WL. The 19-amino acid cassette of cyclooxygenase-2 mediates entry of the protein into the endoplasmic reticulum-associated degradation system. *J Biol Chem.* 2006 Nov 24;281:35770-8.
64. Nie M, Pang L, Inoue H, Knox AJ. Transcriptional regulation of cyclooxygenase 2 by bradykinin and interleukin-1beta in human airway smooth muscle cells: involvement of different promoter elements, transcription factors, and histone h4 acetylation. *Mol Cell Biol.* 2003 Dec;23:9233-44.
65. Shanmugam N, Gaw Gonzalo IT, Natarajan R. Molecular mechanisms of high glucose-induced cyclooxygenase-2 expression in monocytes. *Diabetes.* 2004 Mar;53:795-802.
66. Subbaramaiah K, Lin DT, Hart JC, Dannenberg AJ. Peroxisome proliferator-activated receptor gamma ligands suppress the transcriptional activation of cyclooxygenase-2. Evidence for involvement of activator protein-1 and CREB-binding protein/p300. *J Biol Chem.* 2001 Apr 13;276:12440-8.
67. Besnard P, Niot I, Poirier H, Clement L, Bernard A. New insights into the fatty acid-binding protein (FABP) family in the small intestine. *Mol Cell Biochem.* 2002 Oct;239:139-47.
68. Galli C. Origin of fatty acids in the body: endogenous synthesis versus dietary intakes. *Eur J Lipid Sci Technol.* 2006;108:521-5.
69. van der Vusse GJ. Albumin as fatty acid transporter. *Drug Metab Pharmacokinet.* 2009;24:300-7.
70. Gropper SS SJ, Groff JL. *Advanced Nutrition and Human Metabolism.* 5th ed. Belmont, CA: Wadsworth Cengage Learning; 2009.
71. Hajri T, Abumrad NA. Fatty acid transport across membranes: relevance to nutrition and metabolic pathology. *Annu Rev Nutr.* 2002;22:383-415.



72. Kris-Etherton PM, Taylor DS, Yu-Poth S, Huth P, Moriarty K, Fishell V, Hargrove RL, Zhao G, Etherton TD. Polyunsaturated fatty acids in the food chain in the United States. *Am J Clin Nutr.* 2000 Jan;71:179S-88S.
73. Distel RJ, Robinson GS, Spiegelman BM. Fatty acid regulation of gene expression. Transcriptional and post-transcriptional mechanisms. *J Biol Chem.* 1992 Mar 25;267:5937-41.
74. Martin H. Role of PPAR-gamma in inflammation. Prospects for therapeutic intervention by food components. *Mutat Res.* 2009 Oct 2;669:1-7.
75. Tull SP, Yates CM, Maskrey BH, O'Donnell VB, Madden J, Grimble RF, Calder PC, Nash GB, Rainger GE. Omega-3 Fatty acids and inflammation: novel interactions reveal a new step in neutrophil recruitment. *PLoS Biol.* 2009 Aug;7:e1000177.
76. Thompson BR, Mazurkiewicz-Munoz AM, Suttles J, Carter-Su C, Bernlohr DA. Interaction of adipocyte fatty acid-binding protein (AFABP) and JAK2: AFABP/aP2 as a regulator of JAK2 signaling. *J Biol Chem.* 2009 May 15;284:13473-80.
77. Coe NR, Simpson MA, Bernlohr DA. Targeted disruption of the adipocyte lipid-binding protein (aP2 protein) gene impairs fat cell lipolysis and increases cellular fatty acid levels. *J Lipid Res.* 1999 May;40:967-72.
78. Chmurzynska A. The multigene family of fatty acid-binding proteins (FABPs): function, structure and polymorphism. *J Appl Genet.* 2006;47:39-48.
79. Kazemi MR, McDonald CM, Shigenaga JK, Grunfeld C, Feingold KR. Adipocyte fatty acid-binding protein expression and lipid accumulation are increased during activation of murine macrophages by toll-like receptor agonists. *Arterioscler Thromb Vasc Biol.* 2005 Jun;25:1220-4.
80. Distel RJ, Ro HS, Rosen BS, Groves DL, Spiegelman BM. Nucleoprotein complexes that regulate gene expression in adipocyte differentiation: direct participation of c-fos. *Cell.* 1987 Jun 19;49:835-44.
81. Rauscher FJ, 3rd, Sambucetti LC, Curran T, Distel RJ, Spiegelman BM. Common DNA binding site for Fos protein complexes and transcription factor AP-1. *Cell.* 1988 Feb 12;52:471-80.
82. Watson JD BT, Bell SP, Gann A, Levine M, and Losick R. *Molecular Biology of the Gene.* 6th ed. Cold Spring Harbor, New York: Cold Spring Harbor Laboratory Press; 2008.

83. Wilson AG. Epigenetic regulation of gene expression in the inflammatory response and relevance to common diseases. *J Periodontol*. 2008 Aug;79:1514-9.
84. Vaquero A LA, and Reinberg D The Constantly Changing Face of Chromatin. *Sci Aging Knowl Environ*. 2003 April 2003;9:re4.
85. Lu SC, Wu HW, Lin YJ, Chang SF. The essential role of Oct-2 in LPS-induced expression of iNOS in RAW 264.7 macrophages and its regulation by trichostatin A. *Am J Physiol Cell Physiol*. 2009 May;296:C1133-9.
86. Park YK, Rasmussen HE, Ehler SJ, Blobaum KR, Lu F, Schlegel VL, Carr TP, Lee JY. Repression of proinflammatory gene expression by lipid extract of *Nostoc commune* var *sphaeroides* Kutzing, a blue-green alga, via inhibition of nuclear factor-kappa B in RAW 264.7 macrophages. *Nutr Res*. 2008;28:83-92.
87. Rasmussen HE, Blobaum KR, Park YK, Ehlers SJ, Lu F, Lee JY. Lipid extract of *Nostoc commune* var. *sphaeroides* Kutzing, a blue-green alga, inhibits the activation of sterol regulatory element binding proteins in HepG2 cells. *J Nutr*. 2008;138:476-81.
88. Rasmussen HE, Blobaum KR, Park YK, Ehlers SJ, Lu F, Lee JY. Lipid extract of *Nostoc commune* var. *sphaeroides* Kutzing, a blue-green alga, inhibits the activation of sterol regulatory element binding proteins in HepG2 cells. *J Nutr*. 2008 Mar;138:476-81.
89. Hotamisligil GS, Johnson RS, Distel RJ, Ellis R, Papaioannou VE, Spiegelman BM. Uncoupling of obesity from insulin resistance through a targeted mutation in aP2, the adipocyte fatty acid binding protein. *Science*. 1996 Nov 22;274:1377-9.
90. Uysal KT, Scheja L, Wiesbrock SM, Bonner-Weir S, Hotamisligil GS. Improved glucose and lipid metabolism in genetically obese mice lacking aP2. *Endocrinology*. 2000 Sep;141:3388-96.
91. Maeda K, Cao H, Kono K, Gorgun CZ, Furuhashi M, Uysal KT, Cao Q, Atsumi G, Malone H, et al. Adipocyte/macrophage fatty acid binding proteins control integrated metabolic responses in obesity and diabetes. *Cell Metab*. 2005 Feb;1:107-19.
92. Makowski L, Boord JB, Maeda K, Babaev VR, Uysal KT, Morgan MA, Parker RA, Suttles J, Fazio S, et al. Lack of macrophage fatty-acid-binding protein aP2 protects mice deficient in apolipoprotein E against atherosclerosis. *Nat Med*. 2001 Jun;7:699-705.
93. Boord JB, Maeda K, Makowski L, Babaev VR, Fazio S, Linton MF, Hotamisligil GS. Adipocyte fatty acid-binding protein, aP2, alters late atherosclerotic lesion

- formation in severe hypercholesterolemia. *Arterioscler Thromb Vasc Biol.* 2002 Oct 1;22:1686-91.
94. Layne MD, Patel A, Chen YH, Rebel VI, Carvajal IM, Pellacani A, Ith B, Zhao D, Schreiber BM, et al. Role of macrophage-expressed adipocyte fatty acid binding protein in the development of accelerated atherosclerosis in hypercholesterolemic mice. *FASEB J.* 2001 Dec;15:2733-5.
  95. Yeung DC, Xu A, Cheung CW, Wat NM, Yau MH, Fong CH, Chau MT, Lam KS. Serum adipocyte fatty acid-binding protein levels were independently associated with carotid atherosclerosis. *Arterioscler Thromb Vasc Biol.* 2007 Aug;27:1796-802.
  96. Tuncman G, Erbay E, Hom X, De Vivo I, Campos H, Rimm EB, Hotamisligil GS. A genetic variant at the fatty acid-binding protein aP2 locus reduces the risk for hypertriglyceridemia, type 2 diabetes, and cardiovascular disease. *Proc Natl Acad Sci U S A.* 2006 May 2;103:6970-5.
  97. Pelton PD, Zhou L, Demarest KT, Burris TP. PPARgamma activation induces the expression of the adipocyte fatty acid binding protein gene in human monocytes. *Biochem Biophys Res Commun.* 1999 Aug 2;261:456-8.
  98. Necela BM, Su W, Thompson EA. Toll-like receptor 4 mediates cross-talk between peroxisome proliferator-activated receptor gamma and nuclear factor-kappaB in macrophages. *Immunology.* 2008 Nov;125:344-58.
  99. Christy RJ, Yang VW, Ntambi JM, Geiman DE, Landschulz WH, Friedman AD, Nakabeppu Y, Kelly TJ, Lane MD. Differentiation-induced gene expression in 3T3-L1 preadipocytes: CCAAT/enhancer binding protein interacts with and activates the promoters of two adipocyte-specific genes. *Genes Dev.* 1989 Sep;3:1323-35.
  100. Herrera R, Ro HS, Robinson GS, Xanthopoulos KG, Spiegelman BM. A direct role for C/EBP and the AP-I-binding site in gene expression linked to adipocyte differentiation. *Mol Cell Biol.* 1989 Dec;9:5331-9.
  101. Lo YY, Cruz TF. Involvement of reactive oxygen species in cytokine and growth factor induction of c-fos expression in chondrocytes. *J Biol Chem.* 1995;270:11727-30.
  102. Haliday EM, Ramesha CS, Ringold G. TNF induces c-fos via a novel pathway requiring conversion of arachidonic acid to a lipoxygenase metabolite. *EMBO J.* 1991;10:109-15.
  103. Medzhitov R. Toll-like receptors and innate immunity. *Nat Rev Immunol.* 2001 Nov;1:135-45.

104. Guha M, Mackman N. LPS induction of gene expression in human monocytes. *Cell Signal*. 2001 Feb;13:85-94.
105. Janssen-Heininger YM, Poynter ME, Baeuerle PA. Recent advances towards understanding redox mechanisms in the activation of nuclear factor kappaB. *Free Radic Biol Med*. 2000 May 1;28:1317-27.
106. Forman HJ, Torres M. Redox signaling in macrophages. *Mol Aspects Med*. 2001 Aug-Oct;22:189-216.
107. Cakir Y, Ballinger SW. Reactive species-mediated regulation of cell signaling and the cell cycle: the role of MAPK. *Antioxid Redox Signal*. 2005 May-Jun;7:726-40.
108. Schnyder-Candrian S, Quesniaux VF, Di Padova F, Maillet I, Noulin N, Couillin I, Moser R, Erard F, Vargaftig BB, et al. Dual effects of p38 MAPK on TNF-dependent bronchoconstriction and TNF-independent neutrophil recruitment in lipopolysaccharide-induced acute respiratory distress syndrome. *J Immunol*. 2005 Jul 1;175:262-9.
109. Zhang J, Kalkum M, Chait BT, Roeder RG. The N-CoR-HDAC3 nuclear receptor corepressor complex inhibits the JNK pathway through the integral subunit GPS2. *Mol Cell*. 2002 Mar;9:611-23.
110. Lee SK, Kim JH, Lee YC, Cheong J, Lee JW. Silencing mediator of retinoic acid and thyroid hormone receptors, as a novel transcriptional corepressor molecule of activating protein-1, nuclear factor-kappaB, and serum response factor. *J Biol Chem*. 2000 Apr 28;275:12470-4.
111. Wagner BL, Valledor AF, Shao G, Daige CL, Bischoff ED, Petrowski M, Jepsen K, Baek SH, Heyman RA, et al. Promoter-Specific Roles for Liver X Receptor/Corepressor Complexes in the Regulation of ABCA1 and SREBP1 Gene Expression. *Mol Cell Biol*. 2003;23:5780-9.
112. Li J, Wang J, Nawaz Z, Liu JM, Qin J, Wong J. Both corepressor proteins SMRT and N-CoR exist in large protein complexes containing HDAC3. *EMBO J*. 2000 Aug 15;19:4342-50.
113. Guenther MG, Lane WS, Fischle W, Verdin E, Lazar MA, Shiekhata R. A core SMRT corepressor complex containing HDAC3 and TBL1, a WD40-repeat protein linked to deafness. *Genes Dev*. 2000 May 1;14:1048-57.
114. Jakobsson T, Venteclef N, Toresson G, Damdimopoulos AE, Ehrlund A, Lou X, Sanyal S, Steffensen KR, Gustafsson JA, Treuter E. GPS2 is required for cholesterol efflux by triggering histone demethylation, LXR recruitment, and coregulator assembly at the ABCG1 locus. *Mol Cell*. 2009 May 14;34:510-8.

115. Jepsen K, Hermanson O, Onami TM, Gleiberman AS, Lunyak V, McEvelly RJ, Kurokawa R, Kumar V, Liu F, et al. Combinatorial roles of the nuclear receptor corepressor in transcription and development. *Cell*. 2000 Sep 15;102:753-63.