

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

M. Eugene Rudd Publications

Research Papers in Physics and Astronomy

12-1997

Electron-Impact Total Ionization Cross Sections of CH and C₂H₂

Yong-Ki Kim

National Institute of Standards and Technology, Gaithersburg, MD

M. Asgar Ali

Howard University, Washington, DC

M. Eugene Rudd

University of Nebraska - Lincoln, erudd@unl.edu

Follow this and additional works at: <https://digitalcommons.unl.edu/physicsrudd>

 Part of the [Physics Commons](#)

Kim, Yong-Ki; Ali, M. Asgar; and Rudd, M. Eugene, "Electron-Impact Total Ionization Cross Sections of CH and C₂H₂" (1997). *M. Eugene Rudd Publications*. 18.

<https://digitalcommons.unl.edu/physicsrudd/18>

This Article is brought to you for free and open access by the Research Papers in Physics and Astronomy at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in M. Eugene Rudd Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Electron-Impact Total Ionization Cross Sections of CH and C₂H₂

Volume 102

Number 6

November–December 1997

Yong-Ki Kim

National Institute of Standards and Technology,
Gaithersburg, MD 20899-0001

M. Asgar Ali

Department of Chemistry,
Howard University,
Washington, DC 20059

and

M. Eugene Rudd

Department of Physics and
Astronomy,
University of Nebraska-Lincoln,
Lincoln, NE 68588-0111

Electron-impact total ionization cross sections for the CH radical and C₂H₂ (acetylene) have been calculated using the Binary-Encounter-Bethe (BEB) model. The BEB model combines the Mott cross section and the asymptotic form of the Bethe theory, and has been shown to generate reliable ionization cross sections for a large variety of molecules. The BEB cross sections for CH and C₂H₂ are in good agreement with the available experimental data from ionization thresholds to hundreds of eV in incident energies.

Key words: acetylene; hydrocarbon; radical; total ionization cross section.

Accepted: October 23, 1997

1. Introduction

The Binary-Encounter-Bethe (BEB) model [1] has been successfully used to calculate the total ionization cross sections of a wide range of molecules [2–4] including H₂, O₃, and SF₆. The BEB model combines a modified form of the Mott cross section with the asymptotic form of the Bethe theory (i.e., high incident energy) for electron-impact ionization of an atom.

These two theories are combined in such a way that not only the ionization cross section has the proper asymptotic form but so does the stopping cross section, which is the product of the cross section and the energy

transfer from the incident particle to the target [1]. Although the theoretical model in its differential form requires the knowledge of continuum dipole oscillator strengths df/dE , we assumed a simple form for df/dE to facilitate the integration over the ejected electron energy. The resulting expression for the ionization cross section per molecular orbital, which we refer to as the BEB model, has a remarkably simple form [1]:

$$\sigma_{\text{BEB}} = \frac{S}{t + u + 1} \left[\frac{\ln t}{2} \left(1 - \frac{1}{t^2} \right) + 1 - \frac{1}{t} - \frac{\ln t}{t + 1} \right], \quad (1)$$

where $t = T/B$, $u = U/B$, $S = 4\pi a_0^2 NR^2/B^2$, a_0 is the Bohr radius ($= 0.5292 \text{ \AA}$), R is the rydberg energy ($= 13.6057 \text{ eV}$), T is the incident electron energy, and N , B , and U are the electron occupation number, the binding energy, and the average kinetic energy of the orbital, respectively.

In this article, we compare the BEB cross sections for the CH radical in its ground state $^2\Pi$ and C_2H_2 also in the ground state $^1\Sigma_g^+$ with available experimental data. Both molecules are used in modeling the behavior of edge plasmas in the divertor region of a tokamak [5].

2. Results and Discussion

The molecular constants B and U were calculated using the 6-311+G(d,p) basis set in the GAMESS code [6]. Although one can in principle use theoretical values for all B 's, theoretical values for the lowest ionization potential (IP) are usually inaccurate. Therefore, we used experimental values [7] only for the lowest IP to match the observed thresholds. Theoretical results and comparisons to experimental data presented below are also available on a NIST web page for electron-impact ionization cross sections at:

<http://physics.nist.gov/PhysRefData/Ionization/Xsection.html>

Note that the above address is case sensitive.

2.1 CH ($^2\Pi$)

The molecular orbital constants B , U , and N for the CH radical are listed in Table 1, and the BEB model is compared in Fig. 1 to the only available experimental data by Tarnovsky et al. [8]. Although the experiment was performed using deuterium instead of hydrogen, i.e., with CD, our theory is insensitive to the use of the isotope. As Tarnovsky et al. [8] have pointed out, their experiment did not detect fast deuterons produced, which may partly account for the difference between theory and experiment near the peak. Otherwise, theory and experiment are in excellent agreement, particularly for $T < 30 \text{ eV}$. This is the region of T for which most other theories have difficulty in reproducing observed cross sections. The BEB model produces reliable cross sections for many molecules at low T ($\approx 50 \text{ eV}$ or less) [2–4].

Table 1. Molecular orbital (MO) constants for CH ($^2\Pi$): B = binding energy, U = kinetic energy, N = occupation number. The B value marked by * is an experimental value

MO	B (eV)	U (eV)	N
1σ	309.23	436.80	2
2σ	23.48	36.40	2
3σ	13.03	32.96	2
1π	10.64*	26.19	1




Fig. 1. Comparison of the BEB cross section to experiment for CH. Solid curve, BEB cross section; circles, experimental data by Tarnovsky et al. (for CD) [8]. The ordinate is the ionization cross section, and the abscissa is the incident energy.

2.2 Acetylene, C₂H₂ (¹Σ_g⁺)

The molecular orbital constants for acetylene are listed in Table 2, and the BEB model is compared in Fig. 2 to the available experimental data [9–11]. The experimental data by Zheng and Srivastava [9] and those by Tate and Smith [10] agree well with each other but they are approximately 15 % higher than the theory at the peak. On the other hand, the data by Gaudin and Hagemann [11] are in excellent agreement with the theory from $T = 100$ eV to 2 keV. Since the BEB model provides a convenient analytic form for the ionization cross section for the entire range of T shown in Fig. 2, and the data by Tate and Smith agree well with the theory for $T < 30$ eV, we are confident that the BEB cross sections will be appropriate and useful for plasma modeling applications.

Table 2. Molecular orbital (MO) constants for C₂H₂ (¹Σ_g⁺): B = binding energy, U = kinetic energy, N = occupation number. The B value marked by * is an experimental value

MO	B (eV)	U (eV)	N
1σ _g	305.62	435.15	2
1σ _u	305.50	436.31	2
2σ _g	28.18	49.60	2
2σ _u	20.80	32.79	2
3σ _g	18.55	33.64	2
1π _u	11.4*	28.99	4

Acknowledgment

The work at NIST was supported in part by the Office of Fusion Energy Science of the U.S. Department of Energy.

3. References

- [1] Y.-K. Kim and M. E. Rudd, *Phys. Rev. A* **50**, 3954 (1994).
- [2] W. Hwang, Y.-K. Kim, and M. E. Rudd, *J. Chem. Phys.* **104**, 2956 (1996).
- [3] Y.-K. Kim, W. Hwang, N. M. Weinberger, M. A. Ali, and M. E. Rudd, *J. Chem. Phys.* **106**, 1026 (1997).
- [4] M. A. Ali, Y.-K. Kim, W. Hwang, N. M. Weinberger, and M. E. Rudd, *J. Chem. Phys.* **106**, 9602 (1997).
- [5] H. Tawara, Chap. 16 in *Atomic and Molecular Processes in Fusion Edge Plasmas*, R. K. Janev, ed., Plenum Press, New York (1995).
- [6] M. W. Schmidt, K. K. Baldrige, J. A. Boatz, S. T. Elbert, M. S. Gordon, J. H. Jensen, S. Koseki, N. Matsunaga, K. A. Nguyen, S. J. Su, T. L. Windus, M. Dupuis, and J. A. Montgomery, *J. Comput. Chem.* **14**, 1347 (1993).
- [7] S. G. Lias, J. F. Liebman, R. D. Levin, and S. A. Kafafi, NIST Positive Ion Energetics Database, Version 2.0, Standard Reference Database 19A, National Institute of Standards and Technology, Oct. 1993. This database lists adiabatic IPs.
- [8] V. Tarnovsky, A. Levin, H. Deutsch, and K. Becker, *J. Phys. B* **29**, 139 (1996).
- [9] S.-H. Zheng and S. K. Srivastava, *J. Phys. B* **29**, 3235 (1996).
- [10] J. T. Tate and P. T. Smith, *Phys. Rev.* **39**, 270 (1932).
- [11] A. Gaudin and R. Hagemann, *J. Chim. Phys.* **64**, 1209 (1967).




Fig. 2. Comparison of BEB cross section to experiments for C₂H₂. Solid curve, BEB cross section; circles, experimental data by Zheng and Srivastava [9]; triangles, data by Tate and Smith [10]; diamonds, data by Gaudin and Hagemann [11]. The ordinate is the ionization cross section, and the abscissa is the incident energy.

***About the authors:** Yong-Ki Kim is a physicist in the Atomic Physics Division of the NIST Physics Laboratory. M. Asgar Ali is a professor in the Chemistry Department at Howard University. M. Eugene Rudd is a professor emeritus in the Department of Physics and Astronomy at the University of Nebraska-Lincoln. The National Institute of Standards and Technology is an agency of the Technology Administration, U.S. Department of Commerce.*