

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

2008 Bird Strike Committee USA/Canada, 10th
Annual Meeting, Orlando, Florida

Bird Strike Committee Proceedings

8-2008

Improving the DNA Identification Success Rate of Birdstrike Remains through Better Field Collecting Methods

Faridah Dahlan
Smithsonian Institution

Carla J. Dove
Smithsonian Institution

Marcy Heacker
Smithsonian Institution

Follow this and additional works at: <https://digitalcommons.unl.edu/birdstrike2008>



Part of the [Environmental Health and Protection Commons](#)

Dahlan, Faridah; Dove, Carla J.; and Heacker, Marcy, "Improving the DNA Identification Success Rate of Birdstrike Remains through Better Field Collecting Methods" (2008). *2008 Bird Strike Committee USA/Canada, 10th Annual Meeting, Orlando, Florida*. 9.
<https://digitalcommons.unl.edu/birdstrike2008/9>

This Article is brought to you for free and open access by the Bird Strike Committee Proceedings at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in 2008 Bird Strike Committee USA/Canada, 10th Annual Meeting, Orlando, Florida by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Improving the DNA Identification Success Rate of Birdstrike Remains through Better Field Collecting Methods

Faridah Dahlan, Carla J. Dove, and Marcy Heacker

Smithsonian Institution, P.O. Box 37012, Washington, DC 20013-7012
USA

DNA barcoding is one of the 'tools' used by the Smithsonian Feather Lab to identify birdstrike remains. In a study conducted during fall migration 2006, 48% of the birdstrike cases received for identification from the U.S. Air Force (USAF) and the Federal Aviation Administration (FAA) were submitted for DNA analysis. Using DNA 'barcoding' techniques, we identified nearly 68% of those cases to species-level which included 128 species of birds from 14 different Orders, and 12 bats. In the cases where we failed to obtain DNA (approx. 32%), the primary cause was due to fungi growing on the collected samples. Therefore, in 2007 we suggested using alcohol instead of water for swipes, and introduced DNA-fixing FTA® cards for collecting 'snarge' samples. These collection changes yielded more samples with minimal fungal contaminants and better success for DNA extractions (76%). In 2008, we optimized our molecular lab protocols and are now using additional primers other than CO1 ('barcoding') to achieve 85% success. Combined with the morphological feather identification tools, we have a success rate of nearly 100% in identifying birdstrike remains. This poster presents the entire process of 'snarge' sampling with emphasis on the DNA identification process and the improved collecting methods.