

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Faculty Publications from the Harold W. Manter
Laboratory of Parasitology

Parasitology, Harold W. Manter Laboratory of

1-1-2007

Research Note: *Hymenolepis horrida* (Cestoda: Hymenolepididae) and *Catenotaenia peromysci* (Cestoda: Anoplocephalidae) in Voles from the Canadian Rockies

Yeen Ten Hwang

Saskatchewan Biodiversity Assessment and Planning Unit, yeenten.hwang@gov.sk.ca

Scott Lyell Gardner

University of Nebraska - Lincoln, slg@unl.edu

John S. Millar

University of Western Ontario

Follow this and additional works at: <https://digitalcommons.unl.edu/parasitologyfacpubs>



Part of the [Parasitology Commons](#)

Hwang, Yeen Ten; Gardner, Scott Lyell; and Millar, John S., "Research Note: *Hymenolepis horrida* (Cestoda: Hymenolepididae) and *Catenotaenia peromysci* (Cestoda: Anoplocephalidae) in Voles from the Canadian Rockies" (2007). *Faculty Publications from the Harold W. Manter Laboratory of Parasitology*. 9.
<https://digitalcommons.unl.edu/parasitologyfacpubs/9>

This Article is brought to you for free and open access by the Parasitology, Harold W. Manter Laboratory of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Publications from the Harold W. Manter Laboratory of Parasitology by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Research Note

***Hymenolepis horrida* (Cestoda: Hymenolepididae) and *Catenotaenia peromysci* (Cestoda: Anoplocephalidae) in Voles from the Canadian Rockies**

YEEN TEN HWANG,^{1,3} SCOTT L. GARDNER,² AND JOHN S. MILLAR¹

¹ Department of Biology, University of Western Ontario, London, Ontario N6A 5B7, Canada, and

² Harold W. Manter Laboratory of Parasitology, W529 Nebraska Hall, University of Nebraska State Museum, and School of Biological Sciences, University of Nebraska–Lincoln, Lincoln, Nebraska 68588-0514, U.S.A.

ABSTRACT: Cestodes from *Clethrionomys gapperi* (n = 34) and *Microtus longicaudus* (n = 8) from Kootenay National Park, British Columbia, Canada are reported. Two species of cestodes, *Hymenolepis horrida* (*sensu lato*) and *Catenotaenia peromysci*, were found in *Clethrionomys gapperi*, and 1 species of cestode, *H. horrida*, was found in *Microtus longicaudus*. *Clethrionomys gapperi* represents a new host record for *C. peromysci*. The Canadian Rockies is a new locality record for both cestode species.

KEY WORDS: *Clethrionomys gapperi*, *Microtus longicaudus*, *Hymenolepis horrida*, *Catenotaenia peromysci*, red-backed vole, long-tailed vole, cestodes, boreal forest, biodiversity.

The red-backed vole (*Clethrionomys gapperi*) occurs transcontinentally through Canada, and its northwestern distribution overlaps with that of *Clethrionomys rutilus*. It also occurs in western North America from the Rocky Mountains south to southwestern Arizona and New Mexico, and in the east from the Appalachian Mountains south into northern Georgia. *Clethrionomys gapperi* is an omnivorous, opportunistic feeder inhabiting mesic environments in coniferous, deciduous, and mixed forests (Merritt, 1981). The long-tailed vole (*Microtus longicaudus*) occurs throughout most of the western United States and Canada, with its geographic range extending from New Mexico and Arizona to as far north as southeastern Alaska. *Microtus longicaudus* is primarily herbivorous and occurs in a wide variety of habitats, including coniferous and hardwood forests, meadows, and riparian areas (Smolen and Keller, 1987). Both species occur in suitable habitat in the Canadian Rockies.

Thirty-four specimens of *C. gapperi* and 8 specimens of *M. longicaudus* were collected from July 20 to August 23, 2005, at Kootenay National Park, British Columbia, Canada (50°68'N, 115°93'W). They were live-trapped with Longworth traps baited with sunflower seeds. Captured animals were trans-

ported alive to the laboratory at the Kananaskis Field Station (University of Calgary) where they were euthanized by CO₂ inhalation less than 4 hr after capture and examined for parasites. The lungs, liver, stomach, small intestine, cecum, large intestine, and bladder were examined for cestodes. Cestodes were relaxed in fresh water and fixed in 70% ethanol. For identification, cestodes were stained with Semichon's acetic-carmin solution, dehydrated in ethanol, cleared in terpineol or xylene, and mounted in Damar Gum.

Seventeen specimens of *Hymenolepis horrida* (*sensu lato*) and 2 specimens of *Catenotaenia peromysci* were found in *C. gapperi*. Fourteen specimens of *H. horrida* were found in *M. longicaudus*. Voucher specimens of parasites were deposited in the Harold W. Manter Laboratory of Parasitology (HWML), Lincoln, Nebraska.

Clethrionomys gapperi

Thirty-four red-backed voles (21 males, 13 females; mean body mass = 21.4 g) were collected from July 20 to August 23, 2005, at Kootenay National Park, British Columbia, Canada (Nearctic biogeographic region).

***Hymenolepis horrida*
(Linstow, 1901) Lühe, 1910**

Prevalence, mean intensity, and range: Hosts infected, 9 of 34 (26%, 1.9 ± 0.51; 1–5).

Site of infection: Small intestine.

Other reported hosts: *Apodemus agrarius* (Shimalov, 2002), *Arvicola terrestris* (Feliu et al., 1997), *Chionomys nivalis* (Feliu et al., 1997), *Clethrionomys bedfordiae* (Asakawa, 1993; Sakata et al., 2001), *C. gapperi* (Schiller, 1952a, b; Kinsella, 1967), *C. glareolus* (Feliu et al., 1997), *C. rutilus* (Rausch, 1952; Schiller, 1952a; Asakawa, 1993), *C. wrangeli* (Rausch,

³ Corresponding author.

1952; Schiller, 1952b), *Dicrostonyx groenlandicus* (Rausch, 1950; Kuns and Rausch, 1950; Rausch, 1952), *Eothenomys smithii* (Asakawa et al., 1992), *Lemmus amurensis* (Revin, 1983), *L. bungei* (Haukisalml and Henttonen, 2001), *L. sibiricus* (Luzhkov, 1964; Gubanov and Fedorov, 1970; Shahmatova and Yudina, 1989; Yushkov, 1995; Haukisalml and Henttonen, 2001), *L. trimucronatus* (Rausch, 1950, 1952; Schiller, 1952b; Haukisalml and Henttonen, 2001), *Meriones tristrami* (Schmidt and Wertheim, 1988), *Microtus arvalis* (Feliu et al., 1997), *Microtus chrotorrhinus* (Schiller, 1952b), *Microtus longicaudus* (Rausch, 1952; Schiller, 1952b; Kinsella 1967), *Microtus miurus* (Rausch, 1952; Schiller, 1952b; Haukisalml et al., 1995), *Microtus montanus* (Kuns and Rausch, 1950), *Microtus ochrogaster* (Rausch, 1951), *Microtus oeconomus* (Rausch, 1952; Haukisalml et al., 1995), *Microtus pennsylvanicus* (Rausch, 1952; Schiller, 1952b), *Microtus richardsoni* (Kuns and Rausch, 1950; Schiller, 1952b), *Perognathus californicus* (Voge, 1952), *Peromyscus boylei* (Voge, 1952, 1955), *P. californicus* (Voge, 1952, 1955), *P. truei* (Voge, 1952, 1955), *Pitymys* (syn. *Microtus*) *duodecimcostatus* (Feliu et al., 1997), *P. pyrenaicus* (Feliu et al., 1997), *Sciurus vulgaris* (Shimalov and Shimalov, 2002), *Tamiasciurus hudsonicus* (Schiller, 1952b), *Thomomys bottae* (Voge, 1952), *T. bulbivorus* (Gardner, 1985), *T. monticola* (Howard and Childs, 1959), *T. talpoides* (Frandsen and Grundmann, 1961), *T. umbrinus* (Frandsen and Grundmann, 1961), and *Zapus princeps* (Kinsella, 1968).

Geographic range: Asia: Japan (Asakawa et al. 1992; Asakawa, 1993; Sakata et al., 2001); Africa: Sinai (Egypt—Schmidt and Wertheim, 1988); Europe: Belarus (Shimalov, 2002; Shimalov and Shimalov, 2002), Russia (Luzhkov, 1964; Gubanov and Fedorov, 1970; Revin, 1983; Shahmatova and Yudina, 1989; Yushkov, 1995; Haukisalml and Henttonen, 2001), Spain (Feliu et al., 1997); North America: Alaska (U.S.A.—Rausch, 1950, 1952; Schiller, 1952b; Haukisalml et al., 1995), California (U.S.A.—Voge, 1952, 1954; Howard and Childs, 1959), Montana (U.S.A.—Schiller, 1952b; Kinsella, 1968), Oregon (U.S.A.—Gardner, 1985), Quebec (Canada—Schiller, 1952b), Tennessee (U.S.A.—Rausch, 1951), Utah (U.S.A.—Frandsen and Grundmann, 1961), Washington (U.S.A.—Schiller, 1952a), Wyoming (U.S.A.—Kuns and Rausch, 1950).

Specimens deposited: HWML# 48382, 48387, 48388, 48389, 48390, 48391, 48392, 48393.

Remarks: *Hymenolepis horrida* (*sensu lato*) is a

widespread parasite of arvicoline rodents (i.e., voles and lemmings) in the northern hemisphere, with a Holarctic distribution (Doran, 1954; Rausch, 1957). Recent DNA analyses have concluded that *H. horrida* may not be a single species but may consist of multiple species (cf. the “*Arostrilepis horrida* complex”) with similar morphologies (Hoberg et al., 2003). Kootenay National Park, British Columbia, Canada, is a new locality record for the species in Canada.

Catenotaenia peromysci

Prevalence, mean intensity, and range: Hosts infected, 1 of 34 (3%, 2).

Site of infection: Small intestine.

Type host and type locality: *Peromyscus maniculatus rufinus* (Merriam), Rio Arriba Country, New Mexico (Smith, 1953).

Other reported hosts: *Peromyscus maniculatus sonoriensis* (Grundmann et al., 1976).

Geographic range: Colorado (U.S.A.—Smith, 1953), New Mexico (U.S.A.—Smith, 1953), Utah (U.S.A.—Grundmann et al., 1976), Wyoming (U.S.A.—Smith, 1953).

Specimens deposited: HWML# 48385.

Remarks: *Catenotaenia peromysci* appears to be closely related to *C. pusilla*. However, in general, many structures of *C. peromysci* are larger than those of *C. pusilla* (Smith, 1953). *Clethrionomys gapperi* is a new host record for *C. peromysci*. Rausch and Tiner (1948) reported *C. pusilla* infecting *C. gapperi*, but this was later corrected as *Catenotaenia dendritica* (Doran, 1954). *Catenotaenia dendritica* has been found rarely in *C. gapperi* (Kinsella, 1967). Our specimens do not conform to descriptions of *C. dendritica* and fit most closely the descriptions of *C. peromysci* (length: 62 to 67 mm; width: 0.82 mm; scolex: 0.28 mm; number of testes: 65; cirrus sac length to width: 0.15 to 0.07; primary uterine branches: 24). *Clethrionomys gapperi* represents a new host record for this species.

Microtus longicaudus

Eight long-tailed voles (6 males, 2 females; mean body mass = 27.5 g) were collected from July 20 to August 23, 2005, at Kootenay National Park, British Columbia, Canada (Nearctic biogeographic region).

Hymenolepis horrida
(Linstow, 1901) Lühe, 1910

Prevalence, mean intensity, and range: Hosts infected, 4 of 8 (50%, 4.7 ± 1.2 ; 3–7).

Site of infection: Small intestine

Specimens deposited: HWML# 48381, 48383, 48384, 48386.

Remarks: Taxonomic summary, distributional summary, and other remarks are presented under *C. gapperi*.

We report the cestode community found in voles, *Clethrionomys gapperi* and *Microtus longicaudus*, in the Kootenay National Park of the Canadian Rockies. *Hymenolepis horrida* has a wide geographic distribution and broad host range (Rausch, 1957). Schiller (1952b) noted exceptional morphological variation and diversity in *H. horrida* infecting various arvicolid rodents in the Nearctic region. It was established as the type species for the genus *Arostrilepis* based on morphological evidence (Mas-Coma and Tenora, 1997). However, the type species was found in the Common Pine Vole, *Microtus subterraneus*, from Hungary (Mas-Coma and Tenora, 1997). However, recent molecular studies on *H. horrida* from different parts of its distribution have shown that it perhaps encompasses multiple species in the *Arostrilepis* complex (Hoberg et al., 2003). Therefore, we retain the name *H. horrida* until taxonomic revision of the species in the “complex” in both New and Old World rodents is published.

The prevalence and intensity of *H. horrida* were higher in *M. longicaudus* than in *C. gapperi*. Perhaps individuals of *M. longicaudus* have a greater probability of coming into contact with infected arthropod intermediate hosts than do individuals of *C. gapperi*. Another possibility is that *C. gapperi* may have greater resistance to infection by *H. horrida*. We examined other sympatric small mammals for the presence of parasites, including *Phenacomys intermedius*, *Peromyscus maniculatus*, *Tamias amoenus*, and *Spermophilus lateralis*, and found no *H. horrida*. Thus, both red-backed and long-tailed voles probably have at least a partial niche overlap in this part of the Rockies (Millar et al., 1985).

Although both species of mammals share the same *Hymenolepis* species, only *C. gapperi* was infected with *C. peromysci*. The significance of the observation that *C. peromysci* infected only *C. gapperi* cannot be determined in this study.

Many thanks to S. Gil for field collection of parasites, Terry Haverkost and David Scott Tinnin for assistance in parasite identification in the HWML, and Kootenay National Park staff for allowing us to work within the park. Funding was provided by the G8 Legacy Fund to YTH and the Natural Science and Engineering Research Council to JSM. Logistic support for identification of parasites was provided by the staff and facilities of the Harold W. Manter Laboratory of Parasitology, University of Nebraska State Museum, University of Nebraska, Lincoln.

LITERATURE CITED

- Asakawa, M.** 1993. Parasitic helminths obtained from rodents on Nemuro Peninsula and Notsuke-saki, Hokkaido, Japan. *Kokuritsu-Kahaku-Senpou* 26:75–82.
- Asakawa, M., F. Tenora, S. Fukumoto, K. Kano, and T. Tomonari.** 1992. Faunal and zoogeographical study on the parasitic helminths of voles and field mice in Shikoku, Japan. *Bulletin of Tokushima Prefectural Museum* 2:51–75.
- Doran, D. J.** 1954. A catalogue of the protozoa and helminths of North American rodents. II. Cestoda. *American Midland Naturalist* 52:469–480.
- Feliu, C., F. Renaud, F. Catzeflis, J.-P. Hugot, P. Durand, and S. Morand.** 1997. A comparative analysis of parasite species richness of Iberian rodents. *Parasitology* 115:453–466.
- Frandsen, J. C., and A. W. Grundmann.** 1961. Endoparasitism in isolated populations of rodents of the Lake Bonneville basin, Utah. *Journal of Parasitology* 47: 391–396.
- Gardner, S. L.** 1985. Helminth parasites of *Thomomys bulbivorus* (Richardson) (Rodentia: Geomyidae), with the description of a new species of *Hymenolepis* (Cestoda). *Canadian Journal of Zoology* 63:1463–1469.
- Grundmann, A. W., R. G. Warnock, and D. L. Wassom.** 1976. Some mechanisms of natural regulation of parasitic helminth populations. *American Midland Naturalist* 95:347–360.
- Gubanov, N. M., and K. P. Fedorov.** 1970. [Helminth fauna of Muridae of Yakuti]. Pages 18–47 in *Fauna Sibirii*. Novosibirsk: Nauka. (In Russian.)
- Haukisalmi, V., and H. Henttonen.** 2001. Biogeography of helminth parasitism in *Lemmus* Link (Arvicolinae), with the description of *Paranoplocephala fellmani* n. sp. (Cestoda: Anoplocephalidae) from Norwegian lemming *L. lemmus* (Linnaeus). *Systematic Parasitology* 49:7–22.
- Haukisalmi, V., H. Henttonen, and G. O. Batzli.** 1995. Helminth parasitism in the voles *Microtus oeconomus* and *M. miurus* on the North Slope of Alaska: host specificity and the effects of host sex, age and breeding status. *Annales Zoologici Fennici* 32:193–201.
- Hoberg, E. P., S. J. Kutz, K. E. Galbreath, and J. A. Cook.** 2003. Arctic biodiversity: from discovery to faunal baselines—revealing the history of a dynamic

- ecosystem. *Journal of Parasitology* 89 (Supplement): S84–S95.
- Howard, W. E., and H. E. Childs.** 1959. Ecology of pocket gophers with emphasis on *Thomomys bottae mewa*. *Hilgardia* 29:277–358.
- Kinsella, J. M.** 1967. Helminths of Microtinae in Western Montana. *Canadian Journal of Zoology* 45:269–274.
- Kinsella, J. M.** 1968. Helminths of the Western Jumping Mouse (*Zapus princeps*) in Western Montana. *Journal of Parasitology* 54:309.
- Kuns, M. L., and R. Rausch.** 1950. An ecological study of helminths of some Wyoming voles (*Microtus* spp.) with a description of a new species of *Nematospiroides* (Heligmosomidae: Nematoda). *Zoologica* 35:181–188.
- Luzhkov, A. D.** 1964. [Parasitic worms of lemmings and voles of the Yamal peninsula]. Materialy k Nauchnoj Konferenicii Vsesojuznogo Obschestva Gel'mintologov 1:234–238. (In Russian.)
- Mas-Coma, S., and F. Tenora.** 1997. Proposal of *Arostreplepis* n. gen. (Cestoda: Hymenolepididae). *Research and Reviews in Parasitology* 57:93–101.
- Merritt, J. F.** 1981. *Clethrionomys gapperi*. *Mammalian Species* 146:1–9.
- Millar, J. S., D. G. L. Innes, and V. A. Loewen.** 1985. Habitat use by non-hibernating small mammals of the Kananaskis Valley, Alberta. *Canadian Field-Naturalist* 99:196–204.
- Rausch, R.** 1950. Observations on the cyclic decline of Lemmings (*Lemmus*) on the Arctic coast of Alaska during the spring of 1949. *Arctic* 3:166–177.
- Rausch, R.** 1951. Biotic interrelationships of helminth parasitism. *Public Health Reports* 66:928–934.
- Rausch, R.** 1952. Studies on the helminth fauna of Alaska. XI. Helminth parasites of microtine rodents—taxonomic considerations. *Journal of Parasitology* 45:415–444.
- Rausch, R.** 1957. Distribution and specificity of helminths in microtine rodents: evolutionary implications. *Evolution* 11:361–368.
- Rausch, R., and J. D. Tiner.** 1948. Studies on the parasitic helminths of the North Central States. I. Helminths of Sciuridae. *American Midland Naturalist* 39:728–747.
- Revin, Yu V.** 1983. [Materials on the ecology of the Amur lemming (*Lemmus amurensis*) in South Yakutia.] *Zoologicheskii Zhurnal* 62:922–929. (In Russian.)
- Sakata, K., I. A. Nevedomskaya, N. Kondo, and M. Asakawa.** 2001. Parasitic helminth fauna of wild rodents on Uenkotan, Kunashiri I. The Memoirs of the Preparative Office of Nemuro Municipal Museum 15:115–118.
- Schiller, E. L.** 1952a. Studies on the helminth fauna of Alaska. V. Notes on Adak rats (*R. norvegicus* Berkenhout) with special reference to helminth parasites. *Journal of Mammalogy* 33:38–49.
- Schiller, E. L.** 1952b. Studies on the helminth fauna of Alaska. X. Morphological variation in *Hymenolepis horrida* (von Linstow, 1901) (Cestoda: Hymenolepididae). *Journal of Parasitology* 36:554–568.
- Schmidt, G. D., and G. Wertheim.** 1988. *Inermicapsifer beveridgei* n. sp. (Cestoidea: Anoplocephalidae) from *Procapra capensis* (Hyracoidea) in Israel, with notes on two species of *Hymenolepis*. *Journal of Parasitology* 74:487–488.
- Shahmatova, V. I., and S. A. Yudina.** 1989. [Helminths of Muridae of Tajmyra.] Pages 145–178 in K. P. Fedorov, ed. *Ekologiya gel'mintov pozvonochnykh Sibiri*. Novosibirsk: Nauka. (In Russian.)
- Shimalov, V. V.** 2002. Helminth fauna of the striped field mouse (*Apodemus agrarius* Pallas, 1778) in ecosystems of Belorussian Polesie transformed as a result of reclamation. *Parasitology Research* 88:1009–1010.
- Shimalov, V. V., and V. T. Shimalov.** 2002. Helminth fauna of the red squirrel (*Sciurus vulgaris* Linnaeus, 1758) in Belorussian Polesie. *Parasitology Research* 88:1008.
- Smith, C. F.** 1954. Four new species of cestodes of rodents from the high plains, central, and southern rockies and notes on *Catenotaenia dendritica*. *Journal of Parasitology* 40:245–254.
- Smolen, M. J., and B. L. Keller.** 1987. *Microtis longicaudus*. *Mammalian Species* 271:1–7.
- Voge, M.** 1952. Variation in some unarmed Hymenolepididae (Cestoda) from rodents. University of California Publications in Zoology 57:1–52.
- Voge, M.** 1955. A list of cestode parasites from California mammals. *American Midland Naturalist* 54:413–417.
- Yushkov, V. F.** 1995. [Fauna of the European North-East of Russia.] Page 225 in *Gel'minty mlekopitayuschih*. St. Petersburg: Rossijskaya Akademiya Nauk. (In Russian.)