

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

---

5 - Fifth Eastern Wildlife Damage Control  
Conference (1991)

Eastern Wildlife Damage Control Conferences

---

February 1991

## MODIFICATION OF RACCOON FOOD RESOURCES IN PARKS TO REDUCE RABIES RISKS TO HUMANS

Stephen C. Frantz

*Wadsworth Center for Laboratories and Research, New York State Department of Health, Albany, NY*

Follow this and additional works at: <https://digitalcommons.unl.edu/ewdcc5>



Part of the [Environmental Health and Protection Commons](#)

---

Frantz, Stephen C., "MODIFICATION OF RACCOON FOOD RESOURCES IN PARKS TO REDUCE RABIES RISKS TO HUMANS" (1991). *5 - Fifth Eastern Wildlife Damage Control Conference (1991)*. 14.  
<https://digitalcommons.unl.edu/ewdcc5/14>

This Article is brought to you for free and open access by the Eastern Wildlife Damage Control Conferences at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in 5 - Fifth Eastern Wildlife Damage Control Conference (1991) by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

# MODIFICATION OF RACCOON FOOD RESOURCES IN PARKS TO REDUCE RABIES RISKS TO HUMANS

STEPHEN C. FRANTZ, Wadsworth Center for Laboratories and Research, New York State Department of Health, Albany, NY 12201-0509

Proc. East. **Wildl. Damage Control Conf.** 5:165. 1992.

In May 1990, the mid-Atlantic raccoon (*Procyon lotor*) rabies epizootic crossed New York State's southern border. By the end of the year, 84 rabid raccoons had been identified from a 400-km "front" in the State's southern tier of counties. By about mid-February 1991, the number of confirmed rabid raccoons increased to 134. Major metropolitan areas are threatened and predictions for the future indicate a pressing need for information and services by various government agencies. High-risk areas currently are park campgrounds, but will be extended as the outbreak progresses to include urban parks and those residential communities having large raccoon populations. Well designed, comprehensive management strategies are needed to reduce human and domestic animal contacts with raccoons during the outbreak. Although many interventions are available, the purpose of this study is to develop methods that will lessen raccoon activity in and around park areas with intense patron use and large raccoon popula-

tions. The objective is to reduce raccoon food resources through improved solid-waste storage/removal, and to educate park patrons not to feed raccoons directly or indirectly. Observation and recording of activity patterns and behavior of raccoons and/or humans at study sites is conducted primarily by remote-controlled, low-light videography. We learned that some raccoons easily climb, open lids, and enter some designs of garbage containers, where they may remain for several hours. Raccoons and other wildlife (e.g., skunks [*Mephitis mephitis*]) also feed upon spilled garbage and drainage from the garbage containers. Wildlife feeding activity at garbage containers has been observed to partially overlap with the daily activity period of park patrons. We are continuing to investigate the problems and remedial interventions of garbage container design, raccoon and human behavior, and implications for rabies risk reduction.