University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Uniformed Services University of the Health
Sciences

U.S. Department of Defense

2003

Mutations in hns reduce the adherence of Shiga toxin-producing E. coli 091:H21 strain B2F1 to human colonic epithelial cells and increase the production of hemolysin

Maria E. Scott

American Red Cross, Rockville, MD

Angela R. Melton-Celsa

Uniformed Services University of the Health Sciences, angela.melton-celsa@usuhs.edu

Alison D. O'Brien

Uniformed Services University of the Health Sciences, alison.obrien@usuhs.edu

Follow this and additional works at: <https://digitalcommons.unl.edu/usuhs>

Part of the [Medicine and Health Sciences Commons](#)

Scott, Maria E.; Melton-Celsa, Angela R.; and O'Brien, Alison D., "Mutations in hns reduce the adherence of Shiga toxin-producing E. coli 091:H21 strain B2F1 to human colonic epithelial cells and increase the production of hemolysin" (2003). *Uniformed Services University of the Health Sciences*. 14.
<https://digitalcommons.unl.edu/usuhs/14>

This Article is brought to you for free and open access by the U.S. Department of Defense at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Uniformed Services University of the Health Sciences by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Short communication

Mutations in *hns* reduce the adherence of Shiga toxin-producing *E. coli* 091:H21 strain B2F1 to human colonic epithelial cells and increase the production of hemolysin

Maria E. Scott^a, Angela R. Melton-Celsa^b, Alison D. O'Brien^{b,*}

^aDepartment of Biochemistry, American Red Cross, Rockville, MD 20855, USA

^bDepartment of Microbiology and Immunology, F. Edward Hébert School of Medicine, Uniformed Services University of the Health Sciences, 4301 Jones Bridge Road, Bethesda, MD 20814-4799, USA

Received 10 October 2002; received in revised form 9 December 2002; accepted 9 December 2002

Abstract

Shiga toxin-producing *Escherichia coli* (STEC) 091:H21 strain B2F1, an isolate from a patient with the hemolytic uremic syndrome (HUS), produces elastase-activatable Shiga toxin (Stx) type 2d and adheres well to human colonic epithelial T84 cells. This adherence phenotype occurs even though B2F1 does not contain the locus of enterocyte effacement (LEE) that encodes the primary adhesin for *E. coli* O157:H7. To attempt to identify genes involved in binding of B2F1 to T84 cells a bank of mini-Tn5 ϕ ACm^r transposon mutants of this strain was generated. Several of these mutants exhibited a reduced adherence phenotype, but none of the insertions in these mutants were within putative adhesin genes. Rather, insertional mutations within *hns* resulted in the loss of adherence. Moreover, the *hns* mutant also displayed an increase in the production of hemolysin and alkaline phosphatase and a loss of motility with no change in Stx2d-activatable expression levels. When B2F1 was cured of the large plasmid that encodes the hemolysin, the resulting strain adhered well to T84 cells. However, an *hns* mutant of the plasmid-cured B2F1 strain exhibited a reduction in adherence to T84 cells. Taken together, these results indicate that H-NS regulates the expression of several genes and some potential virulence factors in the intimin-negative B2F1 STEC strain and that the large plasmid is not required for T84 cell colonization.

Published by Elsevier Science Ltd.

Keywords: B2F1; Adherence; Hemolysin; Non-O157

1. Introduction

Strains of non-O157 STEC have been isolated from patients with hemorrhagic colitis (HC). Some of these isolates have been associated with the hemolytic uremic syndrome (HUS) in infected persons [1–4]. Four clonally-related groups of STEC have been identified and are designated EHEC 1, EHEC 2, STEC 1, and STEC 2 [5,6]. Members of EHEC groups 1 and 2 encode the locus of enterocyte effacement (LEE) and produce intimin, an adherence factor common to both EHEC groups [6,7]. The O157:H7 strains that cause most of the major outbreaks of STEC-related illness typically belong to the EHEC 1 group. STEC 1 and 2 members typically do not encode the gene for

intimin nor do they carry the LEE pathogenicity island [6]. Serotype 091:H21 strain B2F1 [8] lacks the LEE locus, as determined by Southern blot with LEE-locus probes [9], and is categorized as an STEC 1 group member. B2F1 was originally isolated from a patient with HUS (personal communication, Dr M. Karmali). We hypothesized that B2F1, like other clinical isolates of intimin-negative STEC [10–12], has the capacity to adhere to epithelial cells by an intimin-independent mechanism. Therefore, we attempted to define factors important for the adherence of B2F1 to human intestinal epithelial cells. Although an intestinal cell adhesion for the intimin-negative 091:H21 STEC strain was not identified, three new findings were generated from this study. First, we found that the H-NS protein regulates, in a positive manner, the adherence of B2F1 to T84 human colonic epithelial cells, as well as the motility of B2F1. Second, we demonstrated that H-NS negatively regulates the plasmid-encoded hemolysin and chromosomally

* Corresponding author. Tel.: +1-301-295-3400; Fax: +1-301-295-3773.

E-mail address: aobrien@usuhs.mil (A.D. O'Brien).

encoded alkaline phosphatase produced by B2F1. Third, a plasmid-cured strain of B2F1 was used to show that the large plasmid is not required for B2F1 to adhere to T84 cells.

2. Results and discussion

2.1. Rationale for selection of bacterial strain and eukaryotic cells

Strain B2F1, a human 091:H21 isolate that has two copies of the Stx2d-activatable [8,13] adhered to T84 cells at consistently high levels (Fig. 1A). The laboratory *E. coli* control strain DH5 α adhered poorly to T84 cells (not shown). T84 cells were used as the model cell line for adherence for two reasons. First, T84 cells are not sensitive to killing by Stx because they lack the toxin binding receptor [14]. Second, under the appropriate in vitro conditions the T84 human colonic adenocarcinoma cell line grows in a polarized manner and resembles the native intestinal colonic crypt cell from which it was derived [15]. We found that B2F1 adhered with a similar phenotype (large irregularly shaped colonies) to both polarized and non-polarized T84 cells (not shown). Therefore, we elected to use non-polarized T84 cells for most of this study for ease of maintenance and growth but confirmed all adherence results with polarized T84 cells.

2.2. Isolation of mini-Tn5phoA *Cm*^r adherence mutants

B2F1 was mutagenized with a mini-Tn5phoA*Cm*^r transposon to screen for mutants with insertions in potential adhesins. The mini-Tn5phoA*Cm*^r used in this study was generated by replacing the kanamycin resistance gene in

mini-Tn5phoA [16] with a chloramphenicol resistance gene (not shown). Chloramphenicol was used as the selective marker in the mutagenesis rather than kanamycin because B2F1 mutates to kanamycin resistance at a high frequency. We isolated 22,555 chloramphenicol resistant transconjugates of B2F1. Approximately 1% of the chloramphenicol resistant transconjugates were blue when grown on Luria–Bertani (LB) agar with the phosphatase substrate 5-bromo-4-chloro-3-indolyl phosphate (XP). The blue mutants that contained putative active alkaline phosphatase fusions to exported proteins were screened for the capacity to bind to confluent T84 cells. Briefly, wild type and mutant strains were allowed to associate with T84 cells in plastic chamber slides for a total of 3 h at 37 °C in 5% CO₂. Samples were washed 10 times with fresh Dulbecco's PBS supplemented with Ca²⁺/Mg²⁺ fixed and stained with Leukostat (modified Wright's stain from Fisher Scientific, New Hampshire), and screened qualitatively by phase contrast microscopy (assay modified from) [17,18]. Quantitative analysis was done in a similar manner except trypsin was added to the 8-well chamber slides after the 3 h incubation period. The bacterial-eukaryotic cell suspension was then serially diluted and plated on LB agar to obtain viable colony counts. Each sample was done in triplicate. The geometric mean adherent CFU was calculated for each strain tested. Of the 253 mutants screened, two, 34.3 and 34.7, had a reduced adherence phenotype (Fig. 1B shows 34.7). Quantitative analysis of a 3 h adherence assay on polarized T84 cells revealed that approximately 10⁵ CFU/ml of strains 34.3 or 34.7 were recovered from each well of the chamber slide as compared to 10⁷ CFU/ml of wild-type B2F1.

Southern hybridization analysis of *Hind*III-restricted genomic DNA from the mutants (34.7 and 34.3) with reduced adherence showed that each mutant appeared to contain a single insertion in an approximately 6 kb DNA fragment (data not shown). Furthermore, double-restriction enzyme Southern blot analysis of the two mutants revealed that the insertion was in the same DNA fragment (data not shown). Therefore, only 34.7 was analyzed further.

2.3. Sequence analysis of the mini-Tn5phoA*Cm*^r disrupted gene of mutant 34.7

The 6 kb *Hind*III DNA fragment that contained the mini-Tn5phoA*Cm*^r from mutant 34.7 was cloned by classical methods [19] into pBR322 and sequenced. Analysis of that sequence revealed that the transposon had inserted near the start of the B2F1 *hns* gene. However, the sequence analysis also showed that *hns* and *phoA* were not in the same reading frame, a finding that indicated that neither a functional H-NS nor a functional alkaline phosphatase fusion protein could be produced in 34.7. Therefore, we concluded that H-NS regulates the expression of the native alkaline phosphatase in B2F1, since 34.7 produced blue colonies on media with XP. Other investigators have reported similar findings when H-NS was inactivated [20].

Fig. 1. Phase contrast microscopy shows the reduced capacity of strains with mutations in *hns* to adhere to T84 cells as compared to wild-type. The parental wild-type strain B2F1 is shown in panel A and its isogenic plasmid-cured derivative, S11, is shown in panel C; both strains exhibit a wild-type adherence phenotype on T84 cells. In contrast, *hns* mutants 34.7 (panel B) and S11358.19 (panel D) are deficient in adherence. Arrows indicate clusters of adherent bacteria.

2.4. Phenotypic analysis and complementation of mutant 34.7 with native H-NS

Inactivation of H-NS affects the expression of several unlinked and unrelated genes encoded in *E. coli* and *Salmonella typhimurium* [21,22]. Therefore, several characteristics of 34.7 that are potentially relevant to STEC virulence were examined. Mutant 34.7 grew at rates similar to B2F1 and expressed Stx2d-activatable at wild-type levels (not shown), but unlike the parental B2F1 strain, mutant 34.7 was nonmotile (Table 1). However, the non-motile phenotype of 34.7 was most likely not responsible for the failure of 34.7 to adhere to T84 cells, since other mini-Tn5*phoAC*^r non-motile mutants did adhere to T84 cells (not shown). Strain 34.7 also produced large zones of hemolysis on agar that contained 5% washed sheep red blood cells (SRBCs) in contrast to the small turbid zones of hemolysis produced by wild-type B2F1, Table 1. The hemolysin operon is located on the large ~90 kb plasmid carried by the vast majority of EHEC and STEC clinical isolates [23]. However, hemolytic activity is generally not detectable in the cell-free filtrates of wild-type STEC [23], an observation that we confirmed with B2F1 (not shown). In contrast, hemolytic activity was detectable in culture supernatants of 34.7 and was quantifiable by a hemolytic tube assay [24], Table 1. Furthermore, the amount of hemolysin in bacterial culture supernatants and whole cell lysates of 34.7 was greater than that produced by the parent (Table 1), as determined by Western blot analysis with antibody raised against α -hemolysin [24]. Thus, these Western blot data taken together with the hemolytic phenotype of 34.7 noted on SRBC agar and in tube assays indicate that hemolysin production is deregulated in 34.7. Similarly, H-NS in complex with Hha, a protein that belongs to a relatively new family of modulators, controls the expression of the thermoregulated α -hemolysin operon encoded by uropathogenic *Escherichia coli* [25].

The adherence capacity of mutant 34.7 was restored to wild-type levels when *hns* was introduced into 34.7 on a moderate copy number plasmid while transformation with the vector alone failed to complement the adherence defect of 34.7 (data not shown). This finding confirmed that the *hns* mutation in 34.7 was responsible for the defect in adherence. Similarly, the wild-type hemolytic phenotype was restored in mutant 34.7 complemented with wild-type *hns*, as demonstrated by small turbid zones of hemolysis around the colonies on SRBC agar (not shown, see Table 1 for a summary of phenotypic data).

2.5. Construction of a *hns* insertion mutation in a plasmid-cured derivative of B2F1 and assessment of the capacity of that mutant to adhere to T84 cells

We next sought to separate the reduced adherence phenotype of 34.7 from the hyper-hemolytic phenotype exhibited in that mutant by constructing a *hns* mutation in a B2F1 derivative lacking hemolysin. First, a hemolysin negative derivative of B2F1 was generated by curing the large plasmid from a mutant of B2F1 that carried a mini-Tn5*phoAC*^r insertion on the large plasmid. The mutant was cured by repeated passaging of the mutant in LB containing ampicillin/cycloserine. Of 11 chloramphenicol sensitive derivatives of the mutant, one, strain S11, was found not to hybridize to the probe for the large plasmid [26] and did not have a visible large plasmid present in plasmid preparations (not shown). As expected, S11 still adhered to T84 cells with a wild-type phenotype (Fig. 1C) and was not hemolytic on SRBC agar (Table 1). Second, an *hns* mutation was introduced into the non-hemolytic S11 by integration of a mobilizable suicide plasmid that contained a portion of the *hns* coding sequence into the wild-type *hns* in S11. (Note: repeated attempts to transfer the null mutation from 34.7 to S11 by allelic exchange or P1 transduction failed.)

Table 1
Genotypic and phenotypic characteristics of B2F1, S11 and their isogenic *hns* mutants

Strain	Relevant genotype	Phenotype				
		Alkaline phosphatase ^a	Adherence ^b	Hemolysis (titer) ^c	Hemolysin ^d	Motility ^e
B2F1	Wt	+/+	+	-/+(0)	-	+++
34.7	B2F1 <i>hns</i> :: mini-Tn5 <i>phoAC</i> ^r	++	-	++(8)	+	-
34.7 (pHNS)	B2F1 <i>hns</i> :: mini-Tn5 <i>phoAC</i> ^r (pHNS)	-	+	-/+(ND) ^f	ND	+
S11	B2F1 (large plasmid cured)	-/+	+	-(ND)	-	++
S11358.19	S11 <i>hns</i> :: pMS358	+	-	ND (ND)	ND	+
S11358.19 (pHNS)	S11 <i>hns</i> :: pMS358 (pHNS)	-	+	ND (ND)	ND	+

^a Alkaline phosphatase was determined as blue colony color after growth on LB plates with XP.

^b Adherence was measured on T84 cells as described in the text.

^c Hemolysis was detected as a clearing of red blood cells around colonies grown on tryptose blood agar with 10 mM CaCl₂ and 5% SRBCs. The hemolytic titer was determined as the reciprocal dilution where lysis of sheep erythrocytes was evident after 2 h at 37 °C.

^d Hemolysin was detected by Western blot with antibody to α -hemolysin [24].

^e Motility was assayed after growth in semi-solid agar plates at 30 °C overnight.

^f ND-not determined.

Successful insertional inactivation of a gene by this method has been previously reported [27].

The construction of the *hns* mutation in S11 is described briefly as follows. First, a 358 bp portion of *hns* from S11 was amplified by the polymerase chain reaction (PCR). Second, the truncated *hns* PCR product was ligated into a temperature-sensitive suicide plasmid, pSTAMP [28], to generate plasmid pMS358. Third, plasmid pMS358 was electroporated into S11 and transformants that were ampicillin resistant were recovered at 30 °C. Fourth, stable colonies of S11 (pMS358) were incubated overnight at 44 °C on agar containing ampicillin to select for colonies with pMS358 integrated into *hns* encoded on the chromosome of S11. Putative cointegrates were analyzed by PCR and Southern blot analysis to confirm that integration of the plasmid into *hns* had occurred (not shown). Insertion of the plasmid into wild-type *hns* generated two incomplete copies of *hns* only one of which retained an intact promoter region. The H-NS produced from the copy of *hns* with an intact promoter was predicted to lack six amino acid residues at its carboxy-terminal end.

The in vitro growth rate of the S11 *hns* mutant, S11358.19, was similar to that of wild-type S11 (not shown). However, as predicted, S11358.19 showed reduced adherence to T84 cells compared to S11 (Fig. 1D). Furthermore, a quantitative adherence assay demonstrated that 10⁵ CFU/ml of S11358.19 were recovered as compared to 10⁶ CFU/ml of its parental strain S11 and B2F1. (The reason for the lower overall adherence of the wild-type B2F1 in this set of assays as compared to the assays described above is unknown, however, the adherence of B2F1 and S11 was consistently 10-fold higher than that of the *hns* mutant of S11.) These results demonstrate that the absence of H-NS function, not the presence of increased STEC hemolysin, was responsible for the reduction in the adherence of mutant 34.7 to T84 cells. Mutant S11358.19 was also blue on XP agar and exhibited reduced motility, whereas the *hns* mutant of B2F1 was non-motile, Table 1. This result suggested that the activity of H-NS in mutant S11358.19 was not completely ablated. When S11358.19 was transformed with a low copy number plasmid that encoded full-length H-NS protein, the adherence and alkaline phosphatase phenotypes were restored to wild-type levels (not shown). However, full motility was not restored to S11358.19 (pHNS), see Table 1. One explanation for this finding is that the truncated H-NS polypeptide produced by the mutant S11358.19 interacted with wild-type H-NS supplied in trans, which, in turn, resulted in a decrease in H-NS activity. Such a dominant negative phenotype is plausible since H-NS is thought to act as an oligomer or homodimer [22,29].

In summary, *hns* mutations in B2F1 or its plasmid-cured derivative S11 reduced the capacity of those mutants to adhere to T84 cells and resulted in a reduction in motility and an elevation in alkaline phosphatase activity. In B2F1 the *hns* mutation also increased the plasmid-encoded

hemolytic activity. Regulation of STEC-hemolysin expression in STEC or O157:H7 has not been previously described. Furthermore, genes encoded by the large plasmid have not been reported to be controlled by chromosomal gene products in STEC. In the enteropathogenic *E. coli* (EPEC) regulation of attachment and effacement genes by Ler, which is homologous to H-NS of *Salmonella*, has been reported [30]. Bustamante and colleagues showed that Ler activates transcription of LEE genes by competing for binding with H-NS and that in the absence of Ler H-NS represses the expression of LEE genes important in AE lesion formation [31]. For O157:H7 EHEC strains, *hns* has been suggested to be a repressor of the *espA* operon [32], the products of which are involved in early steps of adherence. Although Ler is also present in *E. coli* O157:H7, intimin-negative STEC do not encode either the LEE pathogenicity island or the Ler gene. Furthermore, our report suggests that H-NS may play a somewhat different role in the regulation of B2F1 adherence. The eventual identification of gene(s) that encode STEC adhesins in LEE-negative strains will facilitate our understanding of the mechanism by which H-NS regulates adherence.

References

- [1] Bokete TN, Whittam TS, Wilson RA, Clausen CR, O'Callahan CM, Moseley SL, Fritsche TR, Tarr PI. Genetic and phenotypic analysis of *Escherichia coli* with enteropathogenic characteristics isolated from Seattle children. *J Infect Dis* 1997;175:1382–9.
- [2] Bonnet H, Souweine B, Gauthier G, Rich C, Livrelli V, Sirot J, Joly B, Forestier C. Non-O157:H7 Stx2-producing *Escherichia coli* strains associated with sporadic cases of hemolytic-uremic syndrome in adults. *J Clin Microbiol* 1998;36:1777–80.
- [3] Griffin PM. *Escherichia coli* O157:H7 and other enterohemorrhagic *Escherichia coli*. In: Blaser MJ, Smith PD, Ravdin JI, Greenberg HB, Guerrant RL, editors. *Infections of the gastrointestinal tract*. New York NY: Raven Press Ltd.; 1995. p. 739–61.
- [4] Willshaw GA, Scotland SM, Smith HR, Cheasty T, Thomas A, Rowe B. Hybridization of strains of *Escherichia coli* O157 with probes derived from the *eaeA* gene of enteropathogenic *E. coli* and the *eaeA* homolog from a vero cytotoxin-producing strain of *E. coli* O157. *J Clin Microbiol* 1994;32:897–902.
- [5] Schmidt H, Geitz C, Tarr PI, Frosch M, Karch H. Non-O157:H7 pathogenic Shiga toxin-producing *Escherichia coli*: phenotypic and genetic profiling of virulence traits and evidence for clonality. *J Infect Dis* 1999;179:115–23.
- [6] Whittam TS. Evolution of *Escherichia coli* O157:H7 and other Shiga toxin-producing *E. coli* strains. In: Kaper JB, O'Brien AD, editors. *Escherichia coli* O157:H7 and other Shiga toxin-producing *E. coli* strains. Washington, DC: ASM Press; 1998. p. 195–209.
- [7] McDaniel TK, Jarvis KG, Donnenberg MS, Kaper JB. A genetic locus of enterocyte effacement conserved among diverse enterobacterial pathogens. *Proc Natl Acad Sci USA* 1995;92:1664–8.
- [8] Ito H, Terai A, Kurazono H, Takeda Y, Nishibuchi M. Cloning and nucleotide sequencing of Vero toxin 2 variant genes from *Escherichia coli* O91:H21 isolated from a patient with the hemolytic uremic syndrome. *Microb Pathog* 1990;8:47–60.
- [9] Scott M. Mutations in the histone-like nucleoid structuring regulatory gene (*hns*) decrease the adherence of Shiga toxin-producing *Escherichia coli* O91:H21 strain B2F1 to human colonic epithelial

- cells and increase the production of hemolysin. Ph.D. Dissertation. Uniformed Services University of the Health Sciences, Bethesda, MD. 1999; p. 66–69.
- [10] Dytoc MT, Ismaili A, Philpott DJ, Soni R, Bruton JL, Sherman PM. Distinct binding properties of *eaeA*-negative verocytotoxin-producing *Escherichia coli* O157:H7 of serotype O113:H21. *Infect Immun* 1994; 62:3494–505.
 - [11] Paton AW, Srimanote P, Woodrow MC, Paton JC. Characterization of Saa, a novel autoagglutinating adhesin produced by locus of enterocyte effacement-negative Shiga-toxigenic *Escherichia coli* strains that are virulent for humans. *Infect Immun* 2001;69: 6999–7009.
 - [12] Srimanote P, Paton AW, Paton JC. Characterization of a novel type IV pilus locus encoded on the large plasmid of locus of enterocyte effacement-negative Shiga-toxigenic *Escherichia coli* strains that are virulent for humans. *Infect Immun* 2002;70:3094–100.
 - [13] Melton-Celsa AR, O'Brien AD. Activation of Shiga-like toxins by mouse and human intestinal mucus correlates with virulence of enterohemorrhagic *Escherichia coli* 091:H21 isolates in orally infected, streptomycin-treated mice. *Infect Immun* 1996;64:1569–76.
 - [14] Philpott DJ, Ackerley CA, Kiliaan AJ, Karmali MA, Perdue MH, Sherman PM. Translocation of verotoxin-I across T84 monolayers: mechanism of bacterial toxin penetration of epithelium. *Am J Physiol* 1997;273:G1349–58.
 - [15] Dharmasathaphorn K, Madara JL. Established intestinal cell lines as model systems for electrolyte transport studies. *Methods Enzymol* 1990;192:354–89.
 - [16] De Lorenzo V, Herrero M, Jakubzik U, Timmis KN. Mini-Tn5 transposon derivatives for insertion mutagenesis, promoter probing, and chromosomal insertion of cloned DNA in Gram-negative eubacteria. *J Bacteriol* 1990;172:6568–72.
 - [17] Cravioto A, Gross RJ, Scotland SM, Rowe B. An adhesive factor found in strains of *E. coli* belonging to the traditional infantile enteropathogenic serotypes. *Curr Microbiol* 1979;3:95–9.
 - [18] Winsor Jr. DK, Ashkenazi S, Chiovetti R, Cleary TG. Adherence of enterohemorrhagic *Escherichia coli* strains to a human colonic epithelial cell line (T84). *Infect Immun* 1992;60:1613–7.
 - [19] Maniatis T, Fritsch EF, Sambrook J. Molecular cloning: a laboratory manual. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory; 1989.
 - [20] Woolwine-Ciacchi F, Blomfield IC, Richardson SH, Mizel SB. Salmonella flagellin induces tumor necrosis factor alpha in a human promonocytic cell line. *Infect Immun* 1998;66:1127–34.
 - [21] Atlung T, Ingmer H. H-NS: a modulator of environmentally regulated gene expression. *Mol Microbiol* 1997;24:7–17.
 - [22] Dorman CJ, Hinton JC, Free A. Domain organization and oligomerization among H-NS-like nucleoid-associated proteins in bacteria. *Trends Microbiol* 1999;7:124–8.
 - [23] Schmidt H, Beutin L, Karch H. Molecular analysis of the plasmid-encoded hemolysin of *Escherichia coli* O157:H7 strain EDL 933. *Infect Immun* 1995;63:1055–61.
 - [24] Bauer M, Welch RA. Characterization of an RTX toxin from enterohemorrhagic *Escherichia coli* O157:H7. *Infect Immun* 1996;64: 167–75.
 - [25] Madrid C, Neito JM, Paytubi S, Falconi M, Gualerzi CO, Juarez A. Temperature- and H-NS-dependent regulation of a plasmid-encoded virulence operon expressing *Escherichia coli* hemolysin. *J Bacteriol* 2002;184:5058–66.
 - [26] Levine MM, Xu JG, Kaper JB, Lior H, Prado V, Tall B, Nataro J, Karch H, Wachsmuth K. A DNA probe to identify enterohemorrhagic *Escherichia coli* of O157:H7 and other serotypes that cause hemorrhagic colitis and hemolytic uremic syndrome. *J Infect Dis* 1987;156:157–82.
 - [27] Taylor RK, Miller VL, Furlong DB, Mekalanos JJ. Use of *phoA* gene fusions to identify a pilus colonization factor coordinately regulated with cholera toxin. *Proc Natl Acad Sci USA* 1987;84:2833–7.
 - [28] Teel L, Melton-Celsa AR, Schmitt CK, O'Brien AD. One of two copies of the gene for the activatable Shiga toxin type 2d (Stx2d) in *Escherichia coli* 091:H21 strain B2F1 is associated with an inducible bacteriophage. *Infect Immun* 2002;70:4282–91.
 - [29] Tanaka K, Muramatsu S, Yamada H, Mizuno T. Systematic characterization of curved DNA segments randomly cloned from *Escherichia coli* and their functional significance. *Mol Gen Genet* 1991;226:367–76.
 - [30] Elliott SJ, Sperandio V, Girón JA, Shin S, Mellies JL, Wainwright L, Hutcheson SW, McDaniel TK, Kaper JB. The locus of enterocyte effacement (LEE)-encoded regulator controls expression of both LEE- and non-LEE-encoded virulence factors in enteropathogenic and enterohemorrhagic *Escherichia coli*. *Infect Immun* 2000;68: 6115–26.
 - [31] Bustamante VH, Santana FJ, Calva E, Puente JL. Transcriptional regulation of type III secretion genes in enteropathogenic *Escherichia coli*: Ler antagonizes H-NS-dependent repression. *Mol Micro* 2001; 39:664–78.
 - [32] Beltrametti F, Kresse AU, Guzman CA. Transcriptional regulation of the *esp* genes of enterohemorrhagic *Escherichia coli*. *J Bacteriol* 1999;181:3409–18.