

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Manure Matters (newsletter)

Biological Systems Engineering

2003

Manure Matters, Volume 9, Number 5

Bahman Eghball
USDA-ARS, Lincoln, NE

Christopher Bauer
USDA-ARS, Lincoln, NE

Charles A. Shapiro
University of Nebraska-Lincoln, cshapiro1@unl.edu

Follow this and additional works at: <https://digitalcommons.unl.edu/manurematters>



Part of the [Agronomy and Crop Sciences Commons](#)

Eghball, Bahman; Bauer, Christopher; and Shapiro, Charles A., "Manure Matters, Volume 9, Number 5" (2003). *Manure Matters (newsletter)*. 33.

<https://digitalcommons.unl.edu/manurematters/33>

This Article is brought to you for free and open access by the Biological Systems Engineering at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Manure Matters (newsletter) by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

UNL's Livestock
Environmental
Issues Committee
Includes
representation from
UNL, Nebraska
Department of
Environmental
Quality, Natural
Resources
Conservation Service,
Natural Resources
Districts, Center for
Rural Affairs,
Nebraska Cattlemen,
USDA Ag Research
Services, and
Nebraska Pork
Producers
Association.

Manure Matters

Volume 9, Number 5

Reducing Spatial Variability of Soil Carbon and Phosphorus by Site-Specific Manure Application

Bahman Eghball and Christopher Bauer, USDA-ARS, Lincoln, NE
and Charles A. Shapiro, Univ. of Nebraska, Haskell Ag Laboratory, Concord, NE

Spatial variability can significantly influence crop performance across a field. Manure, a renewable resource, is an excellent source of nutrients that can be substituted for synthetic types of fertilizers. The carbon (C) and nutrients in manure can enhance the physical and chemical properties of soils, especially infertile soils, hence reducing soil spatial variability. Organic C constitute about 58% of organic matter in the soil ($\%OM = \%OC \times 1.724$). Manure application not only provides nutrients for crops but also improves soil quality since the organic matter in manure improves soil physical and chemical properties. Manure also contains lime that will increase soil pH in areas with acidic soil pH. Therefore one would expect more positive effects of manure application on crop yield than chemical fertilizer application. The objective of this study was to evaluate the effects of C and nutrients in manure on soil C and P variability.

The experiment was conducted from 1998 to 2000 on a private farm in Hamilton

County near Phillips, Nebraska on a center pivot irrigated field with continuous corn. A ridge-till system, consisting of stalk chopping, cultivation, and ridging was utilized. The experimental area consisted of Hord silt loam, Invale loamy sand, Thurman fine sandy loam, Ortello fine sandy loam, Uly silt loam, and Alda loam soils. Treatments included strips (40 ft wide and 2200 ft long) of site-specific manure, uniform manure, uniform commercial fertilizer, and a no-treatment check. Each year, organic C levels were determined by grid sampling of the area, with two samples taken, 20 to 30 inches apart, to a depth of 6 inches at each grid point. Samples were collected every 80 ft in the middle of each strip. Each point was geo-referenced using a differential global positioning system (DGPS). In the site-specific manure treatment, manure was applied to areas within the strips that had a soil C concentration $< 1.4\%$ (2.4% organic matter). Fertilizer was applied to areas within site-specific strips that did not receive manure. Manure was applied to provide 170 lb N/acre. The cooperators

**UNL's Livestock
Environmental
Issues Committee**
Includes
representation from
UNL, Nebraska
Department of
Environmental
Quality, Natural
Resources
Conservation Service,
Natural Resources
Districts, Center for
Rural Affairs,
Nebraska Cattlemen,
USDA Ag Research
Services, and
Nebraska Pork
Producers
Association.

Contact:
Chris Henry
217 LW Chase Hall
University of NE
Lincoln, NE 68583
(402) 472-6529
chenry@unl.edu

applied anhydrous ammonia at a rate of 190 lb N/acre for the fertilizer treatment.

Soil organic C in the soil was increased from 1.17% to 1.52% (2.01 to 2.62% organic matter) with site-specific manure application (Fig. 1). The increase was significant when compared to the original soil C level in 1997 or synthetic fertilizer application. There were about 2.8 tons per acre C sequestered in the soil when site-specific manure was applied. Soil C coefficient of variation (CV), an indicator of variability, was reduced from 29.3% to 22.7% by site-specific manure application. About 50% of the field had soil C concentrations < 1.1% in 1997 (Fig. 1). However, by 2000 when three manure applications had been made, the areas with soil C concentrations < 1.1% were located in a small pocket (< 1% of the field) in the north part of the field. The highest organic C level was 1.847% in 1997 and 2.177% in 2000. This indicates that site-specific manure application not only increased soil C level but also reduced variability of soil organic C and therefore of organic matter.

There was a significant correlation between soil C levels and plant-available P concentrations. Therefore, the low soil C areas also had low P levels (Fig. 1 and 2). Site-specific manure application increased soil P level from a field average of 23 ppm to 61 ppm (Fig. 2) and most of the increase was in the low P areas. Even though the field P average before manure application in 1997 was 23 ppm, about 53% of the field had soil P levels that were less than 15 ppm. A soil Bray and Kurtz P level < 15 ppm is considered deficient for corn and P application will be recommended by the University of Nebraska fertilizer recommendation. As is indicated in Fig. 2, the P deficient areas (< 15 ppm) were located in a small area (about 10% of the field) in the south central part of the field after only two years of site-specific manure

application. The CV of P distribution in the field was reduced from 86% to 42%, a reduction of 51%. The soil P increase in the deficient areas caused a significant grain yield increase for site-specific manure application as compared to uniform fertilizer application. Averaged across five years (1998 to 2002), corn grain yield was 180 bu/acre for site-specific manure, 169 bu/acre for uniform manure, 172 bu/acre for chemical fertilizer, and 116 bu/acre for the untreated check.

Site-specific manure application strategy is not only an effective method of manure utilization but also can significantly reduce soil spatial variability and hence increase crop yield and improve soil quality. By site-specific manure application less productive sites within a field are improved and the productive sites would not receive excess nutrients, especially P, that can cause water quality concerns.

UNL's Livestock Environmental Issues Committee Includes representation from UNL, Nebraska Department of Environmental Quality, Natural Resources Conservation Service, Natural Resources Districts, Center for Rural Affairs, Nebraska Cattlemen, USDA Ag Research Services, and Nebraska Pork Producers Association.

Contact:
Chris Henry
217 LW Chase Hall
University of NE
Lincoln, NE 68583
(402) 472-6529
chenry@unl.edu




Figure 1. Soil organic C before (1997) and after 3 years (2000) of site-specific manure application. The light color areas represent C concentration < 1.1%




Figure 2. Soil P level before (1997) and after 2 years (1999) of site-specific manure application.

Issued in furtherance of Cooperative Extension work, Acts of May 8 and June 30, 1914, in cooperative with the U.S. Department of Agriculture. Elbert Dickey, Director of Cooperative Extension, University of Nebraska, Institute of Agriculture and Natural Resources. University of Nebraska Cooperative Extension educational programs abide with the non-discrimination policies of the University of Nebraska