

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Papers in Herpetology

Papers in the Biological Sciences

2007

Geographic Distribution: *Ramphotyphlops braminus* (Brahminy Blindsnake). USA: Florida: Volusia Co

Louis A. Somma

FDAC, louis.somma@fdacs.gov

Paul E. Skelley

FDAC

Follow this and additional works at: <https://digitalcommons.unl.edu/biosciherpetology>



Part of the [Other Ecology and Evolutionary Biology Commons](#), and the [Zoology Commons](#)

Somma, Louis A. and Skelley, Paul E., "Geographic Distribution: *Ramphotyphlops braminus* (Brahminy Blindsnake). USA: Florida: Volusia Co" (2007). *Papers in Herpetology*. 24.

<https://digitalcommons.unl.edu/biosciherpetology/24>

This Article is brought to you for free and open access by the Papers in the Biological Sciences at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Papers in Herpetology by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

RAMPHOTYPHLOPS BRAMINUS (Brahminy Blindsnake). USA: FLORIDA: VOLUSIA Co.: New Smyrna Beach. 23 September 2006. Collected by Stacy Simmons. Verified by Kenneth L. Krysko, Florida Museum of Natural History, University of Florida. One individual found on floor of residence. UF 151313. First county record. Originally turned in to Division of Plant Industry (DPI), Florida Department of Agriculture and Consumer Services, Gainesville, Florida, as a "worm for ID" on 27 September 2006; Entomology Log No. 6879. It is not clear if this single specimen indicates an established breeding colony in this county. However, this invasive, parthenogenic exotic is established in other Florida counties and other regions of the world through its ability to disperse in the soil of transported potted plants (Ernst and Ernst 2003. Snakes of the United States and Canada. Smithsonian Books, Washington and London; Meshaka et al. 2004. The Exotic Amphibians and Reptiles of Florida. Krieger Publishing Company, Malabar, Florida).

Submitted by LOUIS A. SOMMA, Department of Zoology, PO Box 118525, 223 Bartram Hall, University of Florida, Gainesville, Florida 32611-8525, USA (e-mail: las@zoo.ufl.edu); and PAULE E. SKELLEY, Florida State Collection of Arthropods, Division of Plant Industry – Entomology Section, PO Box 147100, Florida Department of Agriculture and Consumer Services, 1911 SW 34th St., Gainesville, Florida, 32614-7100, USA (e-mail: skellep@doacs.state.fl.us).