University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Faculty Publications from the Center for Plant
Science Innovation

Plant Science Innovation, Center for

11-9-2007

Comment on "A G Protein–Coupled Receptor Is a Plasma Membrane Receptor for the Plant Hormone Absciscic Acid"

Christopher Johnston

University of North Carolina, Chapel Hill

Brenda Temple

University of North Carolina, Chapel Hill

Jin-Gui Chen

University of British Columbia, Vancouver, BC V6T 1Z4, Canada

Yajun Gao

University of British Columbia, Vancouver, BC V6T 1Z4, Canada

Etsuko N. Moriyama

University of Nebraska - Lincoln, emoriyama2@unl.edu

See next page for additional authors

Follow this and additional works at: <https://digitalcommons.unl.edu/plantscifacpub>

Part of the [Plant Sciences Commons](#)

Johnston, Christopher; Temple, Brenda; Chen, Jin-Gui; Gao, Yajun; Moriyama, Etsuko N.; Jones, Alan M.; Siderovski, David; and Willard, Francis, "Comment on "A G Protein–Coupled Receptor Is a Plasma Membrane Receptor for the Plant Hormone Absciscic Acid"" (2007). *Faculty Publications from the Center for Plant Science Innovation*. 30.

<https://digitalcommons.unl.edu/plantscifacpub/30>

This Article is brought to you for free and open access by the Plant Science Innovation, Center for at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Faculty Publications from the Center for Plant Science Innovation by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors

Christopher Johnston, Brenda Temple, Jin-Gui Chen, Yajun Gao, Etsuko N. Moriyama, Alan M. Jones, David Siderovski, and Francis Willard

Figure 2. GCR2 has a predicted tertiary structure consistent with a LanC protein. BLASTP search against the structural database (<http://www.rcsb.org>), using GCR2 as the query, identified nisin cyclase (NisC, PDB ID: 2G0D) as the only structural homolog producing a statistically significant alignment [expect (E) value, 4×10^{-4}]. Significant sequence similarity was noted between amino acids 216 and 282 of GCR2 and amino acids 209 and 386 of NisC. A homology model of GCR2 (amino acids 216 to 282) was then generated using Insight-II (<http://www.accelrys.com/products/insight>). Shown is a superposition of the GCR2 homology model (blue) and the corresponding region of NisC (green). The N and C termini are labeled accordingly. Alpha helices observed in the NisC structure are denoted H8 to H14. Arrows indicate two segments in which NisC contains extended inserts relative to GCR2 and are the only areas of the superposition that diverge between the molecules. The proposed catalytic residues are indicated in NisC (yellow sticks) and GCR2 (red sticks). The superposition and image were generated using PyMol (DeLano Scientific, Palo Alto, CA, USA).

indicates that GCR2 is most likely an α - α toroid protein. This fold is a defining structural characteristic of LanC proteins, terpenoid cyclases, glycosidases, and farnesyl transferases (5, 6).

We created a homology model of GCR2 based on the crystal structure of the *Lactococcus lactis* LanC protein, nisin cyclase (NisC) (Figure 2) (6). This homology model has a Profiles-3D self-compatibility score of 27.6%, indicating a valid model with robust statistical confidence (7). The core α -helices of NisC superimpose very well to those of the GCR2 homology model (Figure 2), with an overall root mean square deviation of 4.0 Å. The zinc-coordinating residues of NisC, important for cysteine cyclization, are conserved in the primary sequence (Figure 1A), and these residues in NisC superimpose well with corresponding residues

Figure 3. Simulated surface plasmon resonance binding curves for a 2 nM affinity interaction between GPA1 and GCR2. Simulations of GPA1 binding to immobilized GCR2, using the rate constants published by Liu *et al.* ($k_a = 1.77 \times 10^4 \text{ M}^{-1}\text{s}^{-1}$; $k_d = 3.9 \times 10^{-5} \text{ s}^{-1}$). Simulated injections are plotted for four different concentrations of GPA1 as reported by Liu *et al.* (1). Arrow indicates the injection time course and corresponding association phase. Simulated sensorgrams were generated using BIAeval 3.2 software (GE Healthcare, Uppsala, Sweden), using the 1:1 Langmuir model with maximum binding of 100 RU. (A) Simulated sensorgrams for an interaction that has no bulk buffer shift. (B) Simulated sensorgrams for an interaction occurring with a bulk buffer shift of 100 RU.

of the GCR2 model (Figure 2). This analysis provides a structural argument that GCR2 is a member of the LanC protein superfamily, not the GPCR superfamily.

Notably, a mammalian LanC homolog (LANCL1) was originally misidentified as a GPCR (GPR69A/p40) (8). In subsequent studies, the authors determined that GPR69A was in fact a LanC ortholog and renamed this protein LANCL1 (9). Biochemical studies confirmed LANCL1 to be a peripheral membrane protein (9). Subsequently, the related protein LANCL2 was suggested to be membrane localized due to both myristoylation and lipid binding (10). These data from orthologous proteins suggest that GCR2 is likely to be a peripheral membrane protein. Further evidence against GCR2 having a 7TM topology is provided by the split ubiquitin assays of Liu *et al.* showing that GPA1-Cub interacts equally well with both N- and C-terminal fusions of GCR2 to NubG (1). These results are incompatible with the GCR2 N terminus being extracellular, as is the case with all known GPCRs, and are incompatible with GCR2 having an odd number of transmembrane spans.

Liu *et al.* reported solubilizing recombinant GCR2 from *Escherichia coli* using 0.1% Triton-X100 and purifying GCR2 to homogeneity. The apparent ease of this purification and the methods used are generally contrary to the known arduous biochemistry of GPCR purification, given 7TM helices (11), but are entirely consistent with purifying a soluble cytosolic protein from *E. coli*. *In vitro* protein-protein interaction was reported between GCR2 and the *Arabidopsis* Ga subunit GPA1 using surface plasmon resonance (SPR) (1). However, the presented SPR data are not representative of a *bona fide* interaction (12). Indeed, the data clearly demonstrate an absence of any GCR2/GPA1 interaction, as GPA1 binding to GCR2 is equivalent to that of the negative control BSA [figure S3 in (1)]. The presented sensorgrams are most likely bulk shift artifacts normally corrected by negative control subtraction (12). We were unable to determine how Liu *et al.* (1) measured their rate constants. However, simulated SPR sensorgrams based on their reported values (Figure 3) clearly demonstrate a discrepancy between the data of Liu *et al.* (1) and expected SPR results (12) based on their reported rate constants. The reported off-rate constant ($3.9 \times 10^{-5} \text{ s}^{-1}$) suggests that the GCR2/GPA1 complex has a binding half-life of 5 hours, thoroughly inconsistent with the raw data presented by Liu *et al.* (1), and also suggesting that a surface regeneration step would be necessary to obtain reliable dose-response data.

The classical *in vitro* assay for GPCR/Ga coupling is demonstration of agonist-promoted guanine nucleotide exchange factor activity, either by GTP γ S binding or steady-state GTPase activity (13). Reconstituting interactions between Ga subunits and their cognate GPCRs typically requires lipid-modified Ga and G $\beta\gamma$ subunits and a model membrane (13). The binding of Ga and G $\beta\gamma$ to GPCRs is synergistic, whereas isolated subunits have low affinity for receptor (14). These considerations were not addressed by Liu *et al.* (1). In summary, while it is possible that GCR2 is both an intracellular receptor for abscisic acid and a G protein modulator, we conclude that GCR2 is neither a transmembrane protein nor a G protein coupled receptor, but rather is an *Arabidopsis* homolog of bacterial lanthionine synthetases. We recommend that any putative plant GPCR be rigorously characterized as a *bona fide* G protein coupled receptor using *in vitro* biochemical methods for demonstrating G protein coupling and activation that have been well-established for the analysis of mammalian GPCRs.

References and Notes

1. X. Liu *et al.*, *Science* **315**, 1712 (2007).
2. S. Moller, M. D. Croning, R. Apweiler, *Bioinformatics* **17**, 646 (2001).
3. E. N. Moriyama, P. K. Strope, S. O. Opiyo, Z. Chen, A. M. Jones, *Genome Biol.* **7**, R96 (2006).
4. S. F. Altschul, W. Gish, W. Miller, E. W. Myers, D. J. Lipman, *J. Mol. Biol.* **215**, 403 (1990).
5. D. W. Christianson, *Science* **311**, 1382 (2006).
6. B. Li *et al.*, *Science* **311**, 1464 (2006).
7. R. Luthy, J. U. Bowie, D. Eisenberg, *Nature* **356**, 83 (1992).
8. H. Mayer, J. Breuss, S. Zeigler, R. Prohaska, *Biochim. Biophys. Acta* **1399**, 51 (1999).
9. H. Bauer, H. Mayer, A. Marchler-Bauer, U. Salzer, R. Prohaska, *Biochem. Biophys. Res. Commun.* **275**, 69 (2000).
10. C. Landlinger, U. Salzer, R. Prohaska, *Biochim. Biophys. Acta* **1758**, 1759 (2006).
11. V. Sarramegn, I. Muller, A. Milon, F. Talmont, *Cell. Mol. Life Sci.* **63**, 1149 (2006).
12. R. L. Rich, D. G. Myszka, *J. Mol. Recognit.* **19**, 478 (2006).
13. E. M. Parker, K. Kameyama, T. Higashijima, E. M. Ross, *J. Biol. Chem.* **266**, 519 (1991).
14. W. A. Clark, X. Jian, L. Chen, J. K. Northup, *Biochem. J.* **358**, 389 (2001).
15. We thank S. K. Nair (University of Illinois at Urbana-Champaign) for valuable discussions.