


1915

## A New Sabertoothed Cat from Nebraska

Erwin H. Barbour

Harold J. Cook

Follow this and additional works at: <http://digitalcommons.unl.edu/conservationsurvey>

 Part of the [Geology Commons](#), [Geomorphology Commons](#), [Hydrology Commons](#), [Paleontology Commons](#), [Sedimentology Commons](#), [Soil Science Commons](#), and the [Stratigraphy Commons](#)

---

Barbour, Erwin H. and Cook, Harold J., "A New Sabertoothed Cat from Nebraska" (1915). *Conservation and Survey Division*. 31.  
<http://digitalcommons.unl.edu/conservationsurvey/31>

This Article is brought to you for free and open access by the Natural Resources, School of at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Conservation and Survey Division by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

36c

# NEBRASKA GEOLOGICAL SURVEY

ERWIN HINCKLEY BARBOUR, State Geologist

VOLUME 4

PART 17

A NEW SABER-TOOTHED CAT FROM NEBRASKA

BY

ERWIN H. BARBOUR

AND

HAROLD J. COOK

---

GEOLOGICAL COLLECTIONS OF HON. CHARLES H. MORRILL



208 B

## A NEW SABER-TOOTHED CAT FROM NEBRASKA

BY ERWIN H. BARBOUR AND HAROLD J. COOK

During the field season of 1913, while exploring the Pliocene beds of Brown County, Mr. A. C. Whitford, a Fellow in the Department of Geology, University of Nebraska, discovered the mandible of a new machærodont cat. His work in this region was in the interest of the Nebraska Geological Survey and the Morrill Geological Expeditions.<sup>1</sup>

The known fossil remains of the ancestral Felidæ fall into two natural lines of descent, as pointed out by Dr. W. D. Matthew.<sup>2</sup> Beginning in the Oligocene, we find two genera as contemporary forms,




Fig. 1.—a, *Dinictis squalidens*, Middle Oligocene.  $\times \frac{1}{4}$ . b, *Hoplophoneus primaevus*, Middle Oligocene.  $\times \frac{1}{4}$ . After Matthew.

*Hoplophoneus* and *Dinictis*, which are of immediate interest. *Dinictis* gave rise to the true cats, which have survived to the present time, passing through the genera *Nimravus*, *Pseudælorus*, and *Felis*, in the order named. *Hoplophoneus* gave rise to the "saber-toothed" cats, so called because they developed two immense canine teeth in the upper jaws. *Hoplophoneus* was succeeded by *Machærodus*, and this in turn by *Smilodon*. In the last named, the phylum culminated in relatively

1. A preliminary account of the region and its fossils may be found in the *University Studies*, vol. 14, No. 2, 1914, and in brochures 32 and 33 of the Nebraska Geological Survey.

2. Matthew, W. D., *Phylogeny of the Felidæ*, *Bull. Am. Mus. Nat. Hist.*, vol. 28, Art. 26, 1910.

recent time. It was contemporaneous with primitive man. A third genus, *Eusmilus*, is also present in the Oligocene, but belongs to still another group paralleling *Smilodon*.

The typical *Machærodus* comes from Europe, and it is doubtful whether this genus really occurs in America. Forms very like it do occur, but owing to the fragmentary condition of much of the known material from the later Tertiary formations of this country, our knowledge of them is incomplete, and better specimens will probably show the American "*Machærodus*" to be distinct from the European form. The later *Machærodontidae* stand in great need of revision. The new species herein described falls most readily under the definition of *Eusmilus*, and will be treated as such for the purposes of this paper.




Fig. 2.—a, *Nimravus gomphodus*, Upper Oligocene.  $\times \frac{1}{4}$ . b, *Machærodus palmidens*, Middle Miocene of France.  $\times \frac{1}{4}$ . After Matthew.

However, it shows affinities to *Machærodus*. As a revision of the literature is out of the province of this paper, a discussion of these relationships will not be attempted.

For this new species of *Eusmilus*, the specific name, *whitfordi*, is proposed in recognition of the finder, Mr. A. C. Whitford.

Subfamily—*Machærodontinae*

Genus —*Eusmilus*

Species —*whitfordi*, sp. nov.

?

Dentition—                      
2—1—2—1

Type, right lower jaw, No. 7-2-4-13 complete with the exception of  $P_3$ , and the crowns of the incisors and canine.

The jaw is about the size of *Felis concolor*, but of very different proportions. The symphyseal flange is very deep, and indicates a long, slender canine. The symphyseal attachment is narrow, and extends far down the flange, almost vertically. The coronoid process is extremely reduced, the top being on a level with the crowns of the cusps of  $P_4$  and  $M_1$ . It is relatively small and more reduced than in *Smilodon*. The angle of the jaw is strongly developed. The masseteric




Fig. 3.—a, *Felis concolor*, "Rocky Mountain Lion," Puma.  $\times \frac{1}{4}$ . b, *Smilodon californicus*, Pleistocene.  $\times \frac{1}{4}$ . After Matthew.

fossa is deep and rough. The diastema is proportionally short between  $P_3$  and the canine.  $P_3$  is represented by the alveolus, and was evidently vestigial.  $P_4$  is well developed, with three principal cusps. The notch between the paraconid and protoconid is shallow, and there may have been a rudimentary posterior cusp. If present, it was very small. Both cutting teeth are well worn down, so that it is impossible to tell to what extent the cusps are developed. The animal was less than half the size of the Pleistocene *Smilodon californicus*, found in the famous Rancho la Brea beds at Los Angeles.

The University of Nebraska  
Lincoln, September 15, 1914.

Distributed January 18, 1915.




Fig. 4.—Saber-toothed tiger, *Smilodon neogaeus*. Lower Pleistocene. Modified after Osborn and Knight.



*EUSMILUS WHITFORDI*, SP. NOV.

- a. Mandible, outer surface.
  - b. Same, inner surface.
  - c. Same, crown view.
- Incisors restored.